



Thermal Time Approach under Different Sowing Windows for Greengram

**K. Sabaridheepa^a, S. P. Ramanathan^{a*}, S. Kokilavani^a, Ga Dheebakaran^a
and K. Boomiraj^a**

^a *Agro Climate Research Centre, Tamil Nadu Agricultural University, Coimbatore-641 003, India.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i1131209

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/91613>

Original Research Article

Received 29 August 2022
Accepted 05 September 2022
Published 07 September 2022

ABSTRACT

Crop phenology and yield are highly influenced by prevailing weather condition during the crop growth phases. A research work was carried out at Agro Climate Research Centre, Tamil Nadu Agricultural University, Coimbatore during June to August 2022 to determine the suitable sowing window for Greengram. The field experiment was laid out in CO 8 Greengram variety with two sowing windows of D₁ - 15 June 2022 (24th SMW) and D₂ - 22 June 2022 (25th SMW). The best time of sowing was established by using Growing Degree Days (GDD), Helio thermal Units (HTU), Photo thermal Units (PTU) and Heat Use Efficiency (HUE). The study revealed that the 25th Standard Meteorological week (SMW) recorded higher yield due to higher values of GDD, HTU, PTU and HUE at different crop growth stages.

Keywords: *Greengram; sowing dates; GDD; HTU; PTU and HUE.*

1. INTRODUCTION

Greengram (*Vigna radiata* L.) commonly known as 'Mung bean' has become the third most important pulse crop next to the Bengal gram and Pigeon pea. It is native of India and Central Asia,

a region where it has been grown since prehistoric times. The green gram seeds are extremely nutritious, which is almost 2.5–3.0 times higher in protein content than cereals [1]. In India, Mungbean was grown on an area of about 5.13 million hectares with the production of about

*Corresponding author: E-mail: ramanathan.sp@tnau.ac.in;

3 million tonnes and productivity of 601 kg/ha [2], Green gram was considered as “nutritional powerhouse” due to high protein content, among many pulses [3]. It is a leguminous crop and helps in enhancing the soil fertility by absorbing atmospheric nitrogen through root nodules. Rhizobium bacteria form nodules on the root of green gram and fix atmospheric nitrogen. Through nitrogen fixation, the green gram can increase soil fertility [4]. It supplies an excellent green fodder to the animals and also used as green manure crop.

Agriculture being backbone of economy in the country and mainly dependent on climatic factors that determines the success or failure of crop productivity. Crop Phenology and yield are highly influenced by prevailing weather condition during the crop growth phase. A change in climate had registered a deleterious effect on crop growth, development and production during the past [5]. Interception of photosynthetically active radiation, air and leaf temperature, relative humidity, prevailing wind speed, CO₂ concentration and soil moisture availability are all environmental factors which might have a major impact on crop growth during growing season [6,7].

Temperature is a determinable weather parameter which plays a pivotal role, that directly have consequences on agro based ecosystem. Plants that grows above a temperature range are characterised either by lower and upper temperature thresholds as its growth thresholds. Under certain limits, temperature forms the linear relationships in crop development [8]. As temperature exceeds to the species optimum level, the growth rate of vegetative phase (leaf appearance) tends to increase. Most plant species have a higher optimum temperature for vegetative stage than for reproductive stage. The thermal environment seems to have an influence on the growth of all crop systems. Sowing window has been recognized as the most important non-monetary input factor because it ensures complete harmony between the vegetative and reproductive phases on the one side, and the climatic rhythm on the other [9]. It is precious and important for better and efficient utilization of available plant growth. Sowing time represents the most important factor that influences mung bean yield among numerous agronomic practises [10].

Many investigators used temperature based agrometeorological indices for evaluating plant growth and development viz., Growing Degree Days (GDD), Photothermal Units (PTU), Helio-

thermal Units (HTU) and Heat Use Efficiency (HUE) [11,12,13]. GDDs is a model that converts raw climatic data into make crop management decisions [14]. Temperature affects many physiological processes, including respiration, photosynthesis, germination, flowering, protein denaturation, and physical processes such as transpiration and evaporation. The heat unit concept assumes that plant growth and temperature have a direct and linear relationship. It begins with the assumption that plant growth is determined by the 'total amount of heat' to which it is exposed during its life.

Present study was aimed to assess the growth and yield of green gram with heat units based agroclimatic models under various sowing windows.

2. MATERIALS AND METHODS

The research trial was conducted during the *kharif* (June - August) season of 2022 at Wetland Farm, Tamil Nadu Agricultural University, Coimbatore and the study area was situated in Western Agro-climatic zone of Tamil Nadu at 11° N latitude and 77° E longitude with an altitude of 426.7 m above mean sea level. The soil of experimental field was clay in texture with a pH of 7.2. The field experiment was laid out in a Split Plot Design, replicated thrice with CO 8 Greengram variety of 55-60 days (short duration crop) by keeping two sowing windows of D₁ - 15 June 2022 (24th Standard Meteorological Week (SMW)) and D₂ - 22 June 2022 (25th SMW). The daily weather data like maximum, minimum temperature, bright sunshine hours required to compute the thermal time was obtained from Principal A Class Agro-meteorological Observatory, Coimbatore.

2.1 Thermal Based Agro Meteorological Indices

Growing Degree Days (GDD)

It is calculated using the following formula proposed by [6].

$$GDD = \frac{T_{max} + T_{min}}{2} - T_{base}$$

where,

T_{max} - daily maximum temperature (°C)
 T_{min} - daily minimum temperature (°C) and
 T_{base} - base temperature of 10 °C was used.

Helio Thermal Unit (HTU)

It is determined using the formula indicated by [15].

$$HTU = GDD \times \text{Actual Bright Sunshine hours } (^{\circ}\text{C hr})$$

Photo Thermal Unit (PTU)

Adding the day length element and employing the GDD idea as a photo-thermal unit in climatic analogy studies, [16] aimed to improve the GDD concept. GDD multiplied by the number of bright sunlight hours that can possibly be obtained (N).

$$PTU = GDD \times \text{Day Length (N)}$$

where day length (N) refers to maximum possible sunshine hours.

Heat use efficiency (HUE)

Net amount of dry matter generated per thermal unit, or growing degree day, is known as heat use efficiency [17]. The efficiency with which the available heat is utilised for grain yields has been characterised as yield per growing degree day or thermal time.

$$\text{Heat use efficiency (HUE)} = \frac{\text{Seed Biomass / Yield (kg ha}^{-1}\text{)}}{\text{GDD}}$$

The T-test was carried between the two mean of the different sowing dates at the level of 5 per cent significance to find the best sowing window for Green gram.

3. RESULTS AND DISCUSSION

3.1 Growing Degree Days (GDD) for Greengram

Data pertaining to thermal units of green gram at varied crop growth stages under different sowing window is presented in Table 1. It was observed that highest numbers of Growing Degree Days (GDD) were noticed on 24th SMW when compared to 25th SMW.

During flowering and pod initiation phase, highest thermal accumulation was obtained in 24th SMW (D₁ – 418^oC at flowering and 355^oC at pod initiation stage) followed by 25th SMW (D₂ – 411^oC at flowering and 333^oC at pod initiation stage. At physiological maturity phase, GDD accumulation recorded maximum (1745^oC) in 24th SMW followed by (162^oC) in 25th SMW. Similar result was reported by [18]. Consistently, a decreasing trend in HTU and PTU for attaining all Phenological stages was observed in 25th SMW.

Table 1. Growing degree days, helio thermal unit and photo thermal unit for various stages of greengram under different sowing dates

Crop stages	15 th June			22 nd June		
	GDD (°C)	HTU (°C)	PTU (°C)	GDD (°C)	HTU (°C)	PTU (°C)
Germination	108	794	1241	105	377	1206
Flower Initiation	418	1485	1252	411	1506	4765
Pod Initiation	355	1797	4186	333	1730	3926
Maturity	162	958	1916	175	1226	2063
Mean	260.8	1258.5	2148.8	256.0	1209.8	2990.0
SE±	149.1026	464.5231	1394.361	140.6129	592.2164	1639.99

D1-Date of sowing-15 June 2022, D2-22 June 2022, GDD-Growing Degree Days, HTU- Helio Thermal Unit, PTU- Photo Thermal Unit, SE± - Standard deviation

Table 2. Heat use efficiency under different sowing dates of Greengram

Crop stages	Yield (kg ha ⁻¹)	HUE (kg ha ⁻¹ °C)	DMP (kg ha ⁻¹)	HUE (kg ha ⁻¹ °C)
D1	871	0.83	2598	2.49
D2	809	0.77	2412	2.31
Mean	840	0.80	2505	2.40
SE±	43.84	0.04	131.52	0.13
T value	5.894		3.871	
P value	0.004*		0.018*	

D1-Date of sowing-15 June 2022, D2-22 June 2022, HUE-Heat use efficiency, DMP-Dry matter production, SE± - Standard deviation. *P<0.05 – 5% level of significance

3.2 Heat Use Efficiency (HUE) for Green Gram

Data recorded on heat use efficiency for dry matter (kg ha^{-1}) as well as grain yield are presented in Table 2. The 24th SMW recorded the highest HUE for grain yield ($0.83 \text{ kg ha}^{-1} \text{ }^{\circ}\text{C}$) followed by 25th SMW ($0.77 \text{ kg ha}^{-1} \text{ }^{\circ}\text{C}$). The higher HUE for dry matter ($2.49 \text{ kg ha}^{-1} \text{ }^{\circ}\text{C}$) was noticed on 24th SMW. This could be attributed by maximum consumption of heat units absorbed during vegetative growth phases. The lower HUE for dry matter ($2.31 \text{ kg ha}^{-1} \text{ }^{\circ}\text{C}$) was found on 25th SMW. This was due to lower accumulation of thermal units. The findings are consistent with those found by [19].

Since P value is less than 0.05 at 5 per cent level of significance for both grain yield and biomass yield, D1 is significant than the D2. A t-test was carried out for the yield and biomass of the Green gram crop and results indicated that the 24th SMW has given significantly higher grain yield and biomass than the 25th SMW. Sowing window significantly influenced the grain yield of green gram crop. Maximum grain yield was recorded in 24th SMW with a yield of 871 kg/ha. It might be due to efficient utilization of GDD, HTU and PTU. The crop grown on 25th SMW attained comparatively lesser yield of 809 kg/ha which is ascribed to continuousness of rains during ensuing weeks. Similar findings are also noticed by [20] and [19].

4. CONCLUSION

Study on heat units of green gram concluded that the sowing of green gram on 24th SMW showed better performance due to optimal thermal indices and higher use efficiency compared to sowing on 25th SMW. Sowing of Green gram during the 24th standard week produced good crop yield in Coimbatore.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kumar B, Dhaliwal SS, Singh ST, Lamba JS, Ram H. Herbage production, nutritional composition and quality of teosinte under Fe fertilization. *International Journal of Agriculture & Biology*. 2016;18(2).
2. Indiastat. Rice production in India; 2021. Accessed 21, July 2022. Available:<http://www.indiastat.com/>.
3. Singh U, Verma P, Praharaj CS, Srivastava RP, Parihar AK. Dietary Supplementation through pulses for nutritional security. *Indian Farming*. 2018; 66(8).
4. Ram A, Dev I, Kumar D, Uthappa AR, Tewari RK, Singh R, Sridhar KB, Singh M, Srivastava M, Kumar V, Chaturvedi OP. Effect of tillage and residue management practices on blackgram and greengram under bael (*Aegle marmelos* L.) based agroforestry system. *Indian Journal of Agroforestry*. 2016;18(1):90-5.
5. Swaminathan MS, Kesavan PC. Agricultural research in an era of climate change. *Agricultural Research*. 2012; 1(1):3-11.
6. Kaur A, Dhaliwal LK. Agroclimatic indices of rice (*Oryza sativa* L.) under different dates of planting. *Progressive Research*. 2014;9(1):222-7.
7. Makone P, Patel JG, Desai CK, Das S, Pal V, Paramar JK. Influence of weather parameters on summer greengram (*Vigna radiata* (L.) Wilczek) at Sardarkrushinagar. *Journal of Agrometeorology*. 2015; 17(1):142-4.
8. Steduto P, Hsiao TC, Fereres E, Raes D. Crop yield response to water. Rome: Food and Agriculture Organization of the United Nations; 2012. Accessed 11, July 2022. Available:<https://www.fao.org/3/i2800e/i2800e00.htm>
9. Subash N, Ram Mohan HS, Sikka AK. Decadal frequency and trends of extreme excess/deficit rainfall during the monsoon season over different meteorological subdivisions of India. *Hydrological Sciences Journal*. 2011;56(7):1090-109.
10. Fraz RA, Iqbal J, Bakhsh MA. Effect of sowing dates and planting patterns on growth and yield of mungbean (*Vigna radiata* L.) cv. M-6. *Int. J. Agri. Biol*. 2006;8(3):363-5.
11. Shamim MO, Devendra S, Gangwar KK, Vipin K. Agro meteorological indices in relation to phenology, biomass accumulation and yield of rice genotypes under Western Plain zone of Uttar Pradesh. *J. Agrometeorol*. 2013;15(2):50-7.
12. Abhilash DC, Singh R, Premdeep RS. Agrometeorological indices and phenology

- of Basmati Rice (*Oryza sativa* L.) under different dates of transplanting. International Journal of Current Microbiology and Applied Science. 2017; 6(3):212-22.
13. Chaudhari NV, Kumar N, Parmar PK, Dakhore KK, Chaudhari SN, Chandrawanshi SK. Thermal indices in relation to crop phenology and yield of rice (*Oryza sativa* L.) grown in the south Gujarat region. J. Pharmacogn. Phytochem. 2019;8(2):146-9.
 14. Pathak TB, Stoddard CS. Climate change effects on the processing tomato growing season in California using growing degree day model. Modeling Earth Systems and Environment. 2018;4(2):765-75.
 15. Chakravarty NV, Sastry PS. Phenology and accumulated heat unit relationships in wheat under different planting dates in the Delhi region. Agricultural science progress. 1983;1:32-42.
 16. Nuttonson MY. Some preliminary observations of phenological data as a tool in the study of photoperiodic and thermal requirements of various plant material. Vernalization and photoperiodism. Chronica Botanica Co., Waltham, Mass. 1948;129-43.
 17. Haider SA, Alam MZ, Alam MF, Paul NK. Influence of different sowing dates on the phenology and accumulated heat units in wheat. J. Biol. Sci. 2003;3(10):932-9.
 18. Praveen KV, Patel SR, Choudhary JL, Bhelawe S. Heat unit requirement of different rice varieties under Chhattisgarh Plain zones of India. Journal of Earth Science & Climatic Change. 2014;5(1).
 19. Pandey IB, Pandey RK, Dwivedi DK, Singh RS. Phenology, heat unit requirement and yield of wheat (*Triticum aestivum*) varieties under different crop-growing environment. Indian Journal of Agricultural Sciences. 2010;80(2):136-40.
 20. Kaur A, Pannu RK, Buttar GS. Impact of nitrogen application on the performance of wheat (*Triticum aestivum*) and nitrogen use efficiency under different dates of sowing. Indian Journal of Agronomy. 2010;55(1):40-5.

© 2022 Sabaridheepa et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/91613>