



# Effect of FYM and Different Levels of Panchgavya on Growth and Establishment of Papaya (*Carica papaya* L.) cv. Red Lady

Saurav Kumar <sup>a++\*</sup>, Samir E. Topno <sup>a#</sup> and Vijay Bahadur <sup>a†</sup>

<sup>a</sup> Department of Horticulture, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj-211007, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i6891>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/117203>

Original Research Article

Received: 09/03/2024  
Accepted: 12/05/2024  
Published: 13/05/2024

## ABSTRACT

Today, with the advancement of environment pollution and health that is obtained from the improper use of inorganic fertilizer, production and use of organic fertilizer is considered as an important approach in the field of soil science in the world's interest to investors. Research was conducted during the period 2023 on central research farm of department of horticulture, (NAI) SHUATS, Prayagraj. Field trial was designed on Randomized Block Design with three replications and ten treatments. Seedling were raised in polybag and 45 days old seedling were transplanted to the main field. Manure and fertilizer were applied as per treatment in ten equal split doses at 30,60 and

<sup>++</sup> M. Sc. Scholar;

<sup>#</sup> Assistant Professor;

<sup>†</sup> Associate Professor;

\*Corresponding author: E-mail: [skm.chanda@gmail.com](mailto:skm.chanda@gmail.com);

**Cite as:** Kumar, S., Topno, S. E., & Bahadur, V. (2024). Effect of FYM and Different Levels of Panchgavya on Growth and Establishment of Papaya (*Carica papaya* L.) cv. Red Lady. *Journal of Advances in Biology & Biotechnology*, 27(6), 322–326. <https://doi.org/10.9734/jabb/2024/v27i6891>

90days intervals. The treatment T<sub>9</sub> (RDF) FYM+ Panchgavya (90ml/l) registered significantly maximum vegetative growth, chlorophyll content and survivality %. On the basis of our experimental findings, it can be concluded that the best result was found in treatment T<sub>9</sub> (RDF) FYM+ Panchgavya (90ml/l) in term of vegetative growth parameters viz., plant height (cm), Number of leaves per plant, stem diameter (mm), leaf area (cm<sup>2</sup>) and root to shoot ratio, leaf area index (LAI), chlorophyll content and survivality % of papaya (*Carica papaya* L.) cv. Red lady.

**Keywords:** Growth; NPK; organic manure; Panchgavya; papaya; survival.

## 1. INTRODUCTION

Papaya (*Carica papaya* L.) is an evergreen herbaceous commercial fruit crop of tropical and subtropical regions and belongs to the family Caricaceae. Papaya is one among the fruits which has attained a great popularity in recent years because of gynodioecious nature, its easy cultivation, quick returns, adoptability to diverse soil and climatic conditions and above all its attractive delicious wholesome fruits having multifarious uses. [1]

Organic manure binds soil particles into structural units called aggregates. These aggregates help to maintain a loose, open, granular condition. Organic matter increased water-holding capacity of soil and reduced surface run off and erosion due to infiltration. Organic matter serves as a reservoir of chemical elements that are essential for plant growth. Most of the soil nitrogen occurs in organic combination. Also, a considerable quantity of phosphorus and sulfur exist in organic forms upon decomposition, organic matter supplied the nutrients needed by growing plants, as well as many hormones and antibiotics [2].

FYM being a bulky organic manure improves soil aeration in addition to the supply of essential plant nutrients and organic matter thereby increasing the soils biological activities. FYM also provided room for better microbial establishment along with the accumulation of excess humus content [3]. So, conjunctive use of chemical fertilizers, cow-based bio-enhancers and organic fertilizers sustain crop production and maintain soil health. Cow based bio-enhancers like panchagavya [4]. This experiment was carried to study the objectives which are; To find out the suitable treatment in relation to establish Papaya and growth of payaya.

## 2. MATERIALS AND METHODS

The experiment was carried out at Horticulture Research Field, Department of Horticulture, Sam

Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, during the period May to August 2023. Its comes under subtropical climate zone prevailing in the winter and summer.

It is situated at latitude of 25.85°N and longitude of 81.15°E. The altitude of this place is 78 m from MSL. During the winter months, the temperature drops down as low as 2°C while in the summer the temperature reaches above 48°C hot desiccating winds are regular feature during summers whereas there may be occasional spell of frost during the winters. Papaya c.v. Red lady were collected from Department of Horticulture, Sam Higginbottom University of Agriculture Technology and Science, Praygaraj.

Seedling were raised in plogybag and 45 days old seedling were transplanted to the main field. The experiment was laid out in Randomized Block Design with tenth treatments and three replications. The treatment comprised of tenth FYM and Panchgavya. Likewise, T<sub>0</sub> Control, T<sub>1</sub> (RDF) FYM + Panchgavya (10ml/l), T<sub>2</sub> (RDF) FYM + Panchgavya (20ml/l), T<sub>3</sub> (RDF) FYM + Panchgavya (30ml/l), T<sub>4</sub> (RDF) FYM + Panchgavya (40ml/l), T<sub>5</sub> (RDF) FYM + Panchgavya (50ml/l), T<sub>6</sub> (RDF) FYM + Panchgavya (60ml/l), T<sub>7</sub> (RDF) FYM + Panchgavya (70ml/l), T<sub>8</sub> (RDF) FYM + Panchgavya (80ml/l) and T<sub>9</sub> (RDF) FYM+ Panchgavya (90ml/l). The seedlings of papaya were transplanted in the field adopting a spacing of 2 x 2 m. The data recorded on vegetative growth viz., plant height (cm), number of leaves per plant, stem diameter (mm), leaf area (cm<sup>2</sup>), leaf area index (LAI), chlorophyll content and root to shoot ratio were analyzed statistically [5].

## 3. RESULTS AND DISCUSSION

### 3.1 Plant Height

Data of vegetative growth parameters of papaya influenced by application of RDF, FYM and

different levels of panchgavya are presented in Table 1. Observation on plant height (cm), Number of leaves per plant, stem diameter (cm), leaf area (cm<sup>2</sup>) and leaf area index LAI recorded at 30 DAT intervals from 30 to 90 DAT showed significant variation among the treatment. Statistical analysis at 90 DAT revealed a highest plant height of (59.56cm) was found in treatment T<sub>9</sub> (RDF) FYM + Panchgavya (90ml/l), which was on par with T<sub>8</sub> (RDF) FYM+ Panchgavya (80ml/l) and differed significantly from other treatments. It was followed by (RDF) FYM + Panchgavya (60ml/l), (RDF) FYM + Panchgavya (70ml/l), (RDF) FYM + Panchgavya (40ml/l) and (RDF) FYM + Panchgavya (50ml/l). Whereas the minimum plant height (38.87) was found in treatment T<sub>0</sub> Control [6].

### 3.2 Number of Leaves Per Plant

Statistical analysis at 90 DAT revealed a highest number of leaves per plant of (18.76) was found in treatment T<sub>9</sub> (RDF) FYM+ Panchgavya (90ml/l), which was on par with T<sub>8</sub> (RDF) FYM + Panchgavya (80ml/l) and differed significantly from other treatments. It was followed by (RDF) FYM+ Panchgavya (70ml/l), (RDF) FYM + Panchgavya (60ml/l) and (RDF) FYM + Panchgavya (50ml/l). Whereas the minimum number of leaves per plant (12.68) was found in treatment T<sub>0</sub> Control [6].

### 3.3 Stem Diameter

Statistical analysis at 90 DAT revealed a highest stem diameter (cm) of (2.64) was found in treatment T<sub>9</sub> (RDF) FYM+ Panchgavya (90ml/l), which was on par with T<sub>8</sub> (RDF) FYM + Panchgavya (80ml/l) and differed significantly from other treatments. It was followed by (RDF) FYM + Panchgavya (50ml/l), (RDF) FYM + Panchgavya (60ml/l), (RDF) FYM + Panchgavya (70ml/l). Whereas the minimum stem diameter (mm) (2.02) was found in treatment T<sub>0</sub> Control [6].

### 3.4 Leaf Area

Statistical analysis at 90 DAT revealed a highest leaf area (cm<sup>2</sup>) of (143.72) was found in treatment T<sub>9</sub> (RDF) FYM + Panchgavya (90ml/l), which was on par with T<sub>8</sub> (RDF) FYM + Panchgavya (80ml/l) and differed significantly from other treatments. It was followed by (RDF)

FYM + Panchgavya (50ml/l), (RDF) FYM + Panchgavya (60ml/l), (RDF) FYM + Panchgavya (70ml/l). Whereas the minimum leaf area (cm<sup>2</sup>) (106.55) was found in treatment T<sub>0</sub> Control [6].

### 3.5 Leaf Area Index (LAI)

Statistical analysis at 90 DAT revealed a highest leaf area index (LAI) of (3.59) was found in treatment T<sub>9</sub> (RDF) FYM+ Panchgavya (90ml/l), which was on par with T<sub>8</sub> (RDF) FYM + Panchgavya (80ml/l) and differed significantly from other treatments. It was followed by (RDF) FYM + Panchgavya (50ml/l), (RDF) FYM + Panchgavya (60ml/l), (RDF) FYM+ Panchgavya (70ml/l). Whereas the minimum leaf area index (LAI) (2.66) was found in treatment T<sub>0</sub> Control [6].

The higher nutrient content and metabolic levels enhanced the growth parameters, ultimately leading to higher yield. The results are also in close conformity with the findings of in papaya [7].

### 3.6 Chlorophyll Content

The data obtained on chlorophyll content (spad value), root to shoot ratio and survival (%) in papaya after application of (RDF) FYM and different levels of panchgavya are illustrated in Table 2. Statistical analysis of the results indicated that the Chlorophyll content (SPAD value) in papaya leaf provided with different levels of panchgavya and (RDF) FYM differed significantly. The treatment T<sub>9</sub> (RDF) FYM + Panchgavya (90ml/l) registered significantly maximum Chlorophyll content (53.78 SPAD value). It was found to be on par with (RDF) FYM + Panchgavya (60ml/l), (RDF) FYM + Panchgavya (70ml/l). Whereas the minimum Chlorophyll content (SPAD value) (49.10) was found in treatment T<sub>0</sub> Control [8].

### 3.7 Root to Shoot Ratio

The treatment T<sub>9</sub> (RDF) FYM + Panchgavya (90ml/l) registered significantly maximum root to shoot ratio (0.245). It was found to be on par with (RDF) FYM+ Panchgavya (60ml/l), (RDF) FYM + Panchgavya (70ml/l). Whereas the minimum root to shoot ratio (0.210) was found in treatment T<sub>0</sub> Control [8].

**Table 1. Effect of FYM and different levels of panchgavya on vegetative growth parameters of papaya (*Carica papaya* L.) cv. Red lady**

Treatment No.	Treatment details	Vegetative growth parameters											
		Plant height (cm)				Number of leaves per plant				Stem diameter (cm)			
		Initial	30 DAS	60 DAS	90 DAS	Initial	30 DAS	60 DAS	90 DAS	Initial	30 DAS	60 DAS	90 DAS
T <sub>0</sub>	Control	15.43	22.51	31.62	38.87	4.47	6.24	9.04	12.68	1.06	1.24	1.48	2.02
T <sub>1</sub>	(RDF) FYM+ Panchgavya (10ml/l)	18.73	30.06	42.41	48.48	5.72	7.11	11.10	13.07	1.11	1.29	1.76	2.25
T <sub>2</sub>	(RDF) FYM+ Panchgavya (20ml/l)	20.27	29.63	38.34	46.76	5.48	7.23	11.53	13.81	1.13	1.33	1.77	2.30
T <sub>3</sub>	(RDF) FYM+ Panchgavya (30ml/l)	19.76	30.73	42.43	48.74	5.86	7.28	12.09	15.36	1.16	1.37	1.81	2.34
T <sub>4</sub>	(RDF) FYM+ Panchgavya (40ml/l)	18.10	30.49	46.67	51.83	5.90	7.43	12.34	15.75	1.21	1.42	1.83	2.37
T <sub>5</sub>	(RDF) FYM+ Panchgavya (50ml/l)	20.11	27.33	42.41	51.08	6.07	8.16	12.57	16.03	1.26	1.52	1.87	2.45
T <sub>6</sub>	(RDF) FYM+ Panchgavya (60ml/l)	19.23	32.49	45.01	55.31	5.27	8.27	13.20	16.40	1.31	1.57	1.92	2.50
T <sub>7</sub>	(RDF) FYM+ Panchgavya (70ml/l)	18.51	32.40	44.02	52.56	6.06	8.38	13.84	17.19	1.35	1.64	1.96	2.55
T <sub>8</sub>	(RDF) FYM+ Panchgavya (80ml/l)	21.73	34.73	49.29	57.53	6.22	8.51	15.37	18.24	1.41	1.74	2.05	2.60
T <sub>9</sub>	(RDF) FYM+ Panchgavya (90ml/l)	23.97	36.47	50.13	59.56	6.42	9.01	15.53	18.76	1.58	1.83	2.09	2.64
	F-Test	-	S	S	S	-	S	S	S	-	S	S	S
	S.Ed.(+)	-	1.24	0.92	1.11	-	0.22	0.23	0.28	-	0.03	0.03	0.03
	C.D at 0.5%	-	2.61	1.93	2.32	-	0.45	0.48	0.59	-	0.06	0.05	0.06
	CV		4.956	2.606	2.653		3.414	2.208	2.174		2.343	1.707	1.501

**Table 2. Effect of FYM and different levels of panchgavya on vegetative growth parameters of papaya (*Carica papaya* L.) cv. Red lady**

Treatment No.	Treatment details	Vegetative growth parameters										
		Leaf area (cm <sup>2</sup> )				LAI (Leaf area index)				Chlorophyll content (SPAD value)	root to shoot ratio	Survival (%)
		Initial	30 DAS	60 DAS	90 DAS	Initial	30 DAS	60 DAS	90 DAS			
T <sub>0</sub>	Control	28.50	57.43	77.45	106.55	0.71	1.44	1.94	2.66	49.10	0.210	46.67
T <sub>1</sub>	(RDF) FYM+ Panchgavya (10ml/l)	29.19	62.63	85.18	125.22	0.73	1.57	2.13	3.13	50.18	0.225	66.67
T <sub>2</sub>	(RDF) FYM+ Panchgavya (20ml/l)	30.57	63.78	85.11	127.50	0.76	1.59	2.13	3.19	50.51	0.241	60.00
T <sub>3</sub>	(RDF) FYM+ Panchgavya (30ml/l)	34.30	61.65	91.08	129.73	0.86	1.54	2.28	3.24	50.43	0.231	66.67
T <sub>4</sub>	(RDF) FYM+ Panchgavya (40ml/l)	34.63	63.36	92.20	131.91	0.87	1.58	2.31	3.30	50.22	0.237	73.33
T <sub>5</sub>	(RDF) FYM+ Panchgavya (50ml/l)	35.07	61.78	97.45	133.28	0.88	1.54	2.44	3.33	50.38	0.229	73.33
T <sub>6</sub>	(RDF) FYM+ Panchgavya (60ml/l)	35.40	64.26	98.54	136.49	0.88	1.61	2.46	3.41	51.49	0.218	73.33
T <sub>7</sub>	(RDF) FYM+ Panchgavya (70ml/l)	36.10	66.70	101.54	137.04	0.90	1.67	2.54	3.43	53.46	0.233	80.00
T <sub>8</sub>	(RDF) FYM+ Panchgavya (80ml/l)	36.64	68.76	103.69	140.43	0.92	1.72	2.59	3.51	53.78	0.232	80.00
T <sub>9</sub>	(RDF) FYM+ Panchgavya (90ml/l)	37.91	70.67	105.53	143.72	0.95	1.77	2.64	3.59	54.39	0.245	93.33
	F-Test	-	S	S	S	-	S	S	S	S	S	S
	S.Ed.(+)	-	1.29	1.28	1.31	-	0.03	0.03	0.03	0.465	0.008	11.62586
	C.D at 0.5%	-	2.71	2.68	2.76	-	0.07	0.07	0.07	0.977	0.017	24.42504
	CV		2.469	1.668	1.224		2.542	1.604	1.184	1.109	4.270	21.312

### 3.8 Survival Percentage

The treatment T<sub>9</sub> (RDF) FYM + Panchgavya (90ml/l) registered significantly maximum survival (%) (93.33). It was followed by (RDF) FYM + Panchgavya (60ml/l), (RDF) FYM + Panchgavya (70ml/l) and (RDF) FYM+ Panchgavya (80ml/l). Whereas the minimum survival (%) (46.67) was found in treatment T<sub>0</sub> Control. The findings of current investigation are in agreement with that of Reddy et al. [8] where they reported that in papaya cv Surya, crop growth was better with 75% recommended dose of fertilizer applied as farm yard manure + vermicompost, which was significantly superior that in 100% recommended dose of fertilizer and no manure/fertilizer treatment and in conformation with that of in sweet orange (*Citrus sinensis* L.) Osbeck) cv Mosambi [9].

### 4. CONCLUSION

From the present investigation it is concluded that treatment (T<sub>9</sub>: (RDF) FYM + Panchgavya 90ml/l) performed best in terms of growth parameters viz., plant height, number of leaves, stem diameter, leaf area and leaf area index, chlorophyll content, root to shoot ratio and survival percentage of Papaya c.v. Red Lady.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Agrawal Baldev and Sahu GD. Effect of organic manure and bio-fertilizer on quality and benefit: Cost ratio of papaya (*Carica papaya* L.) Cv. Red Lady. Journal of Pharmacognosy and Phytochemistry. 2021;10(2):700-702.
2. Devi Okram Bijaya, Singh Yumnam Somi. Effect of Organic Amendments on Growth, Yield and Quality of Papaya (*Carica papaya* L.) cv Vinayak. Environment and Ecology, 2023;41(1B):522—531
3. Mohapatra A, Dash DK, Tripathy P. Comparative analysis of various organic amendments on tree growth and nut yield of cashew (*Anacardium occidentale* L.). International Journal of Agriculture, Environment and Biotechnology. 2016;9(2):225-230.
4. Patel Vrutti and Ramdevputra MV. Effect of different mulches and liquid organic: Inorganic formulations on flowering and growth of papaya (*Carica papaya* L.) cv. GJP-1. International Journal of Chemical Studies. 2021;9(2):765-768
5. Panse VC, Sukhatme PV. Statistical methods for agricultural workers, ICAR, New Delhi (INDIA); 1967.
6. Shrivastava A. Integrated nutrient management (*Carica papaya*L.). Ph.D. Thesis, N.D. University of Agriculture and Technology, Faizabad (U.P.) INDIA; 2008.
7. Chaudhri SM, Shinde SH, Dahiwalkar SD, Dana wale NJ, Shiras HK, Berad SM. Effect of fertigation through drip on productivity of papaya.J. Maharashtra Agric. Univ. 2001;26(1):1820.
8. Reddy YTN, Reju MK, Ganeshmurthy AN, Pannerselvam P, Prasad SRS. Influence of organic practices on growth and fruit yield in papaya cv Surya. J Hort Sci. 2013;8(2): 246— 248.
9. Nanaso GM, Panwar PS. Studies on use of organic manures and biofertilizers on growth, yield and quality in Sweet Orange (*Citrus sinensis* (L.) Osbeck) cv Mosambi. MSc thesis, MPKV, Rahuri; 2020. Available:https://krishikosh.egranth.ac.in/handle/1/5810157148

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/117203>