



Genetic Variability and Correlation Studies in Chilli (*Capsicum annuum* L.) Under Agro-Climatic Condition of Prayagraj

Preksha Yadav ^{a+++*}, Devi Singh ^{a#} and Vijay Bahadur ^{a†}

^a Department of Horticulture, Naini Agriculture Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i64655>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/116957>

Original Research Article

Received: 09/03/2024
Accepted: 12/05/2024
Published: 13/05/2024

ABSTRACT

The present study assesses the extent of genetic variability and correlation in 12 different genotypes of chillies for various compositional and yield properties for identifying predominant genotypes for involvement in coming breeding programs. One of India's most significant crops for both vegetables and spices is the chilli. In the world, India is the top producer of dried chili. Around the world, people utilize chilli as a spice and to make drinks and medications. Since the beginning of time, curry has been an essential component of Indian cuisine. The field trial was conducted from December 2022 to June 2023 at the Horticulture Research Farm, Department of Horticulture,

⁺⁺ M.Sc. Research Scholar;

[#] Associate Professor;

[†] Associate Professor and Head;

*Corresponding author: E-mail: yadavpreksha4@gmail.com;

Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.), India. There is a great need and opportunity to assess and create location-specific cultivars with increased production potential. Twelve distinct genotypes of chillies were gathered from various regions of India for this inquiry in order to examine how well they performed in the Prayagraj agroclimatic conditions. The genetic data on yield and attributes associated with yield would be useful in developing a productive breeding program to enhance the chilli crop and create genotypes and hybrids that show promise.

Keywords: Chilli; genetic variability; capsanthin.

1. INTRODUCTION

“Chilli (*Capsicum annuum* L.) shrubs are perennial and short-lived; they can develop up to 1.5 m in height; their stems are woody at the base, fleshy, and either erects or semi-prostrate, and the shrub comprises of a primary tap root with various lateral roots. The leaves can grow up to 12 cm long and 7.5 cm wide and are changing in shape with a pointed tip. Chilli flowers are independently or in small groups of two or three flowers. They're small and bisexual, with five to six petals each; the flowers of *Capsicum annuum* are white-green. Red chillies get their color from a coloring compound called capsanthin and have a hot, pungent taste due to a chemical called capsaicin, and the numerous small chilli seeds moreover contain capsaicin. Chilli is basically cultivated for fruits which are utilized as vegetables in pharmaceutical as a stimulant and source of oleoresin” [1].

“Fruit yield as well as quality improvement efforts proceed to be the major aims of the chilli improvement program. Fruit yield is a complex inherited character influenced by a few properties of the plant. Area-based screening for upgrading the productivity of this crop is an important step to expanding production” [2]. “A wide range of variability is available in chilli genotypes, which provide great scope for improving fruit yield through systematized breeding. An estimation of the genetic variability show in the germplasm of a crop is a pre-requisite for designing a successful breeding program” [3].

In Uttar Pradesh, chillies are basically developed in the eastern districts, viz., Ballia, Azamgarh, Mirzapur, Basti, Faizabad, and Ghazipur. The larger part of cultivators is still developing local cultivars. Other than soil and climatic components, the cultivar itself is really important in regard of its performance with respect to earliness, disease resistance, and yield [4,5]. Numerous cultivars have been developed and

suggested by different research institutes and State Agricultural Universities, but the adoptability and yielding capacity of the cultivars aren't the same in all regions. So, there's a pressing demand for a appropriate variety in Prayagraj agro-climatic conditions. Therefore, an attempt was made in the present research at the Department of Horticulture, SHUATS, Prayagraj, to assess the extent of genetic variability and correlation in 12 different genotypes of chillies for various compositional and yield properties for identifying predominant genotypes for involvement in coming breeding programs.

2. MATERIALS AND METHODS

The field trial was conducted from December 2022 to June 2023 at the Horticulture Research Farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.) for the genetic variability and correlation studies in chilli under the agro climate conditions of Prayagraj. This research was carried out in a randomized block design (RBD) with 12 genotypes (released varieties and local collection) which were collected from diverse sources (Table 1).

3. RESULTS AND DISCUSSION

Significant differences were obtained among the genotypes for all the characters, showing satisfactory variability among all the genotypes. F1-6106 took a least number of days (37.4) to first flowering taken after by KSP-1350 (42.2), and most extreme days by TMPH-409 (49.93); however, F1-6106 also took a least day to 50% flowering, taken after by Pusa Jwala (48.2), and greatest days by VNR-305 (53.35). F1-6106 took the least number of days (63.34) to the first harvest, taken after by KSP-1350 (65.47) and the greatest days (73.17) taken by VNR-305. The highest plant height was recorded (89.74 cm) in F1-6106, taken after by Pusa Jwala (88.67 cm), and the lowest plant height was recorded (75.8

cm) in VNR-305. The most extreme number of branches recorded (19.33) in F1-6106, taken after by Pusa Jwala (18.07), and the least number of branches recorded was (9.53) in TMPH-484. F1-6106 appears the highest leaf area (29.74 cm²), taken after by Pusa Jwala (29.37 cm²), and the lowest leaf area appeared by VNR-305 (23.32 cm²). The maximum number of fruit plant⁻¹ was recorded (63.99) in F1-6106, taken after by Pusa Jwala (63.21) and the minimum number of fruit plant⁻¹ (52.33) in VNR-305. The maximum fruit weight was recorded in F1-6106 (6.41 g), taken after by F1-hybrid (5.31 g), and the minimum fruit weight was recorded in VNR-305 (2.36 g). The maximum fruit length was recorded in F1-6106 (10.31 cm), taken after by Pusa Jwala (9.98 cm) and the minimum fruit length were recorded in VNR-305 (6.44 cm). The greatest fruit diameter was recorded in F1-6106

(11.00 mm), taken after by Pusa Jwala (10.33 mm) and the least fruit diameter was recorded in VNR-305 (5.83 mm). The highest fruit yield plant⁻¹ (g) was recorded in F1-6106 (382.91 g), taken after by Pusa Jwala (332.14 g), and the least fruit yield plant⁻¹ was recorded (145.32 g) in VNR-305. The highest fruit yield ha⁻¹ (t) was recorded in F1-6106 (12.76 t), taken after by Pusa Jwala (11.07 t), and the least fruit yield plant⁻¹ was recorded (4.84 t) in TMPH-404. The highest TSS content was recorded in VNR-305 (9.73^o), Taken after by hybrid eagle (9.63^o), and the least TSS content was recorded (6.07^o) in F1-6106. The highest ascorbic acid (mg/100g) content was recorded in F1-6106 (196.46 mg/100g), taken after by VNR-305 (187.63 mg/100g), and the lowest ascorbic acid content was recorded (105.56 mg/100g) in hybrid eagle (Table 2 and Table 3) [6,7].

Table 1. List of chilli genotypes and their sources used in study

Notation	Genotypes	Sources
V ₁	KSP-1350	Kalash Seeds Pvt. Ltd
V ₂	KSP-1342	Kalash Seeds Pvt. Ltd
V ₃	Hybrid Eagle	Acsen HyVeg Pvt. Ltd
V ₄	VNR-305	VNR Seed Pvt. Ltd.
V ₅	Shravani 301	Pancha Ganga Seeds Pvt. Ltd.
V ₆	TMPH-484	TriMurti Plant Science Pvt. Ltd.
V ₇	TMPH-404	TriMurti Plant Science Pvt. Ltd.
V ₈	TMPH-407	TriMurti Plant Science Pvt. Ltd.
V ₉	TMPH-409	TriMurti Plant Science Pvt. Ltd.
V ₁₀	F1-Hybrid	Eco Green Hybrid Seeds Co.
V ₁₁	Pusa Jwala	IARI, NEW DELHI
V ₁₂	F1-6106	Eco Care Seeds

Table 2. Mean performance of chilli genotypes for growth and earliness parameters

Genotypes	Plant height (cm)	Number of branches	Leaf area (cm ²)	Days taken to first flowering	Days taken to 50% of flowering	Days to first harvest
V1	79	16.53	24.43	442.2.2	49.56	65.47
V2	83.54	14.27	25.67	43.73	48.53	69.87
V3	76.2	12.26	24.67	44.2	48.94	70.33
V4	75.8	11.0	23.32	48.8	53.35	73.17
V5	82.64	13.8	25.54	46.87	50.67	70.67
V6	76.9	9.53	27.6	42.73	51.1	70.04
V7	83.6	12.0	26.57	48.44	52.6	70.4
V8	82.94	17.0	27.64	47.37	51.27	70.63
V9	80.8	13.6	27.77	49.93	50.37	72.4
V10	85.14	13.07	25.04	42.8	49.8	70.93
V11	88.67	18.07	29.37	43.13	48.2	69.4
V12	89.74	19.33	29.74	37.4	42.47	63.34
S.Ed(±)	5.78	2.09	2.09	0.41	0.73	1.87
CV	8.63	17.98	9.72	1.12	1.80	3.29
CD _{0.05}	1.97	4.33	4.35	0.85	1.52	3.89

Table 3. Mean performance of chilli genotypes for quality and yield parameters

Genotypes	Number of fruit plant ⁻¹	Average fruit weight (g)	Average fruit length (cm)	Average fruit diameter (mm)	Fruit yield plant ⁻¹ (g)	Fruit yield ha ⁻¹ (t)	TSS (Brix ^o)	Ascorbic acid (mg/100g)
V1	59.18	4.47	7.69	9.27	235.19	7.83	6.47	132.1
V2	59.29	4.69	7.38	9.93	250.25	8.34	7.16	135.3
V3	55.4	4.23	7.12	7.01	250.41	8.65	9.63	105.56
V4	52.33	2.36	6.44	5.83	145.62	4.85	9.73	187.63
V5	59.4	3.35	7.94	7.31	209.91	6.99	7.66	126.8
V6	59.19	2.64	6.75	7.24	183.50	6.16	9.63	149.4
V7	52.82	2.43	7.35	7.65	146.05	4.84	6.47	166.6
V8	59.51	2.38	6.99	6.46	149.85	4.91	9.4	186.53
V9	53.56	2.62	7.03	6.12	154.74	5.84	7.7	193
V10	59.58	5.31	8.29	9.44	327.31	10.90	9.6	186.23
V11	63.21	5.11	9.98	10.33	332.14	11.07	7.5	185.3
V12	63.99	6.41	10.31	11.00	382.91	12.76	6.07	196.46
S.Ed(±)	2.73	0.55	0.12	0.45	30.37	3.22	0.23	0.72
CV	5.75	17.47	1.94	6.77	16.11	17.56	3.34	0.57
CD _{0.05}	5.66	1.13	0.26	0.93	62.98	6.69	0.46	1.49

The extent of variability show in 12 genotypes of chilli was measured in terms of range, mean, PCV, GCV, heritability, and genetic advance (Table 4). All the genotypes differed significantly with regard to the distinctive characters examined. A wide range of variation was observed in all the characters. Gupta et al. [8,9,7] moreover detailed a wide range of variation for most extreme of the characters. The genotypic coefficient of variance (GCV) and phenotypic coefficient of variance (PCV) were high for fruit weight (36.40 g and 36.44 g) taken after by fruit

yield plant⁻¹ (31.95 g and 41.67 g), and fruit diameter (21.82 mm and 21.80 mm), appearing great differences for these traits. GCV in common, was lower than the PCV (Table 4), which shown a close affiliation between phenotype and genotype [8,9,10]. Low genotypic coefficient of variance (GCV) and phenotypic coefficient of variance were recorded for days taken to 50% flowering (5.05 and 5.06), taken after by the number of fruits plant⁻¹(6.54 and 6.55) [11,1,10].

Table 4. Genetic variability components for major characters in chilli

Characters	Range	Mean	Coefficient of variance (%)		Heritability (%)	Genetic advance	GAM (%)
			GCV	PCV			
Days taken to first flowering	49.93	44.77	7.79	7.80	7.80	7.17	16.03
Days taken to 50% flowering	53.27	49.78	5.05	5.06	5.06	5.16	10.37
Plant height	58.66	81.92	6.65	5.66	5.66	9.53	11.64
Number of branches	12.62	14.19	21.01	21.04	21.04	6.13	43.21
Number of fruits plant ⁻¹	63.99	58.13	6.54	6.55	6.55	7.83	13.47
Fruit length	7.77	7.73	15.82	15.86	15.86	2.51	32.49
Fruit diameter	11.0	8.15	21.80	21.82	21.82	3.65	44.86
Fruit weight	6.41	3.83	36.40	36.44	36.44	2.86	74.91
Fruit yield plant ⁻¹ (g)	382.91	225.21	31.95	41.67	41.67	115.20	51.19
Leaf area	29.73	26.44	7.39	7.44	7.44	4.01	15.07
TSS	9.73	8.07	17.63	17.74	17.74	2.91	36.09
Ascorbic acid	196.47	162.61	19.21	19.12	19.12	64.38	39.57

Table 5. Correlation coefficient (genotypic) of different characters in chilli

Characters	1	2	3	4	5	6	7	8	9	10	11	12
1	1.0000											
2	0.819**	1.0000										
3	-0.440**	-0.635**	1.0000									
4	-0.473**	-0.690**	0.731**	1.0000								
5	-0.796**	-0.716**	0.685**	0.705**	1.0000							
6	-0.679**	-0.819**	0.856**	0.745**	0.748**	1.0000						
7	-0.821**	-0.792**	0.754**	0.634**	0.784**	0.834**	1.0000					
8	-0.841**	-0.887**	0.645**	0.623**	0.721**	0.836**	0.913**	1.0000				
9	-0.737**	-0.720**	0.555**	0.635**	0.693**	0.769**	0.906**	0.981**	1.0000			
10	-0.295**	-0.520**	0.681**	0.566**	0.547**	0.615**	0.366*	0.256	0.064	1.0000		
11	-0.528**	-0.479**	0.549**	0.384**	0.729*	0.523**	0.380*	0.393*	0.252	0.707	1.0000	
12	0.070	-0.037	0.455	0.316	0.091	0.307	0.054	0.022	0-.112	0.472**	0.384*	1.0000

1-Days taken to 1st flowering, 2- Days taken to 50% flowering, 3-Plant height, 4-Number of branches, 5-Number of fruits plant-1, 6-Fruit length, 7-Fruit diameter, 8- Fruit weight, 9-Fruit yield(g plant-1), 10-Leaf area, 11-TSS, 12-Ascorbic Acid
 *Significant at $P<0.01$; **Significant at $P<0.05$

Heritability is a parameter of huge importance to breeders, as its magnitude indicates the dependability with which a genotype can be recognized through its phenotypic expression (Table 4). Heritability in a wide sense was observed to be high for all the traits examined. Heritability estimates along with genetic advance are more useful than heritability values alone in reading the selection of the best individuals [1,12] In the present studies, fruit yield plant⁻¹, fruit weight, ascorbic acid, and the number of branches plant⁻¹ shown high genetic advance as percentage of the mean along with high heritability. These results shown the impact of additive gene action. High genetic advance for the number of fruits plant⁻¹ and fruit yield plant⁻¹ [8,13,7].

Genotypic correlation of 12 yield and yield attributing characters displayed in (Table 5) shown that fruit yield plant⁻¹ appears a positive and highly significant correlation with plant height (0.555), number of branches (0.635), number of fruit plant⁻¹ (0.693), fruit length (0.769), fruit diameter (0.906) and fruit weight (0.981); similarly, fruit weight shows a positive correlation with plant height (0.645), number of branches (0.623), number of fruits plant⁻¹ (0.721), fruit length (0.836), and fruit diameter (0.913). Fruit diameter appears a positive correlation with plant height (0.754), number of branches (0.634), number of fruit plant⁻¹ (0.784) and fruit length (0.834) though fruit length appears a positive correlation with plant height (0.856), number of branches (0.745) and number of fruit plant⁻¹ (0.748). The number of fruit plant⁻¹ appears a positive correlation with only plant height (0.685) and the number of branches (0.705) [8,14,15] moreover detailed that fruit yield plant⁻¹ had a positive and highly significant correlation with the number of fruits plant⁻¹ and fruit length. The positive association of fruit weight with fruit breadth and fruit length shown that selection of only one of the traits might lead to an increase in the size of fruit [8,16-18].

4. CONCLUSION

The best genotypes for Prayagraj's agroclimatic conditions were determined to be F1-6106 and Pusa Jwala, based on the overall performance of the different genotypes under investigation. These may be employed in breeding efforts to create better cultivars and hybrids, or they can be utilized directly as cultivars. The characters, namely, plant height, fruit length, number of fruits plant⁻¹, fruit weight, and fruit yield, were found to

be the most important traits for improving the genotypes based on mean performance and other genetic parameters of various growth and yield characters. The number of branches plant⁻¹ can be regarded as the second most important character for selection in chilli genotypes. Based on estimates of the correlation coefficient, numbers of fruits plant⁻¹ and fruit weight were adjudged as yield attributing characters that needed to be focused during selection.

ACKNOWLEDGEMENT

This paper forms a part of M.Sc. (Horticulture) Vegetable Science. The thesis of Preksha Yadav was submitted to SHUATS, 2024. The authors are sincerely thankful to Dr. Devi Singh, Associate Prof. Department of Horticulture, SHUATS, Prayagraj.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Samadia DK. Genetic variability studies in chilli germplasm under hot arid ecosystem. Indian J. Hort. 2007;64:477–479
2. Datta S, Jana JC, Evaluation of chilli (*Capsicum annum* L.) germplasm suitable for spices. Paper presented in 1st Indian Horticulture Congress 2004 on "Improving Productivity Quality Postharvest Management and Trade in Horticultural Crop.", 6–9 November, New Delhi; 2004.
3. Parkash C, Estimation of genetic variability and implications of direct effects of different traits on leaf yield in bathua (*Chenopodium album*). Indian J. Agri. Sci. 2012;82:71–74.
4. Gokulakrishnan J, Nagaraja C, Prakash M. Genetic variability and correlation studies in chilli (*Capsicum annum* L.). Plant Archives (09725210). 2020;20(1).
5. Swetha B, Devi HU, Sankari A, Geethanjali S, Sudha M. Variability studies and genetic divergence in chilli (*Capsicum* spp.) genotypes using multivariate analysis. Electronic Journal of Plant Breeding. 2023; 14(3):928-37.
6. Deb P, Datta S & Paul P K, Studies on the performance of chilli genotypes under Terai zone of West Bengal. Indian J. Hort. 2008;65:353–353.

7. Warshamana IK, Vikram A, Kohli UK, Genetic evaluation for physicochemical and yield traits in chilli (*Capsicum annuum* L.) under hilly conditions of Himachal Pradesh. The Hort. J. 2008;21: 62–66.
8. Gupta AM, Singh D, Kumar A, Genetic variability, genetic advance and correlation studies in chilli (*Capsicum annuum* L.). Indian J. Agri. Sci. 2009;79:221–223.
9. Munshi AD, Behera TK, Genetic variability, heritability and genetic advance for some traits in chilli. (*Capsicum annuum* L.) Veg. Sci. 2000;27:39–41.
10. Singh MD, Laisharam JM, Bhagirath T, Genetic variability in local chillies (*Capsicum annuum* L.) of Manipur. Indian J.Hort. 2005;62:203–205.
11. Kumar K, Baswana KS, Partap PS, Genetic variability and heritability studies in chilli. (*Capsicum annuum* L.) Haryana J. Hort. Sci. 1999;28:125–126.
12. Verma SK, Singh RK, Anja RR, Genetic variability and correlation studies in chilli. Prog. Hort. 2004;36:113–117.
13. Sreelathakumary Rajamony L, Variability, heritability and correlation studies in chilli (*Capsicum* spp.) under shade. Indian J. Hort. 2002;59:77–83.
14. Hosamani RM, Shivkumar, Correlation and path coefficient analysis in chilli. Indian J. Hort. 2008;65:349–352.
15. Singh DK, Singh A, Assessment of variability parameters and character association for quantitative traits in chilli (*Capsicum annuum* L.). Prog. Agri. 2011; 11:113-116.
16. Anita Kerketta JP, Collis, Evaluation of Chilli (*Capsicum annuum* L.) Genotypes for Growth, Yield and Quality Characters under Allahabad Agro Climatic Conditions. Int. J. Pure App. Biosci. 2018;6(4):451-455.
17. Amit K, Ahad I, Kumar V, Thakur S. Genetic variability and correlation studies for growth and yield characters in chilli (*Capsicum annuum* L.). Journal of Spices & Aromatic Crops. 2014;23: (2).
18. Rohit Kumar RS, Meena AK, Verma, Hemant Ameta and Alka Panwar, Analysis of Genetic Variability and Correlation in Fennel (*Foeniculum Vulgare* Mill.) Germplasm. Agri Res & Tech: Open Access J. 2017;3(4):ARTOAJ.MS. ID.555616.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/116957>