



Hyer-Ulam-Stability of Generalized Quadratic Functional Equation in Random Normed Spaces

Sushma Devi ^{a*}

^a Kanya Mahavidyalaya, Kharkhoda, Sonapat, Haryana, India.

Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/ARJOM/2023/v19i7671

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/98066>

Original Research Article

Received: 28/01/2023

Accepted: 29/03/2023

Published: 12/04/2023

Abstract

The purpose of this paper to propose a finite variable generalized quadratic functional equation with solution. Also investigate Hyers Ulam stability in Random normed space by direct method.

Keywords: Quadratic functional equation; hyers-ulam stability; random normed space.

Mathematics subject classification: 54E40, 39B82, 46S50.

1 Introduction

Functional equations play an important part in the study of stability. In 1940, the stability problems of functional equations about group homomorphisms was introduced by Ulam [1]. In 1941, Hyers [2] gave an affirmative answer to Ulam's question for additive groups (under the assumption that groups are Banach spaces). Hyers

*Corresponding author: Email: sheoransushma50841@gmail.com, lathersushma@yahoo.com;

theorem was generalized by Aoki [3] for additive mappings and by Rassias [4] for linear mappings by considering an unbounded Cauchy difference $\|f(x + y) - f(x) - f(y)\| \leq \varepsilon(\|x\|^p + \|y\|^p)$ for all $\varepsilon > 0$ and $p \in [0,1)$. Also Rassias [5] generalization theorem was delivered by Gavruta [6] who replaced $\varepsilon(\|x\|^p + \|y\|^p)$ by a control function $\varphi(x, y)$. The paper of Rassias has significantly influenced the development of what we now call the Hyers-Ulam-Rassias stability of functional equations. In 1982, J.M. Rassias [4] followed the modern approach of the Th.M.Rassias theorem in which he replaced the factor product of norms instead of sum of norms.

Recently the stability of many functional equations in various spaces like Banach spaces, modular spaces, fuzzy normed spaces and Random normed spaces etc. have been established by researchers [7-13]. Now we introduce new quadratic functional equation and obtain the Hyers- Ulam stability of quadratic function equation

$$\sum_{i=1}^n \varphi(-2v_i + \sum_{j=1, i \neq j}^n v_j) = (n - 6) \sum_{1 \leq i < j \leq n} \varphi(v_i + v_j) - (n^2 - 8n + 3) \sum_{i=1}^n \varphi(v_i) \tag{1.1}$$

in Random normed spaces.

We adopt the usual terminology, notions and conventions of the theory of random normed spaces as in [14].

Throughout, Δ^+ denotes the distribution functions spaces, i.e., the space of all mapping $f: \mathbb{R} \cup (-\infty, \infty) \rightarrow [0,1]$ such that f is left continuous and increasing on \mathbb{R} , $f(0)=0$ and $f(+\infty)=1$. D^+ subset of Δ^+ consisting of all functions V of Δ^+ for which $\ell^-V(+\infty)=1$ where $\ell^-f(s)$ denotes $\ell^-f(s)=\lim_{t \rightarrow s^-} f(t)$. The space Δ^+ is partially order by usual wise ordering of functions, i.e.,

$$f \leq g \Leftrightarrow f(s) = g(s), \forall s \in \mathbb{R}. \text{ The maximal element for } \Delta^+ \text{ in this order is the distribution function } \epsilon_0(s) = \begin{cases} 0, & \text{if } s \leq 0 \\ 1 & \text{if } s > 0 \end{cases}$$

Definition 1.1: A Random Normed space (RN-space) is a triple (V, μ, T) , where V is a vector space, T is a continuous t-norm and $\mu: V \rightarrow D^+$ satisfying the following conditions:

- (R₁). $\mu_v(t) = \epsilon_0(t)$ for all $t > 0$ if and only if $v=0$,
- (R₂). $\mu_{av}(t) = \mu_v(\frac{t}{|a|})$ for all $v \in V, t \geq 0$ and $a \in \mathbb{R}$ with $a \neq 0$,
- (R₃). $\mu_{v+u}(t + s) \geq T(\mu_v(t), \mu_u(s))$ for all $v, u \in V$ and $t, u \geq 0$.

Definition 1.2. Let (V, μ, T) be a RN- space.

- (i) A sequence $\{v_n\}$ in V is said to be convergent to $v \in V$ if $\lim_{n \rightarrow \infty} \mu_{v_n - v}(t) = 1, t > 0$.
- (ii) A sequence $\{v_n\}$ in V is said to be Cauchy sequence if $\lim_{n \rightarrow \infty} \mu_{v_n - v_m}(t) = 1, t > 0$.
- (iii) A RN-space (V, μ, T) is complete if every Cauchy sequence is convergent in V .

2 General Solution

Theorem 2.1. If any even function $\varphi: V \rightarrow W$ satisfies the functional equation (1.1) for all $v_1, v_2, \dots, v_n \in V$, then the function φ is quadratic.

Proof. Let function $\varphi: V \rightarrow W$ satisfies (1.1). Taking $v_1 = v_2 = \dots = v_n = 0$ in (1.1), we have $\varphi(0) = 0$. Replacing (v_1, v_2, \dots, v_n) by $(v, 0, \dots, 0)$ in (1.1), we get

$$\varphi(-2v) + (n - 1) \varphi(v) = (n - 6)(n-1) \varphi(v) - (n^2 - 8n + 3) \varphi(v) \tag{2.1}$$

$$\varphi(-2v) = (n^2 - 7n + 6 - n^2 + 8n - 3 - n + 1) \varphi(v) \tag{2.2}$$

$$\varphi(-2v) = 4\varphi(v) \tag{2.3}$$

Now taking $v_1=v_2=v$ and $v_3=\dots=v_n=0$ in (1.1), we have

$$2\phi(-v) + (n-2)\phi(2v) = (n-6)\phi(2v) + 2(n-6)(n-2)\phi(v) - 2(n^2 - 8n + 3)\phi(v)$$

$$2\phi(-v) + 4\phi(2v) = 18\phi(v) \tag{2.4}$$

Since ϕ is an even mapping, hence by (2.4) we have

$$\phi(2v) = 2^2\phi(v), \text{ for all } v \in V.$$

Now, replace v by $2v$, we have

$$\phi(2^2 v) = 2^4\phi(v)$$

Continue like this, we generalize

$$\phi(2^n v) = 2^{2n}\phi(v) \tag{2.5}$$

For all $v \in V$ and for any $n \geq 0$.

Similarly, we have

$$\phi(2^{-n} v) = 2^{-2n}\phi(v) \tag{2.6}$$

Remark 2.2: Let V be a linear space and a mapping $\phi: V \rightarrow W$ satisfies the functional equation (1.1), then

- (i) $\phi(r^n v) = r^n \phi(v)$ for all $v \in V, r \in \mathbb{Q}, n \in \mathbb{Z}$.
- (ii) $\phi(v) = v\phi(1)$ for all $v \in V$ if ϕ is continuous.

3 Hyers-Ulam Stability : Direct Method

For notational handiness, we denote (V, Ψ, T) , and (W, Ψ, T) are Complete RN spaces and define a mapping $\phi: V \rightarrow W$ by

$$D\phi(v_1, v_2, \dots, v_n) = \sum_{i=1}^n \phi(-2v_i + \sum_{j=1, i \neq j}^n v_j) - (n-6) \sum_{1 \leq i < j \leq n} \phi(v_i + v_j) + n^2 - 8n + 3i = 1n\phi(v_i) \tag{3.1}$$

for all $v_1, v_2, \dots, v_n \in V$.

Theorem 3.1. If an even mapping $\phi: V \rightarrow W$ with $\phi(0) = 0$ for which there exists a mapping $\Phi: V^n \rightarrow D^+$ for some $0 < \alpha < 4$,

$$\Phi_{2v_1, 2v_2, \dots, 2v_n}(\epsilon) \geq \Phi_{v_1, v_2, \dots, v_n} \left(\frac{\epsilon}{\alpha} \right) \tag{3.2}$$

$$\text{And } \lim_{t \rightarrow \infty} \Phi_{2^t v_1, 2^t v_2, \dots, 2^t v_n}(2^{2t}\epsilon) = 1 \tag{3.3}$$

for all $v_1, v_2, \dots, v_n \in V$ and all $\epsilon > 0$ such that

$$\mu_{D\phi(v_1, v_2, \dots, v_n)}(\epsilon) \geq \Phi_{v_1, v_2, \dots, v_n}(\epsilon). \tag{3.4}$$

Then there exists a unique quadratic mapping $Q_2: V \rightarrow W$ satisfying the functional equation (1.1) with

$$\mu_{Q_2(v) - \phi(v)}(\epsilon) \geq \Phi_{v, 0, \dots, 0}((2^2 - \alpha)2\epsilon) \tag{3.5}$$

for all $v \in V$ and all $\epsilon > 0$. The mapping $Q_2: V \rightarrow W$ is defined by

$$\mu_{Q_2(v)}(\varepsilon) = \lim_{t \rightarrow \infty} \mu_{\frac{\phi(2^t v)}{2^{2t}}}(\varepsilon) \tag{3.6}$$

for all $v \in V$ and all $\varepsilon > 0$.

Proof: Replace (v_1, v_2, \dots, v_n) by $(v, 0, \dots, 0)$ in (3.4), we have

$$\begin{aligned} \mu_{\phi(2v)-2^2\phi(v)}(\varepsilon) &\geq \Phi_{v,0,\dots,0}(\varepsilon). \\ \mu_{\frac{\phi(2v)}{2^2}-\phi(v)}(\varepsilon) &\geq \Phi_{v,0,\dots,0}(2^2\varepsilon). \end{aligned} \tag{3.7}$$

Replace v by $2^t v$ in (3.7) we have

$$\begin{aligned} \mu_{\frac{\phi(2^{t+1}v)}{2^2}-\phi(2^t v)}(\varepsilon) &\geq \Phi_{2^t v,0,\dots,0}(2^2\varepsilon) \\ \mu_{\frac{\phi(2^{t+1}v)}{2^{2(t+1)}}-\frac{\phi(2^t v)}{2^{2t}}}(\varepsilon) &\geq \Phi_{2^t v,0,\dots,0}(2^{2(t+1)}\varepsilon) \\ &\geq \Phi_{v,0,\dots,0}\left(\frac{2^{2(t+1)}\varepsilon}{\alpha^t}\right). \end{aligned} \tag{3.8}$$

for all $v \in V$ and all $\varepsilon > 0$. Since

$$\frac{\phi(2^m v)}{2^{2m}} - \phi(v) = \sum_{i=0}^{m-1} \frac{\phi(2^{i+1}v)}{2^{2(i+1)}} - \frac{\phi(2^i v)}{2^{2i}} \tag{3.9}$$

From (3.8) and (3.9), we have

$$\begin{aligned} \mu_{\frac{\phi(2^m v)}{2^{2m}}-\phi(v)}\left(\sum_{i=0}^{m-1} \frac{\alpha^i}{2^{2(i+1)}} \frac{\varepsilon}{2}\right) &\geq \Phi_{v,0,\dots,0}(\varepsilon). \\ \mu_{\frac{\phi(2^m v)}{2^{2m}}-\phi(v)}(\varepsilon) &\geq \Phi_{v,0,\dots,0}\left(\frac{2\varepsilon}{\sum_{i=0}^{m-1} \frac{\alpha^i}{2^{2(i+1)}}}\right) \end{aligned} \tag{3.9}$$

Replace v by $2^2 v$, we got

$$\mu_{\frac{\phi(2^{m+n}v)}{2^{2(m+n)}}-\frac{\phi(2^n v)}{2^{2n}}}(\varepsilon) \geq \Phi_{v,0,\dots,0}\left(\frac{2\varepsilon}{\sum_{i=n}^{m+n-1} \frac{\alpha^i}{2^{2(i+1)}}}\right)$$

For all $v \in V$ and all $\varepsilon > 0$. As $\Phi_{v,0,\dots,0}\left(\frac{2\varepsilon}{\sum_{i=n}^{m+n-1} \frac{\alpha^i}{2^{2(i+1)}}}\right) \rightarrow 1$ as $m, n \rightarrow \infty$ then $\left\{\frac{\phi(2^n v)}{2^{2n}}\right\}$ is a Cauchy sequence in (W, μ, T) , Since (W, μ, T) is complete RN- space, thus sequence $\left\{\frac{\phi(2^n v)}{2^{2n}}\right\}$ converges to some $Q_2(v) \in W$. fix $v \in V$ and put $n=0$, we obtain

$$\mu_{\frac{\phi(2^m v)}{2^{2m}}-\phi(v)}(\varepsilon) \geq \Phi_{v,0,\dots,0}\left(\frac{2\varepsilon}{\sum_{i=0}^{m-1} \frac{\alpha^i}{2^{2(i+1)}}}\right) \tag{3.10}$$

And so, for every $\delta > 0$, we get

$$\mu_{Q_2(v)-\phi(v)}(\varepsilon + \delta) \geq T\left(\mu_{\frac{\phi(2^m v)}{2^{2m}}-\phi(v)}(\delta), \mu_{\frac{\phi(2^m v)}{2^{2m}}-\phi(v)}(\varepsilon)\right)$$

$$\geq T \left(\mu_{Q_2(v)-\phi(2^m v)}(\delta), \Phi_{v,0,\dots,0} \left(\frac{2\varepsilon}{\sum_{i=0}^{m-1} \frac{\alpha^i}{2^{2(i+1)}}} \right) \right)$$

For all $v \in V$ and all $\varepsilon, \delta > 0$. Taking the limit $m \rightarrow \infty$,

$$\mu_{Q_2(v)-\phi(v)}(\varepsilon + \delta) \geq \Phi_{v,0,\dots,0}(2(2^2 - \alpha)\varepsilon) \tag{3.11}$$

For all $v \in V$ and all $\varepsilon, \delta > 0$. Since δ was arbitrary, taking $\delta \rightarrow 0$,

$$\mu_{Q_2(v)-\phi(v)}(\varepsilon) \geq \Phi_{v,0,\dots,0}(2(2^2 - \alpha)\varepsilon) \tag{3.12}$$

For all $v \in V$ and all $\varepsilon > 0$. Replacing (v_1, v_2, \dots, v_n) by $(2^t v_1, 2^t v_2, \dots, 2^t v_n)$ in (3.4),

$$\mu_{DQ_2(2^t v_1, 2^t v_2, \dots, 2^t v_n)}(\varepsilon) \geq \Phi_{2^t v_1, 2^t v_2, \dots, 2^t v_n}(2^{2t}\varepsilon). \tag{3.13}$$

Since $\lim_{t \rightarrow \infty} \Phi_{2^t v_1, 2^t v_2, \dots, 2^t v_n}(2^{2t}\varepsilon) = 1$, so Q_2 satisfies the functional equation (1.1). To prove the uniqueness of Quadratic mapping Q_2 . Assume that there exists another Quadratic mapping Q'_2 , which satisfies inequality (3.11). Fix $v \in V$. Clearly, $Q_2(2^t v) = 2^{2t}Q_2(v)$ and $Q'_2(2^t v) = 2^{2t}Q'_2(v)$ for all $v \in V$. from (3.10), we have

$$\begin{aligned} \mu_{Q_2(v)-Q'_2(v)}(\varepsilon) &= \mu_{\frac{Q_2(2^t v)}{2^{2t}} - \frac{Q'_2(2^t v)}{2^{2t}}}(\varepsilon) \\ &\geq T \left(\mu_{\frac{Q_2(2^t v)}{2^{2t}} - \frac{\phi(2^t v)}{2^{2t}}}\left(\frac{\varepsilon}{2}\right), \mu_{\frac{Q'_2(2^t v)}{2^{2t}} - \frac{\phi(2^t v)}{2^{2t}}}\left(\frac{\varepsilon}{2}\right) \right) \\ &\geq \Phi_{v,0,\dots,0} \left(\left(\frac{2^2 - \alpha}{\alpha}\right)^t (2^2 - \alpha)4\varepsilon \right) \end{aligned} \tag{3.14}$$

$\lim_{t \rightarrow \infty} 4(2^2 - \alpha) \left(\frac{3}{\alpha}\right)^t \varepsilon = \infty$, we have $\mu_{Q(v)-Q'(v)}(\varepsilon) = 1$ for all $\varepsilon > 0$. Thus, $Q_2(v) = Q'_2(v)$, for all $v \in V$. Hence, the proof is complete.

Theorem 3.2. If an even mapping $\phi: V \rightarrow W$ with $\phi(0) = 0$ for which there exists a mapping $\Phi: V^n \rightarrow D^+$ for some $0 < \alpha < 4$,

$$\Phi_{\frac{v_1 v_2}{2}, \dots, \frac{v_n}{2}}(\varepsilon) \geq \Phi_{v_1, v_2, \dots, v_n}(\alpha\varepsilon) \tag{3.15}$$

$$\text{And } \lim_{t \rightarrow \infty} \Phi_{\frac{v_1 v_2}{2^{2t}}, \dots, \frac{v_n}{2^t}}\left(\frac{\varepsilon}{2^{2t}}\right) = 1 \tag{3.16}$$

for all $v_1, v_2, \dots, v_n \in V$ and all $\varepsilon > 0$ such that

$$\mu_{D\phi(v_1, v_2, \dots, v_n)}(\varepsilon) \geq \Phi_{v_1, v_2, \dots, v_n}(\varepsilon). \tag{3.17}$$

Then there exists a unique quartic mapping $Q_2: V \rightarrow W$ satisfying the functional equation (1.1) with

$$\mu_{Q_2(v)-\phi(v)}(\varepsilon) \geq \Phi_{v,0,\dots,0}\left(\frac{\varepsilon}{(2^2 - \alpha)2}\right) \tag{3.18}$$

for all $v \in V$ and all $\varepsilon > 0$. The mapping $Q_2: V \rightarrow W$ is defined by

$$\mu_{Q_2(v)}(\varepsilon) = \lim_{t \rightarrow \infty} \mu_{2^{2t}\phi\left(\frac{v}{2^t}\right)}(\varepsilon) \tag{3.19}$$

for all $v \in V$ and all $\varepsilon > 0$.

Corollary 3.3. If an even mapping $\phi: V \rightarrow W$ with $\phi(0) = 0$ for which there exists a mapping $\Phi: V \rightarrow D^+$ satisfying

$$\mu_{D\phi(v_1, v_2, \dots, v_n)}(\varepsilon) \geq \Phi_{\sum_{i=1}^n \|v_i\|^\theta}(\varepsilon). \quad (3.20)$$

Then there exists a unique quadratic mapping $Q_2: V \rightarrow W$ satisfying the functional equation (1.1) with

$$\mu_{Q_2(v) - \phi(v)}(\varepsilon) \geq \Phi_{\|v\|^\theta}((2^2 - 2^\theta)2\varepsilon) \quad (3.21)$$

for all $v \in V$, where $\theta < 2$ and all $\varepsilon > 0$

We take $\alpha = 2^{\theta-2}$ and $v_1, v_2, \dots, v_n = \sum_{i=1}^n \|v_i\|^\theta$ in the theorem 3.1.

4 Conclusion

In this paper a new type quadratic functional equation is introduced and proved its stability in Random Normed space. The stability of the equation can be proved in other spaces like Fuzzy normed space, modular space, non-Archimedean normed space etc.

Competing Interests

Author has declared that no competing interests exist.

References

- [1] Ulam SM. Problems in modern mathematics. Science Editions, John Wiley & Sons, Inc., New York; 1964.
- [2] Hyers DH. On the stability of the linear functional equation. Proceedings of the National Academy of Sciences of the United States of America. 2005;27(4):222–224.
- [3] Aoki T. On the stability of the linear transformation in Banach spaces. Journal of the Mathematical Society of Japan. 1950;2(10):64–66.
- [4] Rassias JM. On approximately of linear mappings by linear mappings. J. Funct. Anal. USA. 1982;46:126–130.
- [5] Rassias TM. On the stability of the linear mapping in Banach spaces. Proceedings of the American Mathematical Society. 1978;72(2):297–300.
- [6] Gavruta P. A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings. Journal of Mathematical Analysis and Applications. 1994;184(3):431–436.
- [7] Gajda Z. On stability of additive mappings. Intern. J. Math. Math. Sci. 1991;14:431–434.
- [8] Jin SS, Lee YH. On the stability of the functional equation deriving from quadratic and additive function in random normed spaces via fixed point method, J. Chungcheong Math. Soc. 2012;25:51–63.
- [9] Jung SM. Hyers-Ulam-rassias stability of functional equations in nonlinear analysis, Springer Optimization and Its Applications, Springer, New York, NY, USA; 2011.

- [10] Jung, S.M. and Sahoo P.K., “Hyers-Ulam stability of the quadratic equation of Pexider type,” Journal of the Korean Mathematical Society, 2001, 38(3), 645–656.
- [11] Khamsi MA. Fixed point theory in modular function spaces. Recent advances on metric fixed point theory, Univ. Sevilla, Seville, Ciencias. 1995;48:31–57.
- [12] Rassias Th. M, Shibata K. Variational problem of some quadratic functionals in complex analysis, J. Math. Anal. Appl. 1998;228:234–253.
- [13] Tamilvanan K, Lee JR, Park C. Ulam stability of a functional equation deriving from quadratic and additive mappings in random normed spaces, AIMS Math. 2020;6:908—924.
- [14] Alessa N, Tamilvanan K, Balasubramanian G, Loganathan K. Stability results of the functional equation deriving from quadratic function in random normed spaces, AIMS mathematics. 2021;6(3):2385-2397.
- [15] Rassias Th. M. On the stability of the quadratic functional equation and its applications, Studia Univ. Babeş-Bolyai. 1998;89–124.

© 2023 Devi; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<https://www.sdiarticle5.com/review-history/98066>