



Evaluation of Antioxidant and Xanthine Oxidase Inhibitory Potential of Methanolic Root Extract of *Acorus calamus*- an *In vitro* Study

**Ashfaaq Ahmed ^a, R. Gayathri ^{a*#}, V. Vishnu Priya ^{a†}, J. Selvaraj ^{a#}
and S. Kavitha ^{a‡}**

^a Department of Biochemistry, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai-600 077, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i60B34799

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/81566>

Original Research Article

Received 15 October 2021

Accepted 20 December 2021

Published 23 December 2021

ABSTRACT

Background: Gout is an inflammatory joint disease that elevates the uric acid levels in blood that triggers the formation of urate crystals in the joint, especially since the past 50 years. Xanthine oxidase catalyses oxidative hydroxylation of hypoxanthine to Xanthine to uric acid. show it's higher in men aged above 50 years. *Acorus calamus* is a mid term, perennial, fragrant herb. Therefore, the main approach for the treatment of gout is reducing uric acid.

Aim: To analyze the antioxidant and Xanthine Oxidase inhibitory potential of methanolic root extract of *Acorus calamus*.

Materials and Methods: Preparation of methanolic root extract of *Acorus calamus* was done by hot percolation method. Phytochemical screening test was done. The antioxidant activity was carried out by DPPH radical scavenging assay. Anti gout potential of the herbal extract was analysed by the evaluation of Xanthine oxidase inhibitory potential. The data were analyzed statistically by a one-way analysis of variance (ANOVA) followed by Duncan's multiple range test to see the statistical significance among the groups. The results with $p < 0.05$ level were considered to be statistically significant.

[#]Associate Professor,

[†]Professor,

[‡]Lecturer,

*Corresponding author: E-mail: gayathri.sdc@saveetha.com;

Results: Methanolic root extract of *Acorus calamus* shows a strong presence of phytochemicals such as alkaloids, terpenoids, steroids, saponins, flavonoids. IC₅₀ of antioxidant potential of Methanolic root extract of *Acorus calamus* was found to be 210 µg/ml. IC₅₀ of xanthine oxidase inhibitory potential of methanolic root extract of *Acorus calamus* was found to be 310 µg/ml.

Conclusion: Methanolic root extract of *Acorus calamus* exhibited significant antioxidant and anti-gout potential. Further research on the natural Xanthine oxidase inhibitors especially *in vivo* studies and investigation of active compounds and its pharmacokinetics to be elucidated.

Keywords: Gout; Xanthine oxidase; *Acorus calamus*; pharmacokinetics; DPPH; Innovative technology; Novel method.

1. INTRODUCTION

Acorus calamus also known as sweet flag is a mid-term, perennial, fragrant herb. The plant's rhizomes are brown in colour, twisted, cylindrical, curved and shortly noded, with radiant green leaves, a sword-like structure, which has curly margins and is thick in the middle [1,2]. It is a tall wetland monocot from the family Acoraceae and species *Acorus* [3].

Gout is a chronic disorder or an inflammatory joint disease that elevates the uric acid levels in blood that triggers the formation of monosodium urate crystals in the joint. It can also cause tophi, joint deformities and kidney stones [4]. Its occurrence has increased since the past 50 years, especially in developing countries [5] only 5% of the individuals who had hyperuricemia above 9 mg/dL developed gout. Accordingly, it is proved that the incidence of gout is shared by factors such as genetic predisposition. In India, approximately 0.3% of the population is affected by gout, and the statistics prove that it's higher in men aged above 50 years [6].

Xanthine oxidase is an enzyme that generates reactive oxygen species that catalyses oxidative hydroxylation of hypoxanthine and xanthine to uric acid in further oxidation, leading to inflammation at joints that causes severe pain, redness and soreness [7]. However, it forms uricase which is not a functional human enzyme, as a result people can develop hyperuricemia [8].

Hyperuricemia is the key predictor for development of gout [1] it occurs when serum uric acid levels are more than 0.42 mmol/L and the main cause is due to the unbalanced excretion and production of uric acids, therefore the main approach for treating gout is by reducing the uric acid production [1].

Recently it has been reported that treating gout using medicinal plants is gaining new interest [9], due to its less side effects and lower cost [3]. The synthetic drugs used in the treatment of gout are reported to have various side effects and cannot be used for a longer period of time [10]. Our team has extensive knowledge and research experience that has translated into high quality publications [11-30]. The aim of this current research is to validate the *in vitro* antioxidant and anti-gout potential of methanolic root extract of *Acorus calamus*.

2. MATERIALS AND METHODS

2.1 Preparation of Methanolic Root Extract of *Acorus calamus*

Acorus calamus was purchased from a herbal health care centre. Air dried, crushed and made into powder form. Methanol was added to it. 80% of methanolic extract was obtained. The extract was then prepared by a hot percolation method. Later it was dried and used to analyze the antioxidant and anti-inflammatory potential [31].

2.2 DPPH radical Scavenging Activity

Scavenging of 2, 2-Diphenyl-1-picrylhydrazyl (DPPH) radicals was assessed by the method of Hatano et al (1989). DPPH solution (1.0 ml) was added to 1.0 ml of extract at different concentrations (0.1 to 0.5 mg/ml). The mixture was kept at room temperature for 50 minutes and the activity was measured at 517 nm. Same concentration of ascorbic acid was used as standard. The capability to scavenge the DPPH radical was calculated and expressed in percentage (%) using following formula:

$$\text{DPPH radical scavenging (\%)} = \frac{\text{Control OD} - \text{Sample OD}}{\text{Control OD}} \times 100$$

In vitro anti-inflammatory activity of methanolic root extract of *Acorus calamus* by Xanthine oxidase inhibitory activity: The anti-inflammatory activity of methanolic root extract of *Acorus calamus* was studied by using inhibition of albumin denaturation technique which was studied according to the method of Leela Prakash and Mohan Dass, (2010). The reaction mixture consisted of 1% aqueous solution of bovine albumin fraction and test extracts, the reaction mixture's pH was adjusted using a little amount of 1N HCl. The sample extracts were incubated at 37 °C for 20 min and then heated to 51°C for 20 min, after cooling the samples the turbidity was measured at 660nm. (UV Visible Spectrophotometer Model 371, Elico India Ltd). The experiment was performed in triplicate. The Percentage inhibition of protein denaturation was calculated as follows:

$$\text{Percentage inhibition(\%)} = (\text{Abs Control} - \text{Abs Sample}) \times 100 / \text{Abs control}$$

2.3 Statistical Analysis

The triplicate analysis results of the experiments performed on control and experimented *Acorus calamus* were expressed as mean \pm standard deviation. Results were analyzed statistically by a two-way analysis of variance (ANOVA) and significant differences between the mean values were measured using Tukey's multiple range test using Graph Pad Prism version 5. The results with the $p < 0.05$ level were considered to be statistically significant.

3. RESULTS AND DISCUSSION

From the above study the methanolic root extract of *Acorus calamus* shows a presence of phytochemicals such as alkaloids, terpenoids, steroids, saponins, flavonoids (Table 1). Phytochemicals are secondary metabolites which are exclusively present in plants. The presence of these phytochemicals are responsible for the medicinal property of these plant extracts [32].

The antioxidant activity of *Acorus Calamus* was determined by performing the DPPH radical scavenging assay. They are molecules possessing unpaired electrons emerging from phenolic compounds. Plant extract with phenolic phytochemicals can scavenge the free radicals and thus the antioxidant potential can be estimated [6]. Methanolic root extract of *Acorus calamus* exhibited a significant antioxidant

potential and increased in a dose dependent manner as compared to the standard. IC_{50} of Methanolic root extract of *Acorus calamus* was found to be 210 $\mu\text{g/ml}$ (Fig. 1).

Table 1. Phytochemical screening of methanolic root extract of *Acorus calamus*

Serial no.	Phytochemicals	Presence
1	Amino acids	++
2	Protein	+
3	Alkaloids	-
4	Terpenoids	+
5	Steroids	+
6	Carbohydrates	-
7	Saponins	+
8	Flavonoids	++

A dose dependent Xanthine oxidase inhibitory activity (anti-gout) was observed for the extract in the present study, the standard drug allopurinol showed greater percent of inhibition of Xanthine oxidase than the extract of the compounds with some concentration [33]. IC_{50} of xanthine oxidase inhibitory potential of methanolic root extract of *Acorus calamus* was found to be 310 $\mu\text{g/ml}$ (Graph 2) Gout is a disorder whose prevalence has increased in the last 50 years. Reasons for the disorder can be genetic, high living and excess alcohol consumption. The current treatment for gout predominantly depends on synthetic drugs with serious side effects [34].

Prolonged usage of synthetic drugs can lead to adverse side effects and now research is focussed to explore the rich phytochemicals of herbal extracts which are a part of indigenous medicine [35]. More research and awareness has to be spread to cure the disorder instead of just managing one. Rawlani et al (2014) has stated that conventional radiography is an effective diagnostic method to treat chronic tophaceous gout and ultrasonography has been used recently to assist in both diagnosing and monitoring the disease [36]. Rübenthaler et al has stated that utilizing computed tomography is associated with ionizing radiation, with a little added benefit over ultrasonography and computed tomography, Reiser and Clevert et al stated that magnetic resonance imaging is a great modality to image bones and soft tissues as it has less radiation and better quality [37]. The current common clinical treatments for gout are mainly self medication and anti-inflammatory medications (NSAIDs) by [38].

Antioxidant potential of methanolic root extract of *Acorus calamus*




Fig. 1. Represents antioxidant potential of methanolic root extract of *Acorus calamus* by DPPH Assay against the standard Ascorbic acid.. X-axis represents the concentration in µg/ml while Y-axis represents the inhibitory potential of the extracts. Blue bar represents the standard - Ascorbic acid, purple bar represents methanolic root extract of *Acorus calamus*. Each bar represents Mean ± SEM of 3 independent observations. Significance at $p < 0.05$

Xanthine oxidase inhibitory potential of methanolic root extract of *Acorus calamus*




Fig. 2. Represents anti-inflammatory activity - Xanthine oxidase inhibitory potential of *Acorus calamus* root ethanolic extract compared with the standard - Allopurinol. X-axis represents the concentration in µg/ml and Y-axis represents the inhibitory potential of the extracts. Green bar represents the standard -Allopurinol, purple bar represents methanolic root extract of *Acorus calamus*. Each bar represents Mean ± SEM of 3 independent observations. Significance at $p < 0.05$

4. CONCLUSION

The bioactive compounds found in medicinal plants remain as an important component of research for the development of new drugs with potential to fight against several diseases and disorders. Methanolic root extract of *Acorus calamus* shows significant antioxidant and anti-

gout properties as compared to the standards. Hence it can be used as an antioxidant and anti gout drug for the management of various health ailments. Further studies on the *in vitro* cell line and *in vivo* experimental models need to be carried out in order to ascertain its potential mechanisms of action towards the development of clinical utility.

SOURCE OF FUNDING

The present study was supported by the following agencies.

- Saveetha Dental College
- Saveetha Institute of Medical and Technical Sciences (SIMATS)
- Saveetha University
- Indo Vietnam Association, Pondicherry

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

ACKNOWLEDGEMENT

I would like to acknowledge the support of our guide Dr. Gayathri R and Dr. Selvaraj J and Saveetha dental college, Chennai and Department of biochemistry for providing the research units for carrying out this research.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Zhang W-Z. Why Does Hyperuricemia Not Necessarily Induce Gout? *Biomolecules* [Internet]. 2021;11(2). Available: <http://dx.doi.org/10.3390/biom11020280>
2. *Acorus calamus* Linn [Internet]. SpringerReference. Available: http://dx.doi.org/10.1007/springerreference_67975
3. Altaf M, Manoharadas S, Zeyad MT. Green synthesis of cerium oxide nanoparticles using *Acorus calamus* extract and their antibiofilm activity against bacterial pathogens. *Microsc Res Tech* [Internet]; 2021. Available: <http://dx.doi.org/10.1002/jemt.23724>
4. Choi N, Yang G, Jang JH, Kang HC, Cho Y-Y, Lee HS, et al. Loganin Alleviates Gout Inflammation by Suppressing NLRP3 Inflammasome Activation and Mitochondrial Damage. *Molecules* [Internet]. 2021;26(4). Available: <http://dx.doi.org/10.3390/molecules26041071>
5. Kapoor N, Saxena S. Xanthine oxidase inhibitory and antioxidant potential of Indian Muscodor species [Internet]. *3 Biotech*. 2016;6. Available: <http://dx.doi.org/10.1007/s13205-016-0569-5>
6. Dhiman P, Malik N, Khatkar A, Kulharia M. Antioxidant, Xanthine Oxidase and Monoamine Oxidase Inhibitory Potential of Coumarins: A Review [Internet]. *Current Organic Chemistry*. 2017;21:294–304. Available: <http://dx.doi.org/10.2174/1385272820666161021103547>
7. Wakabayashi T, Ueno S, Nakatsuji T, Hirai T, Niinomi I, Oyama S, et al. Safety profiles of new xanthine oxidase inhibitors: A post-marketing study. *Int J Clin Pharmacol Ther* [Internet]. 2021;9; Available: <http://dx.doi.org/10.5414/CP203898>
8. Port FD, Du Port F. The Signs and Causes of Gout [Internet]. *The Decade of Medicine or The Physician of the Rich and the Poor*. 1988;80–80. Available: http://dx.doi.org/10.1007/978-3-642-73715-2_127
9. Wetherbee DK. Tokens of *Acorus calamus*: Walt Whitman's "Root of Wash'd Sweet-Flag. 1976;48.
10. Kamatani N. Febuxostat ; a non-purine xanthine oxidase inhibitor useful for the treatment of gout and hyperuricemia [Internet]. *Gout and Nucleic Acid Metabolism*. 2011;35:210. Available: <http://dx.doi.org/10.6032/gnam.35.210>
11. Wu F, Zhu J, Li G, Wang J, Veeraraghavan VP, Krishna Mohan S, et al. Biologically synthesized green gold nanoparticles from Siberian ginseng induce growth-inhibitory effect on melanoma cells (B16). *Artif Cells Nanomed Biotechnol*. 2019;47(1):3297–305.
12. Chen F, Tang Y, Sun Y, Veeraraghavan VP, Mohan SK, Cui C. 6-shogaol, a active constituents of ginger prevents UVB radiation mediated inflammation and oxidative stress through modulating Nrf2 signaling in human epidermal keratinocytes (HaCaT cells). *J Photochem Photobiol B*. 2019;197:111518.

13. Li Z, Veeraraghavan VP, Mohan SK, Bolla SR, Lakshmanan H, Kumaran S, et al. Apoptotic induction and anti-metastatic activity of eugenol encapsulated chitosan nanopolymer on rat glioma C6 cells via alleviating the MMP signaling pathway [Internet]. *Journal of Photochemistry and Photobiology B: Biology*. 2020;203:111773. Available:<http://dx.doi.org/10.1016/j.jphoto-biol.2019.111773>
14. Babu S, Jayaraman S. An update on β -sitosterol: A potential herbal nutraceutical for diabetic management. *Biomed Pharmacother*. 2020;131:110702.
15. Malaikolundhan H, Mookkan G, Krishnamoorthi G, Matheswaran N, Alsawalha M, Veeraraghavan VP, et al. Anticarcinogenic effect of gold nanoparticles synthesized from *Albizia lebbek* on HCT-116 colon cancer cell lines. *Artif Cells Nanomed Biotechnol*. 2020;48(1):1206–13.
16. Han X, Jiang X, Guo L, Wang Y, Veeraraghavan VP, Krishna Mohan S, et al. Anticarcinogenic potential of gold nanoparticles synthesized from *Trichosanthes kirilowii* in colon cancer cells through the induction of apoptotic pathway. *Artif Cells Nanomed Biotechnol*. 2019;47(1):3577–84.
17. Gothai S, Muniandy K, Gnanaraj C, Ibrahim IAA, Shahzad N, Al-Ghamdi SS, et al. Pharmacological insights into antioxidants against colorectal cancer: A detailed review of the possible mechanisms. *Biomed Pharmacother*. 2018;107:1514–22.
18. Veeraraghavan VP, Hussain S, Balakrishna JP, Dhawale L, Kullappan M, Ambrose JM, et al. A Comprehensive and Critical Review on Ethnopharmacological Importance of Desert Truffles: *Terfezia claveryi*, *Terfezia boudieri*, and *Tirmania nivea* [Internet]. *Food Reviews International*. 2021;1–20. Available:<http://dx.doi.org/10.1080/87559129.2021.1889581>
19. Sathya S, Ragul V, Veeraraghavan VP, Singh L, Niyas Ahamed MI. An in vitro study on hexavalent chromium [Cr(VI)] remediation using iron oxide nanoparticles based beads. *Environmental Nanotechnology, Monitoring & Management*. 2020;14:100333.
20. Yang Z, Pu M, Dong X, Ji F, Priya Veeraraghavan V, Yang H. Piperine loaded zinc oxide nanocomposite inhibits the PI3K/AKT/mTOR signaling pathway via attenuating the development of gastric carcinoma: In vitro and in vivo studies. *Arabian Journal of Chemistry*. 2020;13(5):5501–16.
21. Rajendran P, Alzaharani AM, Rengarajan T, Veeraraghavan VP, Krishna Mohan S. Consumption of reused vegetable oil intensifies BRCA1 mutations. *Crit Rev Food Sci Nutr*. 2020;1–8.
22. Barma MD, Muthupandian I, Samuel SR, Amaechi BT. Inhibition of *Streptococcus mutans*, antioxidant property and cytotoxicity of novel nano-zinc oxide varnish. *Arch Oral Biol*. 2021;126:105132.
23. Samuel SR. Can 5-year-olds sensibly self-report the impact of developmental enamel defects on their quality of life? *Int J Paediatr Dent*. 2021;31(2):285–6.
24. Samuel SR, Kuduruthullah S, Khair AMB, Shayeb MA, Elkaseh A, Varma SR. Dental pain, parental SARS-CoV-2 fear and distress on quality of life of 2 to 6 year-old children during COVID-19. *Int J Paediatr Dent*. 2021;31(3):436–41.
25. Tang Y, Rajendran P, Veeraraghavan VP, Hussain S, Balakrishna JP, Chinnathambi A, et al. Osteogenic differentiation and mineralization potential of zinc oxide nanoparticles from *Scutellaria baicalensis* on human osteoblast-like MG-63 cells [Internet]. *Materials Science and Engineering: C*. 2021;119:111656. Available:<http://dx.doi.org/10.1016/j.msec.2020.111656>
26. Yin Z, Yang Y, Guo T, Veeraraghavan VP, Wang X. Potential chemotherapeutic effect of betalain against human non-small cell lung cancer through PI3K/Akt/mTOR signaling pathway. *Environ Toxicol*. 2021;36(6):1011–20.
27. Veeraraghavan VP, Periadurai ND, Karunakaran T, Hussain S, Surapaneni KM, Jiao X. Green synthesis of silver nanoparticles from aqueous extract of *Scutellaria barbata* and coating on the cotton fabric for antimicrobial applications and wound healing activity in fibroblast cells (L929). *Saudi J Biol Sci*. 2021;28(7):3633–40.
28. Mickyamaray S, Alfaiz FA, Paramasivam A, Veeraraghavan VP, Periadurai ND, Surapaneni KM, et al. Rhaponticin suppresses osteosarcoma through the inhibition of PI3K-Akt-mTOR pathway. *Saudi J Biol Sci*. 2021;28(7):3641–9.

29. Teja KV, Ramesh S. Is a filled lateral canal – A sign of superiority? [Internet]. Journal of Dental Sciences. 2020;15:562–3. Available:<http://dx.doi.org/10.1016/j.jds.2020.02.009>
30. Kadanakuppe S, Hiremath S. Social and Behavioural Factors Associated with Dental Caries Experience among Adolescent School Children in Bengaluru City, India [Internet]. British Journal of Medicine and Medical Research. 2016;14:1–10. Available:<http://dx.doi.org/10.9734/bjmmr/2016/24021>
31. Worthington OJ. The Phlobatannin of Slash Pine (*Pinus Caribaea, Morelet*). 1936;252 .
32. Fagugli RM, Gentile G, Ferrara G, Brugnano R. Acute renal and hepatic failure associated with allopurinol treatment. Clin Nephrol. 2008;70(6):523–6.
33. Ranjana, Ranjana, Nooreen Z, Bushra U, Jyotshna, Bawankule DU, et al. Standardization and xanthine oxidase inhibitory potential of Zanthoxylum armatum fruits [Internet]. Journal of Ethnopharmacology. 2019;230:1–8. Available:<http://dx.doi.org/10.1016/j.jep.2018.10.018>
34. Nanda BL. Determination of Phytochemicals and Antioxidant Activity of Acorus Calamus Rhizome [Internet]. Vol. 4, Journal of Drug Delivery and Therapeutics;2014. Available:<http://dx.doi.org/10.22270/jddt.v4i6.1005>
35. Roberts CJ. The Effect of Prolonged Drug Usage on Fetal Development. An Epidemiological Approach [Internet]. Advances in Experimental Medicine and Biology. 1972;457–65. Available:http://dx.doi.org/10.1007/978-1-4684-3219-0_39
36. Rawlani S, Rawlani S. Conventional Radiography [Internet]. Oral and Maxillofacial Imaging Techniques. 2014; 5–5. Available:http://dx.doi.org/10.5005/jp/books/12342_2
37. Rübenthaler J, Reiser M, Clevert D-A. Diagnostic vascular ultrasonography with the help of color Doppler and contrast-enhanced ultrasonography [Internet]. Ultrasonography. 2016;35:289–301. Available:<http://dx.doi.org/10.14366/usg.16027>
38. Terkeltaub R. Gout. Novel therapies for treatment of gout and hyperuricemia [Internet]. Arthritis Research & Therapy. 2009;11:236. Available: <http://dx.doi.org/10.1186/ar2738>

© 2021 Ahmed et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/81566>