



Willingness of Voluntary Counseling and Testing Among Long Distance Truck Drivers: A Window of Opportunity in Control of Retroviral Disease

Arodiwe I. O. ^{a*} and Arodiwe E. B. ^b

^a Department of Paediatrics, College of Medicine, University of Nigeria, Enugu Campus, Enugu, Nigeria.

^b Department of Medicine, College of Medicine, University of Nigeria, Enugu Campus, Enugu, Nigeria.

Authors' contributions

This work was carried out in collaboration between both authors. Author AIO conceived the paper, did data collection, conducted data analysis, wrote manuscript and approved final version. Author AEB conducted the data analysis, wrote the manuscript and approved the final version. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AIR/2023/v24i6990

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/106834>

Original Research Article

Received: 26/07/2023

Accepted: 02/10/2023

Published: 12/10/2023

ABSTRACT

Objective: To evaluate the factors influencing the practice of voluntary counselling and testing among long-distance truck drivers as it affects the spread of retroviral disease.

Design: This was a quantitative cross-sectional study conducted in the major bus terminals in Enugu State, Nigeria. A questionnaire-based tool was used to collect data from a population of adult bus drivers.

*Corresponding author: E-mail: ljeoma.arodiwe@unn.edu.ng;

Setting: It was conducted at the 3 main motor parks in Enugu metropolis from January 2019 – March 2020.

Participants: 500 long-distance truck drivers aged 19-65 years.

Intervention: The knowledge and practices of the participants were assessed using a semi-structured interview guide. They were interviewed on various aspects of HIV/AIDS.

Main Outcome Measures: Knowledge and practices. On Knowledge and practice performance scale rating; < 50% is poor, 50 -75% is fair, and > 75% is good.

Results: Among the five hundred (500) long-distance truck drivers (LDTD) studied 68.0% showed a fair knowledge of HIV/AIDS. The prevalent high-risk behaviour was multiple sexual partners (73.2%). The practice of voluntary counselling and testing was poor, as (17.0%) had practised it; however, their risk perception was high at (76.0%). Fortunately, there was also a high willingness to screen 302 (60.4%) which was significantly higher in those with a tertiary level of education (100%) and the married participants (75.8 %) ($\chi^2=174.4$, $p < 0.01$).

Conclusion: There was a high-risk perception in this study population with a high willingness to be tested. We hereby recommend regular screening service at the motor packs of truck drivers in Enugu State.

Keywords: HIV testing; willingness; truck drivers; Enugu.

1. INTRODUCTION

Human Immune Deficiency Virus (HIV) infection and Acquired Immune Deficiency Syndrome (AIDS) have become pandemic and one of the challenges facing African countries [1-3]. It started in Nigeria in the 1980s [4] and is now a generalized epidemic, with prevalence above 1%. Enugu State in which the study was carried has an HIV prevalence of 5.5% [4]. Certain occupations have been associated with an increased risk of acquiring HIV/AIDS and long-distance truck drivers (LDTD) is one such [5,6]. Away for a prolonged period from the comfort of their wives, families and social environments; they engage in risky sexual behaviour, such as regular sex with both female and male commercial sex workers (CSWs), including multiple sex partners and drug abuse [5,6]. Studies have shown that they are most vulnerable to HIV/AIDS and constitute a risk for HIV transmission [6,7].

A study carried out by the USAIDS⁷ indicates that HIV infection prevalence stands at 1-2% among female sex workers and long-distance truck drivers. The prevalence was found to be 10% among injecting drug users and this practice of injecting drugs was also high among long-distance truck drivers [7].

A behavioural surveillance survey conducted in Nigeria in 2014 [8] reported that only 13% had ever been tested for HIV. Some studies have also been done on the attitude and practice of HIV Counseling and Testing (HCT) among LDTD in our environment [9-12] however their willingness to screen was not reported.

HIV Counseling and Testing (HCT) is one of the most important tools for prevention [13-15]. It is a process by which an individual undergoes counselling to enable him/her to make an informed decision about being tested for HIV. It is entirely voluntary or provider-initiated and confidential [16,17]. However, in many countries, less than 50% of the vulnerable groups patronize these services. The reasons are fear of a possible positive result with its associated stigma and discrimination and ignorance of the existence of HCT service [18-20].

Therefore this study aims to evaluate the practice of voluntary counseling and willingness to screen for HIV infection among LDTD. This will help gain insight into their lifestyle that may be related to the spread of the disease and how to mitigate it.

2. MATERIALS AND METHODS

This was a descriptive cross-sectional study of long-distance truck drivers, conducted from January 2019 – March 2020 at the three main parks namely, HolyGhost Luxury Bus Park, 9th Mile Corner and Ogbete main market park. These are very busy truck terminals. These are well known for their transportation business and a link to other neighbouring towns in Enugu State. Good networks of roads connect important centres of trade and industry in the State and are linked between the South-East geo-political zone and the middle belt and beyond.

There was consecutive recruitment of LDTD, who gave their consent. The objectives of the

study were explained to them and confidentiality was assured by the non-inclusion of self-identification in the questionnaire. Documentation was made of their socio-demographic characteristics and their Knowledge of HIV/AIDS, Attitude to risky sexual behaviour and practice of HCT. Four assistants were recruited and trained on methods of data administration and collection. The questionnaire was checked and sorted out manually by the author and the data obtained was analyzed using SPSS statistical software package Version 25.0. The analysis consisted of tabulations and a Chi-square test to examine relationships. Statistical significance was considered present when the value was less than 0.05. Confidence Interval was 95%. On attitude and practice level, < 50% was rated as poor, 50 - 75% as fair, and >75% as good.

3. RESULTS

A total of five hundred (500) LDTD respondents were recruited for the study. They were all males. Table 1 shows that most of the respondents 163 (32.6%) were within the age group 30-39 years. All of them were males. With mean age of respondents = 38.3 ± 8.6 years. Most of the respondents, two hundred and forty-seven 247 (49.4%) had a Secondary level of Education. The majority of them were married 288 (57.6%).

Table 2 shows that they have a generally good knowledge of the prevention of HIV infection. However, there were significantly poor responses to delaying the onset of sexual intercourse and avoiding sex with CSWs as preventive measures. ($\chi^2 = 22.2$, $p < 0.01$; $\chi^2 = 12.5$, $p < 0.01$) respectively.

On their attitude to HCT in relation to educational status, Table 3 shows that those with tertiary education showed a significantly better attitude to HCT than the others. ($\chi^2 = 29.8$, $p < 0.01$). Table 4 showed that the prevalence of high-risk behaviour was high, with multiple sexual partners constituting 366 (73.2%), followed by sex with sex workers 258 (64.0%) and non-use of condoms (54.3%).

Table 5 showed that out of the 500 LDTD that were studied, only 89 (17.0%) had practised the HCT while 411 (83.0%) of them had not. Age groups <20 and ≥ 60 years constituted the highest unscreened group and those with a primary level of education constituted the highest with 171 (87.2%) and these were statistically significant.

The practice of HCT was highest in respondents with the tertiary educational level with all in level 11(100.0%) having practiced it. This difference was significant ($\chi^2 = 54.2$, $p < 0.01$, $df = 3$).

The married were more likely to practice HCT (24.7%) than the other marital status, however, this was not statistically significant ($\chi^2 = 28.2$, $p < 0.10$, $df = 7$). as shown in Table 5.

Fig. 1. showed that the willingness to screen was significantly higher in respondents with a tertiary level of education (100.0%) and least in those with no formal education 8.9%, ($\chi^2 = 10.6$, $p < 0.01$).

The respondents showed a generally good knowledge of the prevention of HIV/AIDS. However, there were poor significant differences in the response to delaying the onset of sexual intercourse and avoiding sex with CSWs as preventive measures. ($\chi^2 = 22.2$, $p < 0.01$; $\chi^2 = 12.5$, $p < 0.01$) respectively.

The willingness to screen was significantly higher in respondents with a tertiary level of education (100.0%) and least in those with no formal education 8.9%, ($\chi^2 = 10.6$, $p < 0.01$).

4. DISCUSSION

The main findings of the study were a generally fairly good knowledge of the prevention of HIV/AIDS among LDTD, ranging from (69% - 100%) across the educational levels. However, this does not translate to good prevention methods. It is interesting to note that sex with sex workers and not being faithful to one's partner had a significantly poor response (2.1% - 27%). This pattern is similar to other studies in the developing countries [21-23]. One of these studies [23] carried out in 17 countries in Africa, showed that more than half of young people did not know how to protect themselves from HIV. **Abstinence**, **Being faithful to one's partner** and the use of **Condoms**; this basic (**ABC**) of HIV prevention is not generally known and accepted [22,23].

The most prevalent high-risk behaviour found in this study was having multiple sexual partners 73.2%, followed by sex with sex workers 64%. This is similar to findings from other studies in developing countries [24,25]. Patronage of commercial sex workers; viewed as an occupational hazard amongst LDTDs may be due to their prolonged absence from their spouses, and hence this predisposition [25].

Table 1. Socio-demographic characteristics of Nigerian long-distance drivers

Socio-demographic characteristics	Frequency (n=500)	Percent (%)
Mean age = 38.3 ± 8.6 years		
Age group (years)		
<20	3	0.6
20-29	137	27.4
30-39	163	32.6
40-49	143	28.6
50-59	42	8.4
≥60	12	2.4
Educational status		
No formal Education	46	9.2
Primary	196	39.2
Secondary	247	49.4
Tertiary	11	2.2
Marital Status		
Single	147	29.4
Married	288	57.6
Divorced	30	6.0
Separated	35	7.0

Table 2. Knowledge of prevention of HIV/AIDS in relation to educational status

Correct prevention means	No formal Education n = 46(%)	Primary n = 196 (%)	Secondary n = 247 (%)	Tertiary n = 11 (%)	χ ²	P-Value
Staying with one faithful uninfected partner	45 (97.0)	192(97.0)	245 (99.0)	10 (98.0)	1.60	0.10
Using condoms every time	45 (97.0)	186(94.0)	231(93.0)	11 (100.0)	2.20	0.50
Abstaining from sex	32 (69.0)	141(71.0)	174 (70.0)	8 (72.0)	0.20	0.90
Delaying the onset of sexual intercourse	9 (18.0)	27(13.0)	32 (12.0)	7 (63.0)	22.2	0.01*
Avoiding sex with CSWs	2 (4.3)	12 (6.1)	9 (3.6)	3 (27.0)	12.5	0.01*
Reducing the number of sexual partners	1 (2.1)	11 (5.6)	20 (8.0)	0 (0.0)	3.5	0.30
Avoiding sex with people who have multiple sexual partners	8 (17.0)	24 (12.0)	42 (17)	1 (9.0)	2.5	0.50
Avoid sharing sharp objects like needles and razors	18 (39.0)	58 (29.0)	86 (34.0)	1 (9.0)	5.0	0.10
Incorrect prevention means						
Praying to God	18 (39.0)	61 (31.0)	80 (32.0)	3 (27.0)	1.2	0.70
Going for checkups	35 (76.0)	136(69.0)	192 (77.0)	8 (72.0)	4.1	0.30
Using antibiotics	32 (69.0)	138(70.0)	185 (74.0)	3 (27.0)	3.2	0.30
Seeking protection from a traditional healer	18 (39.0)	42(17.0)	24 (12.0)	1 (9.0)	2.3	0.50

*Statistically significant

Due to this, long-distance drivers are linked to spreading sexually transmitted infections including HIV in a short period of time [25].

The findings of Wilson et al [23], were quite disturbing. It showed some LDTDs are not only sexually active on the road, they could not

imagine going more than a few days without sex with whomever. They regularly taunt and belittle drivers who do not behave like them. This may lead to a culture of acceptance of these high-risk behaviours by the vulnerable ones who could not stand this 'ridicule' for being virtuous [23].

Table 3. Attitude of Nigerian long-distance drivers to HCT by educational status

Attitude	Educational status				χ^2	P-Value
	No formal Education n = 46(%)	Primary n=196 (%)	Secondary n=247 (%)	Tertiary n=11 (%)		
---Positive						
It is good for HIV prevention.	36 (78.3)	145 (74.0)	196 (79.4)	9 (81.8)	1.9	0.60
It should remain entirely voluntary and confidential.	38 (82.6)	153 (78.1)	202 (81.8)	10 (90.9)	1.9	0.60
It should be widely available	25 (54.3)	111 (56.6)	142 (57.5)	10 (90.9)	5.3	0.12
It should be compulsory for everybody.	1 (2.2)	4 (2.0)	4 (1.6)	3 (27.3)	29.8	0.01
Attitude ---Negative						
It is a breach on one's freedom	8 (17.4)	21 (12.8)	21 (9.7)	0 (0.0)	4.7	0.20
It does not lead to prevention of HIV/AIDS	8 (17.4)	25 (2.8)	24 (9.7)	0 (0.0)	4.1	0.30

Table 4. Prevalence of high-risk behavior among Nigerian long-distance drivers

High-risk behavior #	Freq n=398	% Percent
Multiple sex partner	366	73.2
Sex with sex workers	258	64.0
Do not use condoms	216	54.3
Share sharp objects/unsterile needle	21	5.3
Had unsafe blood transfusion	3	1.6

= Multiple response

Table 5. Demographic characteristics of respondents in relation to practice of HCT

Demographic variables	Practice of HCT		χ^2	p- value
	Yes n (%)	No n (%)		
	89 (17.0%)	411 (83.0%)		
Age group (years)				
<20	0 (0.0)	3 (100.0)	37.2 (5)	0.02
20-29	7 (5.1)	130 (94.9)		
30-39	44 (27.0)	119 (73.0)		
40-49	23 (16.1)	120 (83.9)		
50-59	15 (35.7)	27 (64.3)		
≥60	0 (0.0)	12 (100.0)		
Educational status				
No formal Education	8 (17.4)	38 (82.6)	54.2 (3)	0.01
Primary	25 (12.8)	171 (87.2)		
Secondary	45 (18.2)	203 (81.8)		
Tertiary	11 (100.0)	0 (0.0)		
Marital Status				
Single	6 (4.1)	141(95.9)	28.2 (7)	0.10
Married	71 (24.7)	217 (75.3)		
Divorced	6 (4.1)	25 (83.3)		
Separated	7 (20.0)	28 (80.3)		




Fig. 1. Willingness to screen for HIV infection among respondents

The practice of use of condoms during casual sex was found to be low among them with only 45.7% practicing it. However, this finding was somewhat better than studies done in India (11.0 %)[26] Burkina Faso (18.0%) [27] and Enugu (37.4%)[17] in Nigeria. The differences in practice may be explained by the timing of the studies, as this was done in the early years of the infection. Also, there may be some secular changes in the practice of condom use, as a result of current increasing awareness [23,25].

There was a generally good attitude to HCT in the range of (54.5% - 88.6%) across the group in this index study. However, this does not translate to good practice of HCT. Out of the 500 LDTDS that were studied, only 17.0% had practised HCT. This did not agree with previous studies [17,28] which showed both negative attitudes and practice of HCT. These changes may be due to increasing awareness as already observed. Again, in developed countries with higher literacy levels and resources, the uptake of HCT is very high [29].

The role of the church in the uptake of HCT was striking and commendable [18,21]. Its request for HIV screening tests before conducting marriage among its members took the lead as the reason for screening. This is to curb the spread of the disease among intending couples and their unborn children. However, this is just

the tip of the iceberg as the majority of the respondents were unscreened [21].

In our work, age groups 20-29 and 40-49 years constituted the highest unscreened group with (94.9%) and (83.9%) respectively. These findings were consistent with other findings [17,21,28], where most of the people infected with HIV were young people within the productive, socially active and reproductive age groups. This is the prime age and therefore has serious socio-economic and public health implications [21,28].

The reasons given by Nigerian long-distance drivers for this poor practice of HCT were lack of knowledge of where to go for HCT in about half of them (51%). Studies done in some African countries [19,20] showed that HCT services were not widely available, especially outside the main or urban centres. This was also shown by this index study, where all 89 respondents who had HCT obtained this service at urban centres. In most health institutions in Nigeria, HIV testing is limited to diagnostic purposes and not available to people in the general population interested in knowing their status [15,23]. This has a serious public health implications in the efforts to stem the HIV pandemic. An HIV-positive person with an unknown sero – status is a great danger to the public [15,23]. Our study also showed that fear and anxiety accounted for

37% of why LDTDs do not practice HCT. This is similar to findings from other studies [18-20]. It is a fact that people living with HIV/AIDS have been stigmatized and discriminated against. These negative effects discourage effective prevention by dissuading persons from coming freely for testing [30,31]. These negative attitudes towards people living with HIV/AIDS (PLWHA) are strongly linked to the general low level of knowledge about the disease and in particular, to the causes of HIV and its mode of transmission. In some societies, AIDS is associated with groups whose social and sexual behaviour does not meet public approval [30,31]. The other factors contributing to HIV/AIDS related stigma are; the fact that HIV/AIDS is a life-threatening disease which people are afraid to contract.

Although this index study noted a higher perception of risk for HIV infection, it also, fortunately, showed a high willingness to screen in about 68.0%. This finding is somewhat better in terms of willingness to screen in a study by Wilson, et al. [23]. In their 1995 and 1997 surveys, they found that 37.9 and 39.1 per cent of the drivers, respectively, considered themselves at risk of contracting HIV; only 30% of them were willing to screen for HIV. This poor willingness to screen in their study may be due to the earlier date of the study. This was when little was known about the disease and the driver's perception of individual risk of contracting the disease was poor.

Presently, although treatment modalities are improving, there is yet no cure for the infection or vaccine against its transmission. The only feasible option is to change people's sexual behaviour and attitude and encourage the uptake of HCT services, as this is the entry point of other preventive and curative packages in the fight against HIV/AIDS [4,11].

5. CONCLUSION

The way forward is to embark on behavioural change communication (BCC) via the following:

- Information dissemination about the transmission of HIV and the prevention methods of ABC needs to be targeted at truck stops, toll plazas, and border posts and at the workplaces of truck drivers.
- Misconceptions about the unproven and unscientific means of transmission and negative attitudes should be dissuaded

through adequate information by means of public awareness campaigns.

- There is a need for proper coordination between the individual, health care providers, the mass media, the non-governmental organizations and the government as policy formulators; in conjunction with various truck park representatives who should be given a short-term but well-packaged training in HCT adapted for use among their colleagues.

CONSENT

As per international standard or university standard, Participants' written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

Ethical clearance was given by the UNTH Ethical Committee with number NHREC/05/01/2019 – FWA00002458 – 1RB00002323.

ACKNOWLEDGEMENTS

The authors acknowledged the support of Motor Park authorities, the truck drivers and the families for being the base of the study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Olugbenga-Bello AI, Amusan OA, Oludele EA, Oluogun OA. Knowledge, attitude and uptake of voluntary confidential counseling and testing (VCCT) for HIV/AIDS, among young adults in an urban settlement in southwest Nigeria. *The Social Sciences Medwell Journals*. 2008;3:442-447.
2. Michael H. Merson. The HIV–AIDS pandemic at 25 — The global response. *N Engl J Med*. 2006;354:2414-2417.
3. Nzuobontane D, Blackett KN, Kuaban C. Cardiac involvement in HIV infected people in Yaounde, Cameroon. *Postgraduate Medical Journal*. 2002;78:678-681.
4. National scale up strategy for HIV counseling and testing in Nigeria. *National AIDS/STD Control Programme, Federal Ministry of Health*; 2017.

5. Araoye MO, Kayode O, Akande TM, Ndom RJE. Strategies for preventing STD/AIDS among commercial drivers. *Nigerian Medical Practitioner*. 1999;38:1-3, 5-10.
6. Mupemba K. The Zimbabwe HIV prevention program for truck drivers and commercial sex workers: a behaviour change intervention. 133-137 in *Resistances to Behavioural Change to Reduce HIV/AIDS Infection*, ed. J.C. Caldwell et al. Canberra: Australian National University; 1999.
7. United Nations programme on AIDS, HIV/AIDS – related stigmatization, discrimination and denial forms, contexts and determinants UNAIDS; 2015.
8. United Nations programme on AIDS: 4th National Conference on HIV/AIDS in Nigeria; 2014;40.
9. Okekearu I. Barriers of stigmatization and discrimination, human right Issues and ethical dilemma. Workshop on HIV/AIDS; 2002.
10. Global HIV/AIDS initiative Nigeria (GHAIN): HIV/AIDS-what to know and do. 2005;12.
11. Global HIV/AIDS Initiative Nigeria (GHAIN), counseling and testing. 2004;20.
12. Horizons. Research update: youth and HIV/AIDS Washington, DC. Population council/Horizons; 2000.
13. Inem AV, Adelokun AO, Avankogbe OO. A VCT programme for PMTCT of HIV on practice based research network (PBRN) Hospital in Lagos, Nig. *Nigeria Medical Practitioner*. 43:6.
14. Ofili AN; Knowledge, attitude and practice of long distance drivers towards HIV/AIDS. International Conference on AIDS. Bangkok, Thailand. *Int Conf AIDS*; 2004.
15. Oduwole M, Jeminusi OA, Aderogba OI, Okuboyejo OB. Long distance truck drivers (LDTD's) HIV/AIDS knowledge and attitude on the use of condoms. *Int. Conf AIDS*. 2002; 14(4):7-12;
16. Status of the epidemic. 4th National Conference on HIV/AIDS on Nigeria. 2004;18
17. Onwasigwe CN, Aniebue NP, Ndu AC. HIV/AIDS awareness and sexual behavior of Nigerian long Distance Drivers. *Journal of College of Medicine*. 2001;6:44-46.
18. Mukodzani L, Mupemba K, Marck J. All roads lead to Harare: the response of the Zimbabwe transport industry to HIV/AIDS. In *The Continuing African HIV/AIDS Epidemic*, ed. I.O. Orubuloye, J. C. Caldwell and J.P.M. Ntozi. Canberra: Australian National University. 1999; 131-138
19. Wilson D, Sibanda A, Nyathi B, Lamson N, Sibanda T. Ethnographic and quantitative research to design a community intervention among long-distance truck drivers in Zimbabwe. In *Focusing Interventions among vulnerable groups for HIV infection: experiences from Eastern and Southern Africa*. NARESA Monograph No. 2. Nairobi: Network of AIDS Researchers of Eastern and Southern Africa. 1994;99-108.
20. Rao KS, Pilli RD, Rao AS, Chalam PS. Sexual lifestyle of long distance lorry drivers in India: *B. M. J*. 1999;318:162-163.
21. Malta M, Bastos FI, Pereira-Koller EM, Cunha MD, Marques C, Strathdee SA. A qualitative assessment of long distance truck drivers' vulnerability to HIV/AIDS in Itajai, southern Brazil. *AIDS Care*. 2006;18:489-496.
22. Meda N, Sangare L, Iankoande S, Campaore IP, Catraye J, Sanou PT, Van – Dyck E, Cartoux M, and Sondre RB. The HIV epidemic in Burkina Faso: current Status and the knowledge level of the population about AIDS, 1994-1995. *Rev Epidemiol. Sante. Public*. 1998;46: 14-23.
23. ODU OO, Asekun-Olarinmoye EO, Bamisdele JO, Egbewale BE, Amusom OA, Olowu AO. Knowledge, Attitude to HIV/AIDS and sexual behavior of students in a tertiary institution in South –Western Nigeria. *European Journal of Contraceptive and Reproductive Health Care*. 2008;13(1):90-96.
24. Joint united nations Programme on HIV/AIDS (UNAIDS). HIV/AIDS-related discrimination, stigmatization and denial UNAIDS. 2001;7.
25. De Bruyn T. HIV/AIDS: discrimination and stigma. Definitions and concepts. Ottawa. Canadian HIV/AIDS Legal Network and the Canadian AIDS Society; 1999.
26. Ambati BK, Ambati J Rao AN. Dynamic of knowledge and attitude about AIDS among the educated in Southern India. *AIDS Care*. 1997;3:319–330.
27. Mbugua GG, Muthani LN, Mutura CW, Oogo SA, Waiyaki PG, Linda CP and Hearst N. Epidemiology of HIV infection among long distance truck drivers in Kenya. *East Afr Med. J* Aug. 1995; 72(8):515–8.

28. Obbo C. HIV transmission through social and geographic networks in Uganda. *Soc. Sci Med.* 1993;36 (7): 949–55.
29. Marck J. Long-distance truck drivers' sexual cultures and attempts to reduce HIV risk behaviour amongst them: a review of the African and Asian literature. In: Caldwell J, Caldwell P, Anarfi J, et al. eds. *Resistance to behavioural change to reduce HIV/AIDS Infection.* Canberra, Australia: Health Transition Centre, National Centre for Epidemiology and Population Health, Australian National University. 1999;91–100.
30. The world bank: *Confronting AIDS. Public priorities in a global epidemic.* New York. Oxford University Press; 1999.
31. Kippax S. Effective HIV prevention: the indispensable role of social science. *Journal of the International AIDS Society.* 2012 Apr;15(2):17357.

© 2023 Arodiwe and Arodiwe; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/106834>*