



Studies on Anti-Fungal Activity of Different Macro Algal Extracts against Soil Borne Pathogen *Sclerotium rolfsii* Sacc.

A. Praveen^a and C. Kannan^{a*}

^a Department of Plant Pathology, Faculty of Agriculture, Annamalai University, Annamalai Nagar – 608002, Tamil Nadu, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i193646

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/105023>

Original Research Article

Received: 17/06/2023
Accepted: 22/08/2023
Published: 31/08/2023

ABSTRACT

Management of soil-borne plant pathogens is one of the single greatest challenges facing modern agriculture worldwide. Seaweeds are considered as a source of bioactive compounds as they are able to produce a great variety of secondary metabolites characterized by a broad spectrum of biological activities. The seaweeds derived compounds showing broad range of biological activities such as antiviral, antibacterial and antifungal activities. The antimicrobial activity of macro algal extracts is generally analysed using various organic solvents which always present higher efficiency in extracting bioactive compounds. Among the different the solvent extracted macro algal extracts tested, methanol extracts of *Sargassum wightii* recorded maximum mycelial inhibition (88.22 %) and on par with aqueous extracts recorded 85.78% of mycelial inhibition @ 10% concentration compared to other macro algal extracts.. Among the fifteen macro algal extracts, maximum inhibition of mycelial growth was recorded the *Sargassum wightii* (Brown algae), *Ulva*

*Corresponding author: E-mail: inbakanna11@gmail.com;

lactuca (Green algae) and *Gracilaria salicornia* (red algae) at different concentrations. In this study, we have shown that *Sargassum wightii* has antimicrobial potential against the *S. rolfsii*. Methanol extract was found to be the best solvent for retention of the antimicrobial activity

Keywords: *Sclerotium rolfsii*; macro algae; *Sargassum wightii*; anti-fungal activity; methanol extracts; chloroform extracts; aqueous extracts.

1. INTRODUCTION

Sclerotium rolfsii Sacc. (Teleomorph: *Athelia rolfsii*) is an omnivorous, necrotrophic fungal plant pathogen which has a wide host range of over 500 plant species in 100 families, throughout the globe. This pathogen produce various enzymes viz., phytotoxin (oxalic acid), cell wall degrading enzymes and polysaccharides degrading multiplex enzymes which are affects the host plant system [1]. The typical symptom of this pathogen is rapid wilting and sickly appearance of plants with brownish lesion at the stem base near the soil lane which later girdles the stem [2]. The whitish mycelial growth forms over the infected tissue and often radiates over the soil surface [3]. *S. rolfsii* typically produces abundant white mycelium and small, brown, round sclerotia on the diseased tissue under hot humid conditions [4]. Sclerotia are the primary source for the infection which survives in soil for the several years in dormant stage. These sclerotia infect suitable host and left over in plant residues [5,6].

Seaweeds or “Marine algae” or “Marine macro algae” are the excellent source for plant disease management. These macro algae categorized into three groups based on the presence of pigments such as Green algae (Chlorophyta), Red algae (Rhodophyta) and Brown algae (Phaeophyta) which comprise nearly 9,000 algal species all over the world [7]. Seaweeds contain elaborate secondary metabolites that play a significant role in the host defence against various plant pathogens and parasites. These bioactive compounds (40000 and above) as secondary metabolites like Polysaccharides, polyphenols, carotenoids, proteins, peptides, sterols, terpenes and fatty acids are the main components of macro algal species which induce disease resistance in crop plants [8,9]. Different organic solvents of the algal extracts could provide a potential tool to explore the bioactive compounds responsible for positive effects on plant pathogens and mechanisms of their action [10,11]. The present investigation was carried out to analyse the efficiency of various macro algal

extracts in the management of *S. rolfsii* under in vitro condition.

2. MATERIALS AND METHODS

2.1 Isolation of Pathogen

The pathogen was isolated from the diseased plants showing typical symptoms of groundnut stem rot disease by tissue segment method [10]. The diseased portion of the stem was cut into small bits, surface sterilized with 0.1 per cent NaOCl₂ solution for 30 seconds washed in repeated changes of sterile distilled water. Potato Dextrose Agar medium was prepared and poured @15ml into sterilized Petri dishes. The Petri dishes were incubated in room temperature @ 28±2°C for five days and were observed for the fungal growth. The isolated fungal culture was purified by single hyphal tip method [12]. The purified culture were identified as *Sclerotium rolfsii* based on morphological and colony characteristics such as mycelial growth, mycelial dispersion, sclerotial germination, sclerotial shape, colour, number and arrangement of sclerotia on surface media.

2.2 Collection of Marine Macro Algae

Fifteen Marine macro algae were collected from east coastal areas of Tamil Nadu (Table 1). The collected materials were washed in fresh water to remove the debris, sand and extraneous matter. After draining off the water, the macro algae were wiped with a blotting sheet and air-dried under shade for two weeks, then cut into small pieces and dried in an oven at 45°C for 24 hours. The completely dried material was weighed and ground finely in a mechanical grinder. The powdered sample was stored in freezer for further study [13].

2.3 Preparation of Aqueous Extracts of Marine Macro Algae

100 grams of powdered macro algae was mixed with 100ml of distilled water and autoclaved at 150 lbs pressure for 1 h. The mixture extracts were filtered immediately through a muslin cloth.

Table 1. Macro algae collected from various seashore areas

Sl. No	Marine Macro algae (Seaweeds)	Common name or Group of Macro Algae	Place of collection
1	<i>Dictyota dichotoma</i>	Brown algae	Kanyakumari
2	<i>Sargassum wightii</i>	Brown algae	Pamban
3	<i>Padina gymnospora</i>	Brown algae	Pamban
4	<i>Hydroclathrus hornemanii</i>	Brown algae	Pamban
5	<i>Turbinaria coniodes</i>	Brown algae	Rameshwaram
6	<i>Caulerpa scalpelliformis</i>	Green algae	Pamban
7	<i>Chaetomorpha antennina</i>	Green algae	Mandapam
8	<i>Enteromorpha intestinalis</i>	Green algae	Velankanni
9	<i>Ulva lactuca</i>	Green algae	Rameshwaram
10	<i>Halimeda gracilis</i>	Green algae	Puducherry
11	<i>Acanthophora spicifera</i>	Red algae	Mandapam
12	<i>Gracilaria salicornia</i>	Red algae	Pamban
13	<i>Jania rubens</i>	Red algae	Puducherry
14	<i>Kappaphycus alvarezii</i>	Red algae	Mandapam
15	<i>Hypnea musciformis</i>	Red algae	Mandapam

All the extracts were labelled and stored in separate bottles which were kept in a refrigerator. The extracts thus obtained were crude macro algal extracts. Crude extract of different concentration was prepared with distilled water for further studies.

2.4 Preparation of Solvent Extracts of Marine Macro Algae

Solvent extracts of marine macro algae was prepared by using soxhlet extractor. 20 grams of powdered macro algae was poured in the thimble and extracted successively with each 100 ml of two different solvents viz., Methanol and Chloroform. The extracts were filtered using a Millipore filter unit 0.45 µm pore size, kept in -20° C in airtight brown bottles for further studies [14].

2.5 Poison Food Technique (Groover and Moore, 1962)

Potato dextrose agar media amended with macro algal extracts at different concentrations viz., 2.5%, 5% and 10% were autoclaved and poured into sterile Petri dishes. Then the plates were inoculated with five days old culture disc of *Sclerotium rolfsii* isolate (SrALR). Three replications were maintained in each treatment and incubated at laboratory temperature 28 ± 2°C. Potato dextrose agar media without macro algal extract served as control. The diameter of the mycelial growth was measured in 5 days after incubation (DAI). The per cent inhibition of the test fungi was calculated by the formula of Vincent (17).

$$I = \frac{C - T}{C} \times 100$$

Where,

- I = Per cent inhibition of fungal growth
- C = Growth in Control
- T = Growth in treatment

3. RESULTS

3.1 Antifungal Activity of Macro Algal Extracts Against *Sclerotium rolfsii* (SrALR)

Fifteen different brown, green and red macro algal extracts were selected and evaluate the antifungal activity against *S. rolfsii* by Poison food technique. Among the different macro algal extracts were tested, *Sargassum wightii*, *Ulva lactuca* and *Gracilaria salicornia* showed effective antifungal activity against the *S. rolfsii*. The brown algae *Sargassum wightii* significantly reduced the mycelial growth (1.12 cm) of *S. rolfsii* and recorded highest mycelial inhibition 87.56% at 10% concentration (Table 2 & Plate 1). The green algae *Ulva lactuca* showed maximum reduction in mycelial growth 1.39 cm with highest mycelial inhibition (84.56%) of test pathogen at 10% concentration (Table 3 & Plate 2). Likewise, the red algae *Gracilaria salicornia* effectively inhibit the radial growth (2.08 cm) of *S. rolfsii* with 76.89% of mycelial inhibition at 10% concentration (Table 4 & Plate 3).

Table 2. *In vitro* Evaluation of Brown macro algal extract against *Sclerotium rolfsii* (SrALR)

S. No	Treatment	Mycelial growth (cm) (5DAI)					
		2.5%	Percent inhibition over control	5%	Percent inhibition over control	10%	Percent inhibition over control
1	<i>Dictyota dichotoma</i>	4.53 ^c (12.29)	49.67	3.92 ^d (11.42)	56.45	3.46 ^d (10.71)	61.56
2	<i>Sargassum wightii</i>	2.86 ^a (9.74)	68.23	1.39 ^a (6.77)	84.56	1.12 ^a (6.08)	87.56
3	<i>Padina gymnospora</i>	3.19 ^b (10.28)	64.56	2.36 ^b (8.84)	73.78	1.43 ^b (6.87)	84.12
4	<i>Hydroclathrus hornemanii</i>	5.69 ^d (13.80)	36.78	5.02 ^e (12.94)	44.23	4.79 ^e (12.64)	46.78
5	<i>Turbinaria coniodes</i>	3.40 ^b (10.62)	62.22	2.96 ^c (9.91)	67.12	2.10 ^c (8.33)	76.67
6	Control	9.00					

Mean of three replications

Values in the column followed by same superscript letters do not differ significantly at 5% level by Duncan's multiple range test (DMRT)

Table 3. *In vitro* Evaluation of Green macro algal extract against *Sclerotium rolfsii* (SrALR)

S. No	Treatment	Mycelial growth (cm) (5DAI)					
		2.5%	Percent inhibition over control	5%	Percent inhibition over control	10%	Percent inhibition over control
1	<i>Caulerpa scalpelliformis</i>	5.89 ^e (14.05)	34.56	5.63 ^d (13.72)	37.45	5.00 ^e (12.92)	44.44
2	<i>Chaetomorpha antennina</i>	5.26 ^d (13.26)	41.56	4.91 ^c (12.80)	45.67	4.53 ^d (12.29)	49.67
3	<i>Enteromorpha intestinalis</i>	4.10 ^b (11.67)	54.45	3.62 ^b (10.97)	59.78	2.86 ^b (9.73)	68.23
4	<i>Ulva lactuca</i>	3.39 ^a (10.61)	62.33	2.48 ^a (9.05)	72.45	1.39 ^a (6.77)	84.56
5	<i>Halimeda gracilis</i>	4.47 ^c (12.21)	50.32	3.86 ^b (11.33)	57.12	3.13 ^c (10.19)	65.23
6	Control	9.00					

Mean of three replications

Values in the column followed by same superscript letters do not differ significantly at 5% level by Duncan's multiple range test (DMRT)

Table 4. *In vitro* Evaluation of Red macro algal extract against *Sclerotium rolfsii* (SrALR)

S. No	Treatment	Mycelial growth (cm) (5DAI)					
		2.5%	Percent inhibition over control	5%	Percent inhibition over control	10%	Percent inhibition over control
1	<i>Acanthophora spicifera</i>	4.87 ^b (12.74)	45.89	4.42 ^d (12.14)	50.89	3.91 ^d (11.41)	56.56
2	<i>Gracilaria salicornia</i>	4.02 ^a (11.57)	55.34	3.37 ^a (10.58)	62.56	2.08 ^a (8.30)	76.89
3	<i>Jania rubens</i>	5.27 ^c (13.26)	41.45	4.87 ^e (12.74)	45.87	4.33 ^e (12.01)	51.89
4	<i>Kappaphycus alvarezii</i>	4.56 ^b (12.33)	49.33	3.98 ^c (11.51)	55.78	3.24 ^c (10.36)	64.00
5	<i>Hypnea musciformis</i>	4.19 ^a (11.81)	53.44	3.65 ^b (11.02)	59.45	2.78 ^b (9.60)	69.12
6	Control	9.00					

Mean of three replications

Values in the column followed by same superscript letters do not differ significantly at 5% level by Duncan's multiple range test (DMRT)

Table 5. *In vitro* evaluation of different solvent extracts of macro algae against *Sclerotium rolfsii* (SrALR)

S. No	Treatment	Percent inhibition over control								
		Aqueous extract			Methanol extract			Chloroform extract		
		2.5%	5%	10%	2.5%	5%	10%	2.5%	5%	10%
1	<i>Sargassum wightii</i>	57.89	82.78	85.78	70.22	85.11	88.22	50.11	58.67	74.22
2	<i>Padina gymnospora</i>	48.22	61.11	66.44	50.78	64.44	70.33	43.11	50.00	58.56
3	<i>Enteromorpha intestinalis</i>	40.11	49.67	54.22	42.33	52.33	57.67	33.78	40.89	44.56
4	<i>Ulva lactuca</i>	46.44	70.78	74.67	66.44	73.89	78.56	48.67	56.00	65.67
5	<i>Gracilaria salicornia</i>	43.78	58.78	61.44	46.89	60.44	65.56	39.67	44.22	49.78
6	<i>Hypnea musciformis</i>	35.22	44.22	46.00	37.44	44.67	48.67	29.56	34.78	40.67
7	Control	9.00								

Mean of three replications

Values in the column followed by same superscript letters do not differ significantly at 5% level by Duncan's multiple range test (DMRT)




Plate 1. *In vitro* Evaluation of *Sargassum wightii* (Brown macro algal extract) against *Sclerotium rolfsii* (SrALR)




Plate 2. *In vitro* Evaluation of *Ulva lactuca* (Green macro algal extract) against *Sclerotium rolfsii* (SrALR)




Plate 3. *In vitro* Evaluation of *Gracilaria salicornia* (Red macro algal extract) against *Sclerotium rolfsii* (SrALR)

3.2 *In vitro* Evaluation of Different Solvent Extracts of Macro Algae Against *Sclerotium rolfsii* (SrALR)

Different macro algal extracts were prepared by using two different solvents viz., Methanol and Chloroform, to investigate for their efficacy compared with aqueous macro algal extracts under poison food technique (Table 5). The results revealed that methanol macro algal extracts were significantly inhibited the test pathogen of *S. rolfsii*. Among the six macro algal extracts, methanol extracts of *Sargassum wightii* recorded maximum mycelial inhibition (88.22 %) and on par with aqueous extracts recorded 85.78% of mycelial inhibition @ 10% concentration compared to other macro algal extracts.

4. DISCUSSION

In this present study, our findings are accordance with earlier reports by Karthik et al. [15], they studied efficacy of seaweed liquid fertilizer (SLF) *Turbinaria ornata* and *Ulva reticulata* against the soil borne pathogen *S. rolfsii* and reported that zone of mycelial inhibition was noticed after 72 hours of incubation. Sarkar et al. [16] and Latifeh et al. [17] stated that macro algal fertilizer or seaweed liquid fertilizer from algal group viz., *Sargassum polyphyllum*, *Gelidiopsis* sp., *Padina tetrastomatica*, and *Gracilaria corticata*, which exhibited antagonistic effect against various plant pathogens such as *Rhizoctonia solani*, *Macrophomina phaseolina*, *Sclerotium rolfsii* and *Alternaria solani*. Sujatha et al. [18] stated that the antifungal activity of macro algal extract

against soil borne pathogens in pulses. The results revealed that methanol extract of *S. myricocystum* have a strongest antifungal ability against different soil borne pathogens like *Rhizoctonia solani* and *Macrophomina phaseolina*. Rupapara et al. [19] reported that methanol extracts of *Sargassum johnstonii* showed a significant antimicrobial activity against plant pathogens due to the presence of various bioactive antimicrobial compounds viz., alkaloid, protein carbohydrates, and phenolics. Macro algae contain the different bioactive molecules in secondary metabolites such as terpenes, phenols, alkaloids, fatty acids and polysaccharides that have ability to significantly reduce the mycelial growth of *S. rolfsii* under in vitro conditions [20].

5. CONCLUSION

These studies show that the methanol solvent extraction of macro algae from *Sargassum wightii* has most effective inhibitory action against examined pathogen of *S. rolfsii*. Future researches in *S. wightii* are needed to determine their bioactive compounds and specifically investigate their anti-microbial activity against plant pathogens.

ACKNOWLEDGEMENT

I extend my gratitude to the department of plant pathology, faculty of agriculture, annamalai university for their support.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bowen Z, Dou H, Peiqing L, Rongbo W, Benjin L, Qinghe C. Occurrence of collar rot caused by *Athelia rolfsii* on soybean in China. *Canadian Journal of Plant Pathology*. 2021; 43(1):43-47.
2. Groover RF, Moore RD. Resistance inducing chemicals and disease resistance in plants. *Phytopathology*. 1962;52:876-880.
3. Karthik Pandi V, Gopalakrishnan C, Janahiraman V. Cultural and Morphological variability in *Sclerotium rolfsii* causing stem rot disease. *International Journal of Current Microbiology and Applied Science*. 2017; 6(6):3090-3097.
4. Kumari A, Ghatak A. Variability in Chickpea rot-causing soil-borne necrotrophs, *Sclerotium rolfsii* and *Macrophomina phaseolina*. *Journal of AgriSearch*. 2018;5(4):247-253.
5. Yalda Vasebi, Naser Safaie, Azizollah Alizadeh. Biological control of soybean charcoal root rot disease using bacterial and fungal antagonists in vitro and greenhouse condition. *Journal of Crop Protection*. 2013;2(2):139150.
6. Windels, Carol E, Burnes MP, Kommedahl T. Five-year preservation of fusarium species on silica gel and soil. *Phytopathology*. 1988;78(1):107-109.
7. Chaturvedi S, Kulshrestha S, Bhardwaj K. Role of seaweeds in plant growth promotion and disease management. In *New and Future Developments in Microbial Biotechnology and Bioengineering*. 2022; 217-238
8. Vicente TF, Lemos MF, Felix R, Valentao P, Felix C. Marine macroalgae, a source of natural inhibitors of fungal phytopathogens. *Journal of Fungi*. 2021;7(12):1006.
9. Selma Mabrouki, Fatima Lakhdar. In Vitro Antifungal activity of seaweed extracts against *Sclerotium rolfsii*: A Causal agent of root rot disease on sugar beet (*Beta vulgaris* L.). *International Journal of Pharmaceutical Sciences Review and Research*. 2020;63(2):90-97.
10. Michalak I, Chojnacka K. Production of seaweed extracts by biological and chemical methods. In: Kim S-K, Chojnacka K. (Eds.), *Marine Algae Extracts: Processes, Products, Applications*. Wiley-vch Verlag GmbH & Co. KGaA, Weinheim in press; 2015.
11. Rangaswami G. An agar block technique for isolating soil microorganisms with special reference to pythiaceae fungi. *Scientific Culture*. 1958;24-85.
12. Vincent JM. Distortion of fungal hyphae in the presence of certain inhibitors. *Nature*. 1927;59:850.
13. Aseer M, Sugathan S, George SK, Joseph S, Chippu S, Ramakrishnan G, Mamkoottathil VNP. Bio-potentials of seaweeds collected from southwest coast of India. *Journal of Marine Science*. 2009; 17(1):67-73.
14. Khallil AM, Dagman IM, Fady AA. Antifungal potential in crude extracts of five selected brown seaweeds collected from

- the western Libya coast. Journal of Microbiology and Modern Techniques. 2015;1(1):103
15. Karthik T, Sarkar G, Babu S, Amalraj LD, Jayasri MA. Preparation and evaluation of liquid fertilizer from *Turbinaria ornata* and *Ulva reticulata*. Biocatalysis and Agricultural Biotechnology. 2020;28:01712.
 16. Sarkar G, Jatar N, Goswami P, Cyriac R, Suthindhiran K, Jayasri MA. Combination of different marine algal extracts as bio-stimulant and bio-fungicide. Journal of Plant Nutrition. 2018;41:1163-1171.
 17. Latifeh Pourakbar, Sina Siavash Moghaddam. Antifungal activity of the extract of a macroalgae, *Gracilariopsis persica*, against four plant pathogenic fungi. Plants (Basel). 2021;10(9):1781.
 18. Sujatha K, Mahalakshmi P, Manonmani K. Effect of antifungal activity of seaweed extracts against soil borne pathogens in Pulses. International Journal of Agriculture Innovations and Research. 2014;3(1):135-138.
 19. Rupapara KV, Joshi NH, Vyas KG. Evaluation of antimicrobial activity of crude extracts of seaweed *Sargassum johnstonii*. International Journal of Current Microbiology and Applied Sciences. 2015; 4:300-304.
 20. Junaithal A, Begum M, Selvaraju P, Vijayakumar A. Evaluation of antifungal activity of seaweed extract (*Turbinaria conoides*) against *Fusarium oxysporum*. Journal of Applied and Natural Science. 2016;8(1):60-62.

© 2023 Praveen and Kannan; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/105023>