



Land Use and Land Cover Change Study Using RS-GIS and Its Impact on Biodiversity: Study from Arunachal Pradesh, India

K. Ronald ^a, K. J. Singh ^b and M. S. Lodhi ^{c*}

^a College of Agriculture Sciences, FEEDS Group of Institutions, Hengbung, Kangpokpi, Manipur-795129, India.

^b Department of Civil Engineering, North Eastern Regional Institute of Science and Technology (NERIST), Arunachal Pradesh, India.

^c G. B. Pant National Institute of Himalayan Environment (GBP-NIHE), Almora, Uttarakhand, India.

Authors' contributions

This work was carried out in collaboration among all authors. Author KR designed the study, performed field survey data collection, analysis and manuscript writing. Authors KJS and MSL managed the analyses of the study and finalized the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJEE/2023/v22i4514

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/110457>

Original Research Article

Received: 10/10/2023

Accepted: 13/12/2023

Published: 22/12/2023

ABSTRACT

This study aims to depict the intricate relationship between biophysical parameters, particularly land surface temperature (LST), and their impact on biodiversity at Itanagar Wildlife Sanctuary in Arunachal Pradesh. Focused on the consequences of anthropogenic activities, our research assesses changes in forest cover using Landsat images from 1990, 2003, and 2019. Utilizing satellite data, calculation of indices such as Normalized Difference Vegetation Index (NDVI), Normalized Difference Built-up Index (NDBI), Normalized Difference Water Index (NDWI), and LST were performed. Land use land cover (LULC) was determined through supervised classification,

*Corresponding author: E-mail: mahen29.mail@gmail.com;

achieving accuracies of 75%, 87%, and 83% for 1990, 2003, and 2019, respectively. The analysis highlighted a substantial increase in settlement area from 9.37 km² in 1990 to 22.86 km² in 2019, indicative of significant urbanization impacts. Our findings reveal a pronounced positive correlation between settlement growth and elevated LST, indicating the influence of urbanization on thermal dynamics. Additionally, positive correlations were observed between LST and NDBI, while a negative correlation existed between LST and NDVI. Questionnaire-based surveys complemented remote sensing data, exposing a gradual decline in wildlife sightings, diminishing populations of rare plant species, and the proliferation of invasive plants within and around the Wildlife Sanctuary since 1990. This research underscores the substantial impact of urbanization on the biodiversity of the Itanagar Wildlife Sanctuary. Positive correlations between settlement growth, LST, and biophysical parameters emphasize the urgency for targeted conservation strategies. The decline in wildlife sightings and the encroachment of invasive species underscore the need for proactive measures to balance biodiversity conservation with the challenges posed by urban development. Our integrated approach, combining remote sensing, survey data, and ecological indices, offers valuable insights for sustainable management and conservation efforts in similar habitats globally.

Keywords: LULC; LST; NDIV; NDBI; NDWI; biodiversity conservation; RS-GIS.

1. INTRODUCTION

The rapid urbanization has affected the biodiversity, water balance and socio-economic conditions which resulted as change in climate and biosphere [1,2]. Studies based on conservation and management of biodiversity using remote sensing and GIS technology has become more popular because of its effective and fast result compared to conventional methods. In the last two decades, researchers have focused their attentions towards local and regional climate study which are under influences of anthropogenic activities for understanding the various driving factors related to changing climate [3,4]. The overall affects due to urbanization also contributes to global phenomena such as GHG emission; change in temperature and precipitation and anthropogenic climate change [5,6,7]. The study on changes of land use land cover and its impact on flora and fauna in the biodiversity hotspot of India have become more popular among the researchers in the country and abroad. The rapid changes on the land use land cover pattern in the Western Ghat, one of the important biodiversity hotspots of India, has been studied and reported, the huge transformation due to anthropogenic and natural calamities [8,9]. The monitoring of land use land cover (LULC) changes over a period of time with highly accurate information is very necessary for biodiversity conservation and management especially in the youngest mountain ranges i.e. north-eastern Indian Himalaya. This information will help in understanding the interaction between human activities and environments and its ultimate affects. Furthermore, it will enable to manage the available recourses by utilizing sustainably and

maintaining balance environment [10,11]. Land surface temperature (LST) can be stated as one of the significant parameters for understanding the status of urban environment [12]. Many researches on climate change, urban climate, hydrological cycle, vegetation and environmental research have successfully employed the LST as an important parameter of the study along with indices like NDVI NDBI and NDWI [13,14,15,16]. The mapping of protected areas in Arunachal Pradesh using RS and GIS techniques was found useful for conservation of flora and fauna [17]. The present study has been undertaken to quantify and understand the spatio-temporal changes in the region due to urbanization with special reference to effects on biodiversity and conservation.

2. MATERIALS AND METHODS

2.1 Study Area

The Eastern Himalayan region is known for its rich biodiversity and a sophisticated geomorphic history with pervasive topographic features [18]. The present study site is located in the state of Arunachal Pradesh in the north eastern part of the India, which is also the part of eastern Himalayan biodiversity hotspot [19,20]. The Itanagar Wildlife Sanctuary is situated in the Papumpare districts of Arunachal Pradesh with an area of 140.8km². The sanctuary was created on 14/06/1978 vide Govt. Notification No. FOR 118/78 by superseding the Itanagar Reserved Forest as known earlier (Fig: 1). The Wildlife Sanctuary fall under the survey of India (SOI) topo-sheet No. 83E/12 Subansiri. The sanctuary is located

adjacent to the state capital Itanagar and is surrounded by three rivers i.e., Poma River in the East, Pachin River in the South and Neorch River in the North West. The vegetation of the study area can be classified as north bank tropical evergreen, tropical semi-evergreen forest and Secondary Forest [21]. The forest can be grouped in sub group 2B Northern Tropical Semi Evergreen Forest and Sub group 3C North Indian moist deciduous forest [22]. The sanctuary is home to numbers of diverse mammals, reptiles and birds some notable mammalian like elephants (*Elephas maximus*), tigers (*Panthera tigris*), leopards (*Panthera pardus*), clouded leopards (*Neofelisnebulosa*), sambar (*Cervus unicolor*), barking deer (*Muntiacusmuntjak*), dholes (*Cuonalpinus*), wild boars (*Sus scrofa*), jackals (*Canis aureus*), and small cats [23].

2.2 Satellite Data

The LANDSAT image data are the most widely use satellite data in the world for observation of environmental changes because of its large repository of continuous spatial data with longer time periods [24]. The boundary of the selected wildlife sanctuary was obtained from Department of Environment and Forest, Govt. of Arunachal Pradesh in a PDF file format. After obtaining the boundary of the sanctuary a 1 km. buffer was created in Arc-GIS 10 for the study (Fig 1). LANDSAT satellite images were also acquired (Path/row: 135/41) from USGS earth explorer website for the study. Based on the availability of imageries with less than 10% cloud cover, three set at different time scales (1990, 2003, 2019) were downloaded in GEOTIFF format which are in UTM Zone 46N of WGS 84 datum (Table 1).




Fig. 1. Study site

Table 1. Characteristics of the satellite data used in the study

Date	Path/Row	Satellite	Sensor	No. of Band	Spatial Resolution
1990/05/17	135/41	Landsat 5	TM	7	Thermal= 30x30 Reflective= 30x30
2003/03/10	135/41	Landsat 7	ETM	8	Thermal= 30x30 Reflective= 30x30 Panchromatic= 15x15
2019/04/06	135/41	Landsat 8	OLI_TIRS	11	Thermal= 30x30 Reflective= 30x30 Panchromatic= 15x15

2.3 Estimation of Indices and LST

The pre-processing of the satellite images i.e., geometric, atmospheric correction and data analysis were performed in the ERDAS Imagine 14 and Arc-GIS10 software. The derivations of normalized indices NDVI, NDBI, NDWI were performed by computing the band ratios. These indices were used to understand the relation with various biophysical parameters like greenness, built-up intensity and wetness with respect to thermal responses of the concern environment. The red band (RED)(0.63-0.69mm) and near-infrared (NIR) band (0.76-0.9mm) were processed for normalized difference vegetation index (NDVI) analysis [25]. Normalized difference built-up index (NDBI) is used in detecting the built-up areas as it has higher reflectance of short-wave infrared wavelength (1.55-1.75mm), so it is calculated with NIR [26]. NDWI measures the liquid water molecules in vegetation that interact with solar radiation. NDWI is computed using the green band and the near-infrared band [27]. The thermal infrared band (10.4-12.5mm) was used in retrieving the land surface temperature (LST) of the study area [28,29].

2.4 Classification of Land Use and Land Cover (LULC)

The land use and land cover (LULC) classification is the expression of the earth surface utilized by human and its observation of the earth surface [30]. The analysis of LULC is becoming an important tool for interpreting the rapid changes of the environment over the course of time [31,32]. The supervised

classification method was applied using the maximum likelihood (ML) algorithm, which is one of the extensively used algorithms in remote sensing for its robustness [33]. Six LULC classes were identified for Itanagar wildlife sanctuary. The LULC includes (i) Forest (ii) Jhum land (iii) Barren land (vi) Agriculture land (v) Built-up and (vi) water body/ river. The overall methodology followed for the analysis are represented as flowchart (Fig. 2).

2.5 Collection of Ground Truth Data and Field Survey

The study on flora diversity and faunal interaction with the local community was conducted in the 8 villages (Table 2 and Fig 1). A total of 82 respondents were selected within the age group of 30-55 for formal and informal interviews. The data on sighting of wildlife in and around the village and conflict related to wildlife with the villagers in the past three decades were recorded. The floral diversity of the study area was also recorded by conducting extensive field survey in forest area near the survey villages. The identification of the faunal diversity in the field was done with field guide book [34]. Plant species were identified by following taxonomic literatures and consultation with the experts of regional herbaria (ASSAM and ARUN) of Botanical Survey of India (BSI), Govt. of India. A total of 100 random GPS point samples were collected in the study area describing each LULC classes (Fig 2). Gramintrex30 GPS device was used to collect the location data.




Fig. 2. Flowchart of the methods.

Table 2. List of surveyed villages in Papum Pare district (Arunachal Pradesh)

Sl. No	Village	Dominant Tribal community	Number of Household	Number of respondents
1	Poma	Nyshi	26	15
2	Bat	Nyshi	37	6
3	Ganga	Nyshi	86	15
4	Jollang	Nyshi	71	8
5	Papu nallah	Nyshi	77	10
6	Upia	Nyshi	21	10
7	Sote	Nyshi	40	10
8	Pachin	Nyshi	27	8

2.6 Accuracy Assessment and Quantitative Analysis of LST with Biophysical Parameters

The accuracy assessment of the LULC was performed by using 100 randomly collected GPS points which covers all the classes. Additionally, the calculation of producer's accuracy, user accuracy, overall accuracy and Kappa coefficient was also done using standard methods [35]. However, to understand the dynamics relationship of LST and the biophysical parameters (NDVI, NDBI, NDWI) a total of 100 point were randomly generated in GIS platform which are well distributed in all the classes of LULC. The random points were then used to extract the values of biophysical parameters and LST of all the studied images. The correlation and regression analysis are used to understand how the environments responds to urbanization and its impact of urbanization on biodiversity.

3. RESULTS AND DISCUSSION

3.1 Land Used Land Cover Change Analysis

The temporal LULC map of the Itanagar Wildlife sanctuary was prepared for three different time periods 1990, 2003, 2019 to understand the changes in LULC due to urbanization (Fig 3). A total of six classes of LULC were classified which includes i) Forest- it includes all the forest covers including bamboo mixed forest; ii) Jhum land (shifting agriculture) - it includes all types of mixed cultivation with few sparsely distributed trees, Fallow land with shrubby dominance; iii) Agriculture- this includes mostly the areas under monoculture of paddy specially the wet rice cultivation; iv) Barren- it includes the area with no vegetation or exposed soil either naturally through landslide or by anthropogenic influence; v) Built up area- this represents the settlements

in the area; vi) Water- it includes river and lake. The distributions of LULC classes along the time line (1990, 2003 and 2019) are clearly showing continuous changes with rapid growth in built-up areas (Table 3). The Forest cover of the study area was found to be decreasing gradually as it represents a total of 181. 48 sq. Km (77.88%) in 1990 while it was reduced to 163.98 sq. Km (70.37%) by 2019. However, an opposite trend of gradual increase was found in Jhum with 15.52 sq. Km in 1990, 20.63 sq. Km in 2003 and 22.86 sq. Km in 2019. Moreover, built-up classes were also observed with similar trend, with 9.37 sq. Km, 10.59 sq. Km, and 22.86 sq. Km in 1990, 2003, 2019 respectively. The rural-urban migration in the study area has influence the expansion of built-up areas. According to population census 1991 and 2011, the population has increased from 16, 550 to 59,490 [36]. The increase in population in the area can be attributed to movement of from rural to urban areas for quality life and opportunities. A similar type of example of LULC changes has been observed in the twin cities of Minnesota (USA), where the forest and agriculture land decreased gradually and Built-up areas expanded with the time [37]. The accuracy assessment on LULC classes reveals that the overall accuracy was 75, 87, and 83% for the maps of 1990, 2003 and 2019 respectively while, kappa co-efficient for 1990, 2003 and 2019 are 0.70, 0.84 and 0.79 respectively (Table 4). Moreover, the result on LULC of the study area was found reasonable as it ranges under good or satisfying agreements which are sufficient for analyzing the changes along the 3-decade time span [38].

3.2 Impact of Urbanization on Biodiversity

The rapid urbanization in the world has resulted in the growth of myriad form of structures which modified the energy and water balance by

changing the dynamics of air and surface temperature [39,40]. The analysis on the Land surface temperature (LST) has reveals the thermal changes in the study area during 1990, 2003 and 2019 (Fig 4). It was estimated that the highest surface temperature in Itanagar and Naharlagun area were 28.83°C, 29.21°C and 30.21°C for 1990, 2003 and 2019 respectively. The urban heat island (UHI) can be attributed by the energy transformation in the cities; decrease

of evapotranspiration and production of anthropogenic energy [41]. It was observed that the temperature class of 24-30 °C was spreading in almost all the area of the Itanagar city in year 2019 depicting an urban heat island (UHI). However, there were very less area of UHI with the same temperature class (24-30°C) in 1990 (Fig 4). The increasing built-up area in Itanagar city and deforestation around the sanctuary has resulted in rise of temperature.

Table 3. Land use land cover (LULC) distribution of Itanagar Wildlife Sanctuary

LULC Classes	1990		2003		2019	
	Area (sq.Km)	Area (%)	Area (sq.Km)	Area (%)	Area (sq.Km)	Area (%)
Forest	181.48	77.88	168.77	72.43	163.98	70.37
Jhum (Shifting agriculture)	15.52	6.66	20.63	8.85	22.64	9.72
Agriculture	13.21	5.67	8.97	3.85	9.62	4.13
Barren	10.45	4.49	20.74	8.90	9.65	4.14
Built-up	9.37	4.02	10.59	4.54	22.86	9.81
Water	3.00	1.29	3.39	1.46	4.27	1.83




Fig. 3. LULC of Itanagar Wildlife Sanctuary in 1990, 2003 and 2019.

Table 4. Accuracy assessment LULC classification for 1990, 2003 and 2019

Class	1990		2003		2019	
	PA	UA	PA	UA	PA	UA
Forest	0.94	0.80	0.93	0.79	0.87	0.81
Jhum	0.73	0.73	0.80	0.85	0.87	0.81
Agriculture	0.53	0.47	0.87	1	0.87	0.93
Barren	0.78	0.73	0.90	0.78	0.70	0.82
Built-up	0.68	0.73	0.90	0.90	0.95	0.76
Water	0.85	0.90	0.73	1	0.67	0.90
Overall Accuracy %	75		87		83	
Kappa coefficient	0.70		0.84		0.79	

The three biophysical parameters NDVI, NDBI and NDWI for the year 1990, 2003 and 2019 are present in Fig. 5. The NDVI map for different time periods 1990, 2003 and 2019 clearly indicates the effect of urbanization on vegetation cover in the study area. The mean index values of NDVI are 0.41, 0.41 and 0.22 for 1990, 2003 and 2019 respectively. However, the mean NDBI values are found increasing i.e. -0.01, -0.11 and -0.13 for 1990, 2003, 2019 respectively. Correlation coefficient analysis of LST with NDVI, NDBI and NDWI were found comparatively acceptable (Table 5). The NDVI-LST and NDVI-NDBI were found negatively correlated for the years 1990, 2003, 2019 with the values of -0.42, -0.48, -0.45 and -0.79, -0.68, -0.86 respectively (Table 5). The result has also supported the statement that increasing built-up has influence on vegetation and temperature in the urban areas.

3.3 Changes in the Phyto-Diversity Status

The study on floral diversity of the sanctuary has reveals that the forest represents heterogeneous mixtures of species from semi-evergreen to evergreen forest. Among the tree species *Dipterocarpus macrocarpus*, *Artocarpus chama*, *Magnolia spp.*, *Aglaia spectabilis*, *Castonopsis indica*, *Schimawallichii*, *Duabanga grandiflora*, *Parkiatimoriana*, *Toona ciliate*, *Sapindus murkorossii* represent the higher canopy layers. While, the understory tree layers were *Lagerstroemia speciosa*, *Macaranga denticulata*, *Bauhinia purpurea*, *Mallotussp.*, *Dysoxylumgobara*, *Actephilaexcelsa*, *Callicarpa arborea*, *Musa sp*, *Cyathea sp*. The shrub layers in the area includes *Buddleja asiatica*, *Sterculia hamiltonii*, *Pinangagracilis*, *Phryniumpubinerve*, *Croton caudatus*, *Angiopterisevecta*, *Sambucus hookeri*, *Calamus tenuis*, *Phryniumrheedei*, *Leea indica*, *Phlogocanthuscurvifolius*, *Urena lobata* species. The herbaceous layers were represented by species like *Begonia roxburghii*, *Pteris spp.*, *Diplaziumdilatatum*, *Aolocasia*

acuminata, *Curcuma sp*, *Floscopa scandens*, *Euphorbia hirta*, *Amomum sp*, *Galinsoga parviflora*, *Ipomoea batatas*, *Equisetum debile*, *Ageratum conyzoides*, *Acmella oleracea*, *Asparagus racemosus*. Apart from these compositions of species the sanctuary also represents with some large patch of bamboo mixed forest randomly distributed in most of the areas. Some of the dominant bamboo's species found in the sanctuary are *Dendrocalamus hamiltonii* and *Bambusa pallida*. The practice of shifting cultivation (popularly known as *Jhum cultivation* in the region) has been an integral part of culture and traditions of the local people which is also the major source of vegetables and other food crops. During the field survey, presence of invasive species like *Mikania micrantha*, *Chromolaena odorata*, *Lantana camara*, *Parthenium hysterophorus* was observed in the fallow jhum land. *Chromolaena odorata* and *Parthenium hysterophorus* are also commonly observed along the roads. The occurrence of invasive species in the study areas indicates that the forests are getting disturbed and fragmented, similar other studies conducted earlier for the same study area has also made similar statements [42,43]. The invasive plant species poses serious threats to the native plant community and its ecosystem properties [44]. The extensive field survey conducted in the sanctuary has also observed some rare, threatened and medicinally important species which are distributed with very less population (Table 6). The factors responsible for threatening the rare flora and fauna in the study site are the forest degradation caused by encroachment of forest land for cultivation, building infrastructure and roads. The practice of shorter rotation periods (3-5 years instead of 10-12 years) of Jhum cultivation might have affected the soil nutrient balance and species composition. This also aggravates the risk of soil erosion and frequent landslides in the monsoon season causing habitat fragmentation. The observation

of threats on flora and fauna were also found similar with the findings of others reports [45,46]. The concerns over conservation and sustainable utilization of the resources like medicinal plants were also express by other studies in the state [47,48].

3.4 Anthropogenic Affects on Wildlife

The result on faunal diversity and its interacting with local community was recorded through formal and informal interviews followed by documentation of the events. Among the 8 survey villages only 5 have reported with interaction and conflict with the wildlife. The respondents in Poma village have reported the highest numbers (14 nos.) of interaction and conflict with wildlife followed by Pachin (12 nos.). The reports on human-wildlife conflict were only recoded with Wild Dog, Elephant and King Cobra in Poma, Pachin and Sote villages. Moreover, it also reveals that only 5 species i.e. Monitor Lizard, King Cobra, Mongoose, Wild pig, Jungle cat were the most frequently observed by the local communities during 1990-2019 (Table 7),

these species are reported to be seen regularly in the Jhum land and the nearby forest. However, species like Tiger, Common Leopard, Asiatic Black Bear and Chinese Pangolin were reported to be encountered only during 1990-2000. There was a gradual decrease in interaction of big animals like Tiger, common leopard, wild dog, Asiatic black bear and Elephant according to villagers. One of the villagers in Poma has reported that most of the wild animals were rarely seen at the northern side of the Sanctuary. However, in April, 2020 three individuals from Pachin village have made the headlines in national news for killing a large adult King Cobra. In July, 2020 the King Cobra again made news headline in Itanagar with different ending. An adult King Cobra with a length of 10 feet weighing around 10 kilograms was sighted in the residential area of Itanagar, later it was rescue by the forest department. The regular interaction of wildlife with the local community has been a regular event in and around the Sanctuary. This type of conventional events, are the results of encroachment of forest land for anthropogenic activity causing habitat loss and fragmentation.




Fig. 4. Land surface temperature (LST) of Itanagar wildlife sanctuary in 1990, 2003 and 2019.

Table 5. Correlation coefficient between the biophysical parameters in the year 1990, 2003 and 2019

Environmental Indices	1990	2003	2019
NDVI:LST	-0.42	-0.48	-0.45
NDBI:LST	0.65	0.83	0.69
NDWI:LST	0.30	0.33	0.41
NDVI:NDBI	-0.79	-0.68	-0.86
NDVI:NDWI	-0.98	-0.97	-0.99
NDBI:NDWI	0.68	0.53	0.81




Fig. 5. Biophysical parameters (NDVI, NDBI, and NDWI) in 1990, 2003 and 2019

Table 6. List of important plant species with conservation concern

Sl. No	Scientific Name	Uses	Population Status
1	<i>Abroma augusta</i>	Medicinal Properties: menstrual disorders and gonorrhoea.	Stable
2	<i>Alsophila spinulosa</i>	NA	Decreasing
3	<i>Angiopteris crassipes</i>	NA	Decreasing
4	<i>Aquillariaaagallocha</i>	Essential oil is used in cosmetic and perfume industry.	Threatened**
5	<i>Callicarpa arborea</i>	Medicinal Properties: Stomachache, boils and piles	Decreasing
6	<i>Clerodendrumcolebrookianum</i>	Medicinal Properties: antioxidant and antimicrobial	Decreasing
7	<i>Costusspeciosus</i>	Medicinal Properties: Urinary problem, stimulant and astringent.	Threatened**
8	<i>Dioscorea pentaphylla</i>		Decreasing
9	<i>Dipteriswallichii</i>	NA	Decreasing
10	<i>Dipterocarpus gracilis</i>	Medicinal properties: antiseptic for gonorrhoea and urinary disease	Vulnerable***
11	<i>Gynocordia odorata</i>	Medicinal properties: Diarrhea	Threatened**
12	<i>Hedychium spicatum</i>	Medicinal properties: Anti-inflammatoru. Antispasmodic, and antifungal and antifungal	Threatened**
13	<i>Homalomenaaromatica</i>	Essential oil is used in cosmetic and perfume industry.	Threatened **
14	<i>Hydrocotylejavanica</i>	Medicinal properties: Stomachache, dysentery and antibacterial properties.	Decreasing
15	<i>Livistonajenkinsiana</i>	Traditional housing roof	Decreasing
16	<i>Oroxylum indicum</i>	Medicinal Properties: Stomachache, rheumatism, diarrhoea	Decreasing
17	<i>Piper mullesua</i>	Medicinal Properties: Cough, bronchitis problem and Stomachache	Decreasing
18	<i>Piper pedicellatum</i>	Eaten as vegetables by the locals	Threatened **
19	<i>Terminalia chebula</i>	Medicinal Properties: Dehydration and contipation	Stable
20	<i>Terminalia myriocarpa</i>	Medicinal Properties: Chest pain	Decreasing

Note: **threatened in regional (Ved et al., 2005); ***threatened as per IUCN RED List

Table 7. List of faunal diversity encounter by the local community in and around the wildlife Sanctuary

SI No.	Name of Identified species	Scientific Name and IUCN threatened category	Name of the village/ area	Sighting Year		
				1990-2000	2001-2010	2011-2019
1	Elephant	<i>Elephas maximus</i> ***	A, E, C	████████	████████	
2	Tiger	<i>Panthera tigris</i> ***	A	████████		
3	Common Leopard	<i>Panthera pardus</i> **	A	████████		
4	Asiatic black bear	<i>Ursus thibetanus</i> **	A,D	████████		
5	Clouded leopard	<i>Neofelisnebulosa</i> **	A,B, C	████████	████████	
6	Wild dog	<i>Cuonalpines</i> ***	A,B, C	████████	████████	
7	Jungle cat	<i>Felis chaus</i>	A, B, C, D, E	████████	████████	████████
8	Barking deer	<i>Muntiacusmunjak</i>	A, B, C, D, E	████████	████████	
9	Wild pig	<i>Sus scrofa</i>	A, B, C	████████	████████	████████
10	Chinese Pangolin	<i>Manis pentadactyla</i> ***	C	████████		
11	Mongoose	<i>Herpestes sp.</i>	A, B, C, D, E	████████	████████	████████
12	Indian Porcupine	<i>Hystrix indica</i>	A, B, C,	████████	████████	
13	Monitor Lizard	<i>Varanus bengalensis</i>	A, B, C, D, E	████████	████████	████████
14	King Cobra	<i>Ophiophagus Hannah</i> **	A, B, C, D, E	████████	████████	████████
15	Small Indian Civet	<i>Viverricula indica</i>	A, B, C,	████████	████████	

Note: A- Poma; B- Jollang; C- Pachin; D- Papu Nallah; E- Sote, ***Endangered; **Vulnerable

4. CONCLUSIONS

The application of Remote Sensing (RS) and Geographic Information System (GIS) techniques for assessing changes in land use and land cover in protected areas has proven to be an efficient and invaluable tool. The current study successfully monitored the gradual transformation of land cover into various forms. It revealed a rapid decline in the forest cover of the Itanagar Wildlife Sanctuary from 1990 (77.88%) to 2019 (70.37%), impacting a total area of 17.5 sq. km. Additionally, the built-up area exhibited a substantial increase from 1990 to 2019, measuring 9.37 sq. km, 10.37 sq. km, and 22.86 sq. km, respectively. The research emphasized a correlation between the rise in land surface temperature (LST) and expanding built-up areas, indicative of the formation of an urban heat island (UHI). Correlation analysis further supported a positive correlation between LST and Normalized

Difference Built-Up Index (NDBI) throughout the study years (Table 5).

The study underscored those anthropogenic activities, including Jhum cultivation, forest land encroachment, human-wildlife conflict, rapid invasive species spread, and uncoordinated infrastructure development around the Sanctuary, have significantly impacted biodiversity. Despite Itanagar being the state capital, sustaining economic growth poses a challenge for the Sanctuary. However, achieving a balance between biodiversity conservation and economic development is not unattainable. Both can coexist by promoting sustainable development that prioritizes the restoration and maintenance of ecosystem services. Proposed measures for the Sanctuary encompass habitat evaluation, regular monitoring of wildlife populations and habitats, and the rehabilitation of rare, threatened, and important plants through in-

situ and ex-situ conservation. Sensitization programs aimed at local communities, including awareness and training initiatives, can contribute to effective biodiversity conservation. Furthermore, ecotourism activities such as bird watching, nature walks, and camping can provide livelihood opportunities for the local community. The success of these activities hinges on collaborative efforts involving government departments, research institutes, NGOs, village councils, and other stakeholders, culminating in the formulation of a comprehensive strategy that adheres to the principles outlined above.

ACKNOWLEDGEMENTS

The Authors are thankful to the Department of Environment and Forest, Govt. of Arunachal Pradesh for permission and support. We also thankful to the Chief Wildlife Warden of Itanagar Wildlife Sanctuary for the necessary help and support during fieldwork. The help and information rendered by the local community of different villages is duly acknowledged.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Riebsame WE, Meyer WB, Turner BL. Modeling land use and cover as part of global environmental change. *Climate Change*. 1994;28(1-2):45-64.
2. Threlfall CG, Law B, Banks PB. Sensitivity of insectivorous bats to urbanization: Implications for suburban conservation planning. *Biological Conservation*. 2012; 146(1):41-52.
DOI:10.1016/j.biocon.2011.11.026
3. Adegoke JO, Pielke RA, Eastman J, Mahmood R, Hubbard KG. Impact of irrigation on midsummer surface fluxes and temperature under dry synoptic conditions: A regional atmospheric model study of the US High Plains. *Monthly Weather Review*. 2003;131(3):556-64.
DOI:10.1175/1520-0493(2003)131<0556:IOIOMS>2.0.CO;2
4. Ibrahim F, Rasul G. Urban land use land cover changes and their effect on land surface temperature: Case study using Dohuk City in the Kurdistan Region of Iraq. *Climate*. 2017;5(1):13.
DOI:10.3390/cli5010013
5. Kalnay E, Cai M. Impact of urbanization and land-use change on climate. *Nature*. 2003;423(6939):528-31.
DOI:10.1038/nature01675
6. Grimm NB, Faeth SH, Golubiewski NE, Redman CL, Wu J, Bai X, Briggs JM. Global change and the ecology of cities. *Science*. 2008;319(5864):756-60.
DOI:10.1126/science.115019
7. Arnfield AJ. Two decades of urban climate research: a review of turbulence, exchanges of energy and water, and the urban heat island. *International Journal of Climatology: A Journal of the Royal Meteorological Society*. 2003;23(1):1-26.
DOI:10.1002/joc.859
8. Panigrahy RK, Kale MP, Dutta U, Mishra A, Banerjee B, Singh S. Forest cover change detection of Western Ghats of Maharashtra using satellite remote sensing based visual interpretation technique. *Current Science*. 2010;98(5):657-64.
9. Panaskar S, Nagarajan K, Narwade R. Analysis of Changes in LULC of Western Ghat by Comparing NDVI and NDWI. *Journal of Remote Sensing & GIS*. 2019; 8(4):1-7.
10. Skidmore AK, Bijker W, Schmidt K, Kumar L. Use of remote sensing and GIS for sustainable land management. *ITC Journal*. 1997;3(4):302-315.
Available:https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=6f0280589fcabf2c3d6da8039dce1a5541c23b82
11. Lu D, Moran E, Hetrick S, Li G. Land-use and land-cover change detection. *Advances in environmental remote sensing sensors, Algorithms, and Applications*. 2011;273-290.
12. Xiao H, Weng Q. The impact of land use and land cover changes on land surface temperature in a karst area of China. *Journal of Environmental Management*. 2007;85(1):245-57.
DOI: 10.1016/j.jenvman.2006.07.016
13. Kogan FN. Operational space technology for global vegetation assessment. *Bulletin of the American Meteorological Society*. 2001;82(9):1949-1964.
DOI: 10.1175/1520-0477(2001)082<1949:OSTFGV>2.3.CO;2
14. Rozenstein O, Qin Z, Derimian Y, Karnieli A. Derivation of land surface temperature for Landsat-8 TIRS using a split window algorithm. *Sensors*. 2014;14(4):5768-80.
DOI: 10.3390/s140405768.

15. Sharma R, Chakraborty A, Joshi PK. Geospatial quantification and analysis of environmental changes in urbanizing city of Kolkata (India). *Environmental Monitoring and Assessment*. 2015;187:1-2. DOI:10.1007/s10661-014-4206-7.
16. Bendib A, Dridi H, Kalla MI. Contribution of Landsat 8 data for the estimation of land surface temperature in Batna city, Eastern Algeria. *Geocarto International*. 2017; 32(5):503-13. DOI:10.1080/10106049.2016.1156167.
17. Lodhi MS, Samal PK, Chaudhry S, Palni LM, Dhyanani PP. Land cover mapping for Namdapha National Park (Arunachal Pradesh), India using harmonized land cover legends. *Journal of the Indian Society of Remote Sensing*. 2014;42:461-7. DOI:10.1007/s12524-013-0326-8
18. O'Neill AR. Evaluating high-altitude Ramsar wetlands in the Eastern Himalayas. *Global Ecology and Conservation*. 2019;20:e00715. DOI:10.1016/j.gecco.2019.e00715.
19. Myers N, Mittermeier RA, Mittermeier CG, Da Fonseca GA, Kent J. Biodiversity hotspots for conservation priorities. *Nature*. 2000;403(6772):853-8. DOI:10.1038/35002501.
20. Srivastava RC, Choudhary RK. Floristic scenario of Itanagar wildlife sanctuary-a case study. *Bulletin of Arunachal Forest Research*. 2006;22(1&2):17-21.
21. Kaul RN, Haridasan K. Forest types of Arunachal Pradesh-A preliminary study. *Journal of Economic and Taxonomic Botany*. 1987;9(2):379-89.
22. Champion HG, Seth SK. A revised survey of the forest types of India; 1968.
23. Aiyadurai A, Varma S, Menon V. Human-Predator Conflict in and around Itanagar Wildlife Sanctuary, Arunachal Pradesh, India. Poster presented at Carnivores 2004. Accessed 07 September 2020. Available: https://pdfs.semanticscholar.org/7a1e/54e2f7b0e833507a0688d69901ac5d2ed431.pdf?_ga=2.131967878.824768606.1599367682-2072173107.1596878591
24. Sobrino JA, Jiménez-Muñoz JC, Paolini L. Land surface temperature retrieval from LANDSAT TM 5. *Remote Sensing of Environment*. 2004;90(4):434-40. DOI: 10.1016/j.rse.2004.02.003.
25. Rouse JW, Haas RH, Schell JA, Deering DW. Monitoring vegetation systems in the Great Plains with ERTS. *NASA Spec. Publ.* 1974;351(1):309.
26. Zha Y, Gao J, Ni S. Use of normalized difference built-up index in automatically mapping urban areas from TM imagery. *International Journal of Remote Sensing*. 2003;24(3):583-94. DOI: 10.1080/01431160304987
27. Gao BC. NDWI—A normalized difference water index for remote sensing of vegetation liquid water from space. *Remote Sensing of Environment*. 1996; 58(3):257-66. DOI:10.1016/S0034-4257(96)00067-3.
28. Chen XL, Zhao HM, Li PX, Yin ZY. Remote sensing image-based analysis of the relationship between urban heat island and land use/cover changes. *Remote sensing of Environment*. 2006;104(2):133-46. DOI:10.1016/j.rse.2005.11.016.
29. Malik MS, Shukla JP, Mishra S. Relationship of LST, NDBI and NDVI using landsat-8 data in Kandahimmat watershed, Hoshangabad, India. Accessed 12 September 2020. Available: <http://nopr.niscair.res.in/bitstream/123456789/45657/3/IJMS%2048%281%29%2025-31.pdf>
30. Jansen LJ, Di Gregorio A. Land-use data collection using the "land cover classification system": results from a case study in Kenya. *Land Use Policy*. 2003; 20(2):131-48. DOI: 0.1016/S0264-8377(02)00081-9.
31. Singh P, Kikon N, Verma P. Impact of land use change and urbanization on urban heat island in Lucknow city, Central India. A remote sensing based estimate. *Sustainable Cities and Society*. 2017;32: 100-114. DOI:10.1016/j.scs.2017.02.018.
32. Fatemi M, Narangifard M. Monitoring LULC changes and its impact on the LST and NDVI in District 1 of Shiraz City. *Arabian Journal of Geosciences*. 2019;12:1-2. DOI:10.1007/s12517-019-4259-6.
33. Bayarsaikhan U, Boldgiv B, Kim KR, Park KA, Lee D. Change detection and classification of land cover at Hustai National Park in Mongolia. *International Journal of Applied Earth Observation and Geoinformation*. 2009;11(4):273-80. DOI:10.1016/j.jag.2009.03.004.
34. Menon V, Daniel JC. Field guide to Indian mammals. Dorling Kindersley, India in association with Penguin Book, India; 2003.

35. Congalton RG. A review of assessing the accuracy of classifications of remotely sensed data. *Remote sensing of environment*. 1991;37(1):35-46. DOI:10.1016/0034-4257(91)90048-B.
36. Population.city.com (2020). Accessed 27 August 2020. Available:<http://population.city/india/itanagar/>
37. Yuan F, Sawaya KE, Loeffelholz BC, Bauer ME. Land cover classification and change analysis of the Twin Cities (Minnesota) Metropolitan Area by multitemporal Landsat remote sensing. *Remote sensing of Environment*. 2005; 98(2-3): 317-28. DOI:10.1016/j.rse.2005.08.006.
38. Congalton RG, Green K. Assessing the accuracy of remotely sensed data: principles and practices. CRC Press; 2019.
39. Oke, T.R. *Boundary Layer Climates*. 2nd ed.; Methuen: London, UK, 1987;435.
40. Oke RT. Urban climate and global environmental change. *Applied Climatology*. 1997;273-87.
41. Chandler TJ. London's Urban Climate. *The Geographical Journal*, 1962;128(3):279-. DOI:10.2307/1794042.
42. Kosaka Y, Saikia B, Mingki T, Tag H, Riba T, Ando K. Roadside distribution patterns of invasive alien plants along an altitudinal gradient in Arunachal Himalaya, India. *Mountain Research and Development*. 2010;30(3):252-8.
43. Ronald K, Gajurel PR, Singh B. Assessment of eco-diversity status of *Homalomenaaromatica* (Spreng.) Schott and its habitat in tropical forest of Indian Eastern Himalaya. *Plant Science Today*. 2019;6(2):71-83.
44. Mooney HA, Hamburg SP, Drake JA. The invasions of plants and animals into California. In *Ecology of biological invasions of North America and Hawaii*. New York, NY: Springer New York. 1986;250-272.
45. Kumar A, Sarma K, Krishna M, Devi A. The eastern hoolock gibbon (*Hoolock leuconedys*) in eastern Arunachal Pradesh, India. *Primate Conservation*. 2013; (27):115-23. DOI:10.1896/052.027.0106.
46. Kumari R, Banerjee A, Kumar R, Kumar A, Saikia P and Khan ML. Deforestation in India: Consequences and Sustainable Solutions, In *Forest Degradation Around the World*, by Suratman MZ, Latif ZA, Oliveira GD, Brunzell N, Shimabukuro Y & Santos CACD, Intech Open; 2019. DOI:10.5772/intechopen.85804.
47. Ved DK, Kinhal GA, Ravikumar K, Sankar RV, Haridasan K. Conservation Assessment and Management Prioritisation (CAMP) for wild medicinal plants of North-East India. *Medicinal Plant Conservation*. 2005;11:40-4.
48. Perme N, Choudhury SN, Choudhury R, Natung T, De B. Medicinal plants in traditional use at Arunachal Pradesh, India. *International Journal of Phytopharmacy*. 2015;5(5):86-98.

© 2023 Ronald et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/110457>