



Varietal Screening and Management of Basal Rot Disease of Onion

Sirisha Thakare ^{a*}, Kolase S. V. ^a, Dawale M. B. ^a,
Narute T. K. ^a, Navale A. M. ^a and Chandanshive A. V. ^b

^a Department of Plant Pathology and Microbiology, PGI, MPKV, Rahuri (MH) - 413722, India.

^b Tomato Improvement Project, PGI, MPKV, Rahuri (MH) – 413722, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i234261

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/110688>

Original Research Article

Received: 13/10/2023

Accepted: 19/12/2023

Published: 22/12/2023

ABSTRACT

Onion (*Allium cepa* L.) belongs to the family Amaryllidaceae. It is known as common onion, bulb onion or biennial herb, and is one of the most important vegetable crops in India. However, the crop faces numerous diseases attack, resulting in losses to both quantity and quality. Among the diseases, the basal rot caused by *Fusarium oxysporum* f. sp. *cepae*, is one of the most significant soil-borne diseases. This work aimed to screen various onion varieties for resistance against basal rot in pots under glasshouse condition as well as to assess the efficacy of fungicides against the pathogen under both in vitro and in vivo conditions. It was observed that all onion varieties screened were highly susceptible while Bhima red was just susceptible; none of the varieties were resistant to basal rot. Under in vitro conditions the Fungicides Carbendazim and Tebuconazole, of the given concentration, were cent per cent effective in inhibiting mycelial growth of *Fusarium oxysporum* f. sp. *cepae*. For the field condition, the module of [carbendazim (2 g/kg) followed by tebuconazole (0.1 %) followed by azoxystrobin (0.1 %)] was the most effective.

Keywords: Onion; basal rot; *Fusarium oxysporum* f. sp. *cepae*.

*Corresponding author: E-mail: sirishathakare09@gmail.com;

1. INTRODUCTION

Onion (*Allium cepa* L., $2n = 2x = 16$) a member belongs to the family *Amaryllidaceae* is known as common onion or bulb onion, biennial herb, is one of the most important vegetable crops in India [1]. It is a low latitude horticulture crop and is popularly known as “Queen of Kitchen” [2]. The area, production and productivity of onion in India during the year 2022-23 was 1791.73 thousand ha, 31005.41 thousand tonne and 17.3 MT/ha respectively [3]. Maharashtra was leading state in onion production throughout 2021-22 having 13,301.70 t followed by Madhya Pradesh (4740.60 t) and Karnataka (2779.50 t) [4].

Numerous diseases attack the onion crop, resulting in losses to both quantity and quality. Among the diseases, basal rot of onion which is caused by *F. oxysporum* f. sp. *cepae*, is one of the most significant soil-borne diseases of onion [5], which affects the crop at all stages of development, yield losses up to 50% were recorded in susceptible cultivars [6], 90% losses reported at the seedling stage [7] and 30% losses in the storage [8]. In India, the occurrence of this disease was first reported from Rajasthan [9].

Early symptoms of the disease in the field include curling, yellowing and wilting and the affected plant shows yellowing followed by complete drying of foliage. The bulb of affected plant shows soft rotting and the root decay. The infected plants can be uprooted easily. There will be a whitish mouldy growth on the scale. In addition, when the bulbs are sliced open in excessively hot and dry conditions, diseased bulbs turn discoloured, infected plants wilt, and infected tissue appears brown and watery. The present studies were conducted to screen various onion varieties for resistance against basal rot in pots under glasshouse condition as

well as to assess the efficacy of fungicides against the pathogen *in vitro* and *in vivo*.

2. MATERIALS AND METHODS

2.1 Screening of Onion Varieties against Basal Rot Disease

Onion varieties were collected from Directorate of Onion and Garlic Research, Rajgurunagar, Pune, were tested for resistance against basal rot disease of onion during *Kharif-2022* and 2023. In this experiment, total 23 varieties were screened under artificial epiphytotic conditions under glasshouse, Department of Plant Pathology and Microbiology, PGI, MPKV, Rahuri. Three replications were maintained for each treatment.

Fusarium oxysporum f. sp. *cepae* inoculum was multiplied on sand-maize meal agar medium for the pathogenicity test. The medium was prepared by autoclaving 90 g of dry sieved sand and 10 g of maize meal with 40 ml of distilled water for 15-20 minutes at 1.1 kg/cm² or 15 PSI pressure. In order to achieve uniform growth, the sterilized medium was inoculated with five days old culture of test pathogen and incubated for 14 days at $25 \pm 1^\circ\text{C}$. The inoculums thus prepared and added to the sterilized soil and FYM (3:1) potting mixture @ 10 per cent (w/w) by mixing it with upper layer of soil and allowed for 7 days to infest soil. Forty five days old onion seedlings of all 23 varieties of onion were transplanted into sick soil pots. Periodic observations were recorded on incidence of basal rot by using disease rating scale given by Wellman (1939) and Harrison (1940) as mentioned in the Table 1. The wilt incidence was calculated by using a formula as suggested by Mayee and Datar [10].

$$\text{Per cent wilt incidence} = \frac{\text{Number of wilted plants}}{\text{Total number of plant}} \times 100$$

Table 1. Disease rating scale for basal rot disease of onion

Sr. No.	Reaction/grade	Per cent/Rating (%)
1.	Total Resistance (TR)	0
2.	Highly Resistance (HR)	1-10
3.	Moderately Resistance (MR)	11-30
4.	Moderately Susceptible (MS)	31-50
5.	Susceptible (S)	51-70
6.	Highly susceptible (HS)	71-100

2.2 In vitro Efficacy of Fungicides against *Fusarium oxysporum* f. sp. *cepae*

In vitro studies were undertaken at Laboratory, Department of Plant Pathology and Microbiology, PGI, MPKV, Rahuri following completely randomized design with seven treatments and three replications. Six fungicides (Table 3) were tested against *Fusarium oxysporum* f. sp. *cepae* by using poisoned food technique. Control was maintained without fungicide. The plates were incubated at 25 ± 1°C. The colony's radial growth was recorded and per cent (%) growth inhibition was determined using Vincent's method [11] when the mycelial growth in control plate reached to 90 mm.

$$I = \frac{C - T}{C} \times 100$$

Where,

- I = Per cent inhibition
- C = Radial growth of fungus in control
- T = Radial growth of fungus in treatment

2.3 In vivo Evaluation of Fungicides and Bio-agents (Modules) against Basal Rot Disease of Onion during the Monsoon: Kharif-2022 and 2023

The experiment was conducted on the experimental farm of the Department of Plant Pathology and Microbiology, PGI, MPKV, Rahuri. The onion cultivar "Phule Samarth" was grown following randomized block design (RBD) with nine treatments and three replications (Table 4). Onion seeds were treated with fungicide and raised in nursery. Forty five days after sowing, the seedlings were transplanted in main field after pre-plant dipping with bioagents/fungicides. Spacing of 15 cm × 10 cm was maintained. fungicides drenching was done 30 days after transplanting. Control plots were maintained without any treatment. Disease incidence was recorded from appearance of disease in field upto harvesting of the crop by following rating scale given by Wellman (1939) and Harrison (1940).

2.4 Statistical Analysis

The methodologies outlined by Panse and Sukhatme [12] were used for statistical analysis during the studies.

3. RESULTS AND DISCUSSION

3.1 Screening of Onion Varieties for Resistance against Basal Rot Disease under Glasshouse Condition

Twenty-three varieties of onion were evaluated for resistance to *Fusarium oxysporum* f. sp. *cepae* in glasshouse by using sick pot method. The results of these studies carried out in pot culture are presented in Table 2. All the varieties were found highly susceptible to basal rot disease of onion caused by pathogen *F. oxysporum* f. sp. *cepae* while Bhima red was found susceptible. Average wilt incidence was ranged between 70 to 100 per cent during both years i.e. Kharif-2022 and 2023. Highest mean wilt incidence (100 %) was recorded on cultivar NHRDF red -03, N-2-4-1, Palampur white and PKV white. While lowest mean wilt incidence (70%) was recorded on cultivar Bhima red. The data revealed that all the onion varieties tested in the study were susceptible to basal rot.

Ilhe et al. [1] screened genotypes of onion against *F. oxysporum* f. sp. *cepae* under artificial epiphytotic condition. Among the genotypes screened, the highest disease incidence was found in N-2-4-1 (100 %) followed by Bhima Shakti (90 %), Bhima Super (90 %), Phule Samarth (90 %), B-780 (90 %) and Bhima Shubhra (90 %). The genotypes Bhima Kiran, Bhima Raj, Bhima Shweta exhibited 80 per cent disease incidence. However, lowest incidence was observed on Bhima red (60 %).

3.2 In vitro Evaluation of Fungicides against *Fusarium oxysporum* f. sp. *cepae*

The results presented in Table 3 showed that, the fungicides carbendazim and tebuconazole of the given concentration were cent per cent effective in inhibiting mycelial growth of *Fusarium oxysporum* f. sp. *cepae*. However, the fungicides viz., Azoxystrobin (0.1%), Thiophanate methyl (0.1%), Metiram 55 % + Pyraclostrobin 5 % (0.1%) and Fludioxonil + Sedaxane (0.1%) alone inhibited 87.03, 83.70, 79.63 and 56.29 per cent growth of the pathogen respectively. This shows that Fludioxonil + Sedaxane was less effective.

Table 2. Screening of different varieties of onion against basal rot using sick pot method under glass-house condition during Kharif-2022 and 2023 (Pooled data)

Sr. No.	Variety	Kharif-2022 (Wilt incidence %)	Kharif-2023 (Wilt incidence %)	Mean incidence (%)	Disease Reaction
1.	Agrifound Dark Red	90	90	90	Highly Susceptible
2.	Agrifound White	80	90	85	Highly Susceptible
3.	Arka Kalyan	90	80	85	Highly Susceptible
4.	Arka Yojith	90	90	90	Highly Susceptible
5.	Bhima Dark Red	80	80	80	Highly Susceptible
6.	Bhima Raj	80	90	85	Highly Susceptible
7.	Bhima Red	70	70	70	Susceptible
8.	Bhima Kiran	80	80	80	Highly Susceptible
9.	Bhima Safed	90	90	90	Highly Susceptible
10.	Bhima Shubhra	90	80	85	Highly Susceptible
11.	Bhima Shweta	80	80	80	Highly Susceptible
12.	Bhima Super	90	80	85	Highly Susceptible
13.	GWO-1	90	90	90	Highly Susceptible
14.	NHRDF Red-3	100	100	100	Highly Susceptible
15.	NHRDF Red-4	80	90	85	Highly Susceptible
16.	N-2-4-1	100	100	100	Highly Susceptible
17.	Palampur White	100	100	100	Highly Susceptible
18.	Phule Baswant	90	80	85	Highly Susceptible
19.	Phule Samarth	90	90	90	Highly Susceptible
20.	PKV White	100	100	100	Highly Susceptible
21.	Pusa Madhavi	80	90	85	Highly Susceptible
22.	Pusa Shobha	90	80	85	Highly Susceptible
23.	Pusa White Round	90	90	90	Highly Susceptible

Table 3. Evaluation of fungicides against *Fusarium oxysporum* f. sp. *Cepae*

Tr. No.	Fungicides	Concentration (%)	Mean colony diameter (mm*) after 7 days of inoculation	Per cent growth inhibition
T ₁	Carbendazim 50 % WP	0.1	0.00	100.00
T ₂	Metiram 55 % + Pyraclostrobin 5 % WG	0.1	18.33	79.63
T ₃	Azoxystrobin 23 % SC	0.1	11.67	87.03
T ₄	Thiophanate methyl 70 % WP	0.1	14.67	83.70
T ₅	Tebuconazole 25.9 % EC	0.1	00.00	100.00
T ₆	Fludioxonil + Sedaxane	0.1	39.33	56.29
T ₇	Control	-	90.00	00.00
	S.Em. (±)	-	0.25	-
	CD at 1%	-	0.76	-

* = Average of three replications

Yadav et al. [13], evaluated seven systemic fungicides against *F. oxysporum* f. sp. *cepae* and reported that, tebuconazole and carbendazim were most effective showing 100 % mycelial growth inhibition, followed by thiophanate methyl showing 90.03 per cent

growth inhibition. Chethan et al. [8] who evaluated the efficacy of seven fungicides against *F. oxysporum* f. sp. *cepae* and reported that, tebuconazole was effective showing 92.59 per cent mycelial growth inhibition.

Table 4. *In vivo* disease management of basal rot of onion during *Kharif-2022* and *2023* (Pooled data)

Tr. No.	Treatments	% disease incidence of basal rot of onion					Yield (q/ha)	PDC over control
		15 DAS	Before Transplanting	30 DAT	60 DAT	90 DAT		
T ₁	Carbendazim (2 g/kg) followed by (Metiram+Pyraclostrobin) (0.1 %) followed by Azoxystrobin (0.1 %)	6.00 (14.17)	7.83 (16.25)	10.17 (18.59)	12.33 (20.55)	14.17 (22.11)	118.05	59.12
T ₂	Carbendazim (2 g/kg) followed by (Metiram + Pyraclostrobin) (0.1 %) followed by Thiophanate methyl (0.3 %)	6.00 (14.17)	7.83 (16.25)	11.67 (19.97)	13.50 (21.55)	15.50 (23.18)	112.5	55.29
T ₃	Carbendazim (2 g/kg) followed by Tebuconazole (0.1 %) followed by Azoxystrobin (0.1 %)	6.00 (14.17)	7.83 (16.25)	9.17 (17.62)	10.83 (19.22)	12.33 (20.55)	128.33	64.43
T ₄	Carbendazim (2 g/kg) followed by Tebuconazole (0.1 %) followed by Thiophanate methyl (0.3 %)	6.00 (14.17)	7.83 (16.25)	9.83 (18.27)	11.33 (19.67)	13.17 (21.27)	123.61	62.01
T ₅	(Fludioxonil+Sedaxane) (0.1 %) followed by <i>Trichoderma viridae</i> (5 g/kg) followed by Azoxystrobin (0.1 %)	8.5 (16.95)	10.50 (18.90)	13.17 (21.28)	14.50 (22.38)	16.67 (24.09)	106.66	51.91
T ₆	(Fludioxonil + Sedaxane) (0.1 %) followed by <i>Trichoderma viridae</i> (5 g/kg) followed by Thiophanate methyl (0.3 %)	8.5 (16.95)	10.50 (18.90)	13.67 (21.69)	15.17 (22.92)	17.50 (24.72)	99.44	49.52
T ₇	(Fludioxonil + Sedaxane) (0.1 %) followed by <i>Trichoderma harzianum</i> (5 g/kg) followed by Azoxystrobin (0.1 %)	8.5 (16.95)	10.50 (18.90)	15.50 (23.18)	18.00 (25.10)	21.33 (27.51)	88.60	38.47
T ₈	(Fludioxonil+Sedaxane) (0.1 %) followed by <i>Trichoderma harzianum</i> (5 g/kg) followed by Thiophanate methyl (0.3 %)	8.5 (16.95)	10.50 (18.90)	17.17 (24.48)	19.50 (26.21)	24.33 (29.55)	84.44	29.82
T ₉	Control	12.83 (20.99)	18.33 (25.47)	28.83 (29.22)	29.83 (33.11)	34.67 (36.07)	55.27	00.00
	S.Em. (±)	0.27	0.32	0.30	0.37	0.44	1.77	-
	CD at 5%	0.82	0.97	0.90	1.11	1.32	5.33	-

(Figures in parenthesis are angular transformed values, DAS: Days After Sowing, DAT: Days After Transplanting)

3.3 In vivo Disease Management of Basal Rot of Onion during Monsoon: Kharif-2022 and 2023 (Pooled Data)

The basal rot disease of onion over control was reduced by all treatments as compared to control. Two years cropping season pooled data presented in Table 4, revealed that, after 90 DAT, the per cent disease incidence was ranged from 12.33 to 24.33 per cent as compared to control (34.67 %). Least incidence was observed in treatment T₃ [carbendazim (2 g/kg) followed by tebuconazole (0.1 %) followed by azoxystrobin (0.1 %)] (12.33 %) and T₄ [carbendazim (2 g/kg) followed by tebuconazole (0.1 %) followed by thiophanate methyl (0.3 %)] (13.17 %) which were statistically at par with each other. It was then followed by T₁ [carbendazim (2 g/kg) followed by (metiram+pyraclostrobin) (0.1 %) followed by azoxystrobin (0.1 %)] (14.17 %) and T₂ [carbendazim (2 g/kg) followed by (metiram + pyraclostrobin) (0.1 %) followed by thiophanate methyl (0.3 %)] (15.50 %). Maximum disease incidence (34.67 %) was found in T₉ (Control). Maximum per cent disease control over control (64.43 %) and yield (128.33 q/ha) was found in treatment T₃.

These research findings are confirmatory with Sintayehu et al. [14] who dipped shallot bulb in Prochloraz, Tebuconazole, Carbendazim, Mancozeb for bulb treatment and found that Prochloraz and Carbendazim reduced basal rot incidence (40 and 43 %). Similarly, Mondani et al. [15] evaluated the efficacy of commercial chemicals and bioagents, for reducing disease incidence of *Fusarium* spp. in garlic in the field stage and found that Tebuconazole was the most effective for reducing disease incidence, showing 26.5 per cent reduction of basal plate rots and 44 per cent reduction of bulb rots and 33.4 per cent reduction in visible symptoms on cloves relative to the untreated controls [16].

4. CONCLUSION

In the present study, total 23 onion cultivars that were screened against basal rot disease of onion were found highly susceptible while, one variety 'Bhima red' was found susceptible to the disease. Among six fungicides, tebuconazole (0.1 %) and carbendazim (0.1 %) were found effective. Modules i.e., [carbendazim (2 g/kg) followed by tebuconazole (0.1 %) followed by

azoxystrobin (0.1 %)] and [carbendazim (2 g/kg) followed by tebuconazole (0.1 %) followed by thiophanate methyl (0.1 %)] were found statistically at par with each other and effective against basal rot disease of onion under field conditions.

ACKNOWLEDGEMENTS

The author expressed gratitude for the support and resources offered by the Department of Plant Pathology and microbiology at Mahatma Phule Krishi Vidyapeeth, Rahuri, and ICAR-Directorate of Onion and Garlic Research Pune, which were instrumental in facilitating the research.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ilhe BM, Musmade NA, Kawade SB. Management of basal bulb rot of onion (*Allium cepa* L.). Int J Plant Prot. 2013; 6(2):349-352.
2. Griffiths G, Trueman L, Crowther T, Thomas B, Smith B. Onions: a global benefit to health. Phytother Res. 2002; 16(7):603-615.
3. Anonymous. Agricultural Statistics at a Glance-2022. Department of Agriculture & Farmers Welfare; 2022. Available:www.agricoop.nic.in
4. Anonymous. India Stat Agri-2022. Horticulture; 2022. Available:indiastatagri.com
5. Jagtap JD, Suryawanshi NS. Potential of biocontrol agents against basal rot of onion caused by *Fusarium oxysporum* f. sp. *cepae*. Int J Life Sci. 2015;A5:65-69.
6. Evert KL, Schwartz HF, Epsky ND, Capinera JL. Effect of maggots and wounding on occurrence of *Fusarium* basal rot of onion in Colorado. Plant Dis. 1985;69:878-882.
7. Davis GM, Reddy CS. A seedling blight stage of onion bulb rot. Phytopathol. 1983;73:22-28.
8. Barnocakine SE. Possibilities to control *Fusarium* in onion Zold segtements kutote. Intezet Bulletinjo. 1986;19-33.

9. Mathur, Sankhala. Observed the first time the bulb rot in Indian book: Disease problem in vegetable production by S. K. Gupta and T. S. Thind. 1963;359-360.
10. Mayee CD, Datar VV. Phytopathometry, Marathwada Agricultural University, Parbhani. 1986;146.
11. Vincent JM. Distortion of fungal hyphae in presence of certain inhibitors. Nature. 1927;159:850.
12. Panse VS, Sukhatme PV. Statistical methods for agricultural workers. ICAR. New Delhi. 1985; pp. 97-156.
13. Yadav GMB, Mishra JP. *In vitro* study of onion basal rot caused by *Fusarium Oxyporum*. J Emerg Technol Innov Res. 2020;7(5):2349-5162.
14. Sintayehu A, Sakhuja PK, Fininsa C, Ahmed S. Management of *Fusarium* basal rot (*Fusarium oxysporum* f. sp. *cepae*) on shallot through fungicidal bulb treatment. Crop Prot. 2011;30:560–565.
15. Mondani L, Chiusa G, Battilani P. Efficacy of chemical and biological spray seed treatments in preventing garlic dry rot. Phytopathol Mediterr. 2022;61(1):27-37.
16. Chethan P, Lohate SR, Hasabnis SN, Bansode GM. *In vitro* evaluation of fungicides against the pathogens associated with post-harvest bulb rot of onion. J Pharma Innov. 2021;10(10):2290-2294.

© 2023 Thakare et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/110688>