



Oxidative Stress and Diabetes Mellitus: Unravelling the Intricate Connection: A Comprehensive Review

**Nadeem Rais ^{a+++*}, Akash Ved ^{b#}, Rizwan Ahmad ^{c†}
and Aashna Parveen ^{d‡}**

^a Department of Pharmacy, Bhagwant University, Ajmer, Rajasthan 305004, India.

^b Faculty of Pharmacy, Dr. A.P.J. Abdul Kalam Technical University, Lucknow, Uttar Pradesh 226031, India.

^c Department of Pharmacy, Vivek College of Technical Education, Bijnor, Uttar Pradesh 246701, India.

^d Faculty of Applied Science, Bhagwant Global University, Kotdwar, Uttarakhand 246149, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2024/v36i17493

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/112366>

Review Article

Received: 18/11/2023

Accepted: 24/01/2024

Published: 31/01/2024

ABSTRACT

Background: Diabetes mellitus (DM) is often associated with oxidative stress (OS), which is defined as an imbalance between the body's antioxidant defense systems and the generation of reactive oxygen species (ROS). OS serves as a crucial factor in the intricate relationship between DM and cellular dysfunction, influencing the generation of ROS and subsequent DM complications such as retinopathy, cardiomyopathy, neuropathy, nephropathy, encephalopathy, and peripheral arteriopathy.

⁺⁺ Assistant Professor;

[#] Professor and Head;

[†] Associate Professor;

[‡] Ph.D. Research Scholar;

*Corresponding author: E-mail: drx.nadeemrais@gmail.com;

Objective: This comprehensive review aims to elucidate the complex interplay between OS and DM, providing a thorough understanding of the underlying mechanisms and highlighting emerging therapeutic interventions for the management of OS-related complications in DM. It also explores novel antioxidant-based therapies aiming at specific OS markers and developing personalized interventions, which represents a promising avenue for enhancing treatment efficacy in DM.

Method: The search was conducted on scientific databases and web portals such as PubMed, ScienceDirect, Web of Science, Embase, Google Scholar, EBSCO, DOAJ, etc.

Conclusion: In conclusion, OS and DM are related through a dynamic and intricate interaction involving genetic, molecular, and environmental variables. Interdisciplinary approaches hold the potential to uncover novel biomarkers for early detection, prognosis, and oriented therapeutic interventions, thereby revolutionizing the clinical management of DM-related complications. With research continuing to advance and customized treatments being more widely incorporated into clinical practice, there is hope that the impact of OS-related DM complications will be significantly mitigated in the future. Despite notable progress, certain unexplored facets necessitate deeper investigations into the precise mechanisms through which OS exacerbates the progression of DM.

Keywords: Oxidative stress; diabetes mellitus; reactive oxygen species; insulin resistance; diabetic complications.

1. INTRODUCTION

A physiological state known as oxidative stress (OS) is defined by an imbalance between the body's ability to neutralize reactive oxygen species (ROS) and the amount of ROS it produces. ROS are very reactive substances that can damage deoxyribonucleic acid (DNA), proteins, and lipids. Examples of these molecules are superoxide radicals, hydrogen peroxide, and hydroxyl radicals [1]. Under normal conditions, the body possesses an advanced defense mechanism that includes enzymes, antioxidants, and other substances to offset the damaging effects of ROS. On the other hand, OS develops when the body's ability to combat ROS is outmatched by their creation. This condition can cause tissue damage and cellular dysfunction and contribute to the onset and advancement of a number of diseases, including diabetes mellitus (DM) [2] (Fig. 1).

Within the context of DM, OS is a major player in the disease's aetiology. It has a strong connection to the emergence of DM issues, beta-cell malfunction, and insulin resistance. In order to develop oriented treatment strategies to reduce the consequences of OS, it is imperative to understand how OS affects the progression and complications of DM [3,4]. We attempted to clarify the intricate relationship between OS and DM in this comprehensive review, offering a clear grasp of the underlying mechanisms and highlighting recently developed therapeutic approaches for the treatment of complications in DM related to OS.

2. THE ROLE OF OS IN DM DEVELOPMENT

DM develops primarily as a result of oxidative damage. ROS are produced in excess due to several mechanisms, such as mitochondrial malfunction and nicotinamide adenine dinucleotide phosphate (NADPH) oxidase activity [5]. Consequently, lipid peroxidation (LPO), protein modification, and DNA damage cause cellular destruction. Important components in the pathophysiology of DM, insulin resistance, and decreased beta-cell activity are caused by these processes [6]. To create oriented therapy solutions, it is imperative to comprehend these complex systems.

2.1 Mechanisms of OS generation

A complicated web of biological processes that result in the body producing too many ROS is what causes OS [7]. Several mechanisms contribute to the generation of OS in the context of DM (Fig. 2). DM-related OS is caused by a variety of factors. ROS generation is increased by mitochondrial malfunction, and oxidative damage is intensified by advanced glycation end products (AGEs). Increased ROS levels are caused by elevated NADPH oxidase activity, and the oxidative load is increased by activating the polyol pathway. Importantly, disturbances in the balance of metal ions and inflammatory pathways work together to cause damage to cells and advance issues associated with DM [7,8]. Generally termed the "powerhouses" of the cell, mitochondria are essential for oxidative

phosphorylation, the process that produces energy in cells. But they also contribute significantly to ROS production. High blood sugar in diabetics can cause mitochondrial malfunction, which raises the generation of ROS [9]. AGEs can develop as a result of elevated blood glucose levels, arising from the chemical process between carbohydrates and proteins. These molecules cause direct tissue damage and induce the formation of ROS, contributing to OS [10]. NADPH oxidase, an enzyme complex producing superoxide radicals, sees elevated activity in DM, leading to increased ROS synthesis, especially impactful in vascular tissues due to its role in endothelial dysfunction [11]. The polyol pathway, an additional glucose metabolism mechanism, results in sorbitol and fructose accumulation in DM, exacerbating OS by utilizing NADPH, necessary for antioxidant production [12]. In DM, chronic inflammation characterizes the activation of cyclooxygenase and lipoxygenase, contributing to ROS formation through inflammatory pathways [13]. Transition

metals like iron and copper, imbalanced in DM, catalyze processes like Fenton and Haber-Weiss, further generating damaging radicals [14].

2.2 Cellular Damage and Dysfunction

The pathophysiology of DM and its consequences is aided by OS, which has deleterious effects on cellular structures and activities. Cell membrane integrity is compromised by LPO. Protein mutation influences the structure and function of the protein, affecting vital biological functions. Mutations and genomic destabilization can result from damage to DNA. ROS formation is made worse by mitochondrial malfunction, which impairs energy production. Protein folding and cellular equilibrium are interfered with by endoplasmic reticulum (ER) stress. Cellular discomfort is exacerbated by misregulated signaling pathways. DM problems arise from a lack of cellular balance brought on by the




Fig. 1. Relationship between OS and DM: A physiological state known as OS is defined by an imbalance between the body's generation of ROS and its capacity to counteract them with antioxidants. This condition can lead to cellular malfunction and tissue damage, which can initiate and worsen a variety of diseases, including diabetes mellitus and its complications




Fig. 2. The suggested mechanism through which OS is produced in relation to diabetic complications: Endoplasmic reticulum stress-induced lipotoxicity and mitochondrial dysfunction are factors that lead to insulin resistance in target tissues, pancreatic β -cell failure, and diabetic consequences. Adopted and modified from Luong et al. (2017) [7]

cumulative effect [15,16]. LPO, induced by ROS, produces lipid peroxides and other hazardous byproducts that can damage cell membrane integrity, potentially compromising cell viability and leading to cell death [17]. Protein amino acids reacting with ROS generate AGEs, nitrotyrosine, and protein carbonyls, causing altered proteins to lose typical structure and functionality, leading to cellular dysfunction [18]. DNA strands suffer direct impact from ROS or secondary reactive species, resulting in degradation, crosslinks, alterations, and strand breakage, potentially causing mutations and genomic destabilization if not properly repaired [19,20]. Mitochondria, major ROS producers, are highly susceptible to oxidative damage, which hampers oxidative phosphorylation and energy synthesis, exacerbating cell malfunction [21]. OS compromising ER function leads to protein misfolding, triggering the unfolded protein response, associated with insulin resistance and dysfunctional beta cells [22]. OS affects cellular signaling pathways involving apoptosis, inflammation, and cellular survival, exacerbating cellular damage and dysfunction [23]. The disruption of cellular homeostasis by cumulative degradation and dysfunction results in altered ion gradients, impaired cellular metabolism, and disturbances in intracellular communication [24].

3. OS BIOMARKERS IN DM

Keeping an eye on biomarkers of OS in DM offers vital information about how the condition advances. An increased oxidative load is indicated by elevated ROS and reactive nitrogen species (RNS). Cellular membrane damage can be physically detected by LPO products [25]. Assessing antioxidant enzyme activity unveils the system's capacity to counteract OS. These biomarkers collectively offer a comprehensive assessment of OS levels, aiding in the development of targeted therapeutic interventions for individuals with DM.

3.1 ROS and RNS linked products

Significant roles are played by highly reactive substances known as ROS and RNS in OS and its consequences for DM [25]. ROS encompass various radicals such as superoxide radicals generated in the mitochondrial electron transport chain during cellular respiration, and by enzymes like NADPH oxidase, xanthine oxidase, and cytochrome P450 [26]. Hydrogen peroxide (H_2O_2), though less reactive than superoxide, still poses significant cell harm, produced enzymatically or via superoxide dismutation [27]. Hydroxyl radicals ($\cdot OH$) are produced via the Fenton and Haber-Weiss reactions, which

involve transition metals like iron and copper, and are the most aggressive and damaging ROS [28]. RNS include nitric oxide (NO), crucial for signaling and vasodilation but can combine with superoxide under OS, forming peroxynitrite (ONOO⁻) [29], a potent oxidant causing inflammation and cellular damage [30]. As previously stated, this particular enzyme complex plays a significant role in the formation of superoxide. It is activated in OS circumstances and participates in multiple cellular functions [31]. The two most important isoforms of nitric oxide synthase (NOS) in relation to DM are endothelial NOS (eNOS) and inducible NOS (iNOS). They generate NO, which may be part of pathways that are harmful or beneficial [32,33]. The Mitochondrial electron transport chain (METC), especially during malfunction, becomes a substantial ROS source via superoxide production from oxidative phosphorylation electron leakage. Inflammatory cells like neutrophils and macrophages produce ROS as part of the immune response, intensifying DM-related inflammation and tissue damage. METC is a major ROS source, especially when there is mitochondrial malfunction. Superoxide can be produced when oxidative phosphorylation involves electron leakage [34]. ROS can be produced by inflammatory cells as a component of the immune response, including neutrophils and macrophages. This exacerbates DM-related inflammation and tissue damage [35].

3.2 LPO Products

One important mechanism by which OS damages cellular structures is LPO. It involves the oxidative breakdown of lipids, especially polyunsaturated fatty acids, which produces a number of hazardous byproducts. LPO products are a major cause of tissue damage and cellular dysfunction in the context of DM [36]. One of the most widely recognized and extensively researched byproducts of LPO is malondialdehyde (MDA). It is produced when polyunsaturated fatty acids break down and is a trustworthy indicator of OS. Elevated peroxidation of lipids and oxidative deterioration are indicated by elevated MDA levels [37]. Another extremely reactive aldehyde produced during LPO is 4-hydroxy-2-nonenal (4-HNE). It can interact with proteins, DNA, and different cellular constituents to generate adducts that impede function and change the structure of the cell. 4-HNE has been linked to a number of degenerative diseases, such as DM [38]. Isoprostanes (IsoPs) are substances that

resemble prostaglandins and are created when arachidonic acid undergoes non-enzymatic peroxidation. They function as trustworthy indicators of OS and LPO in biological tissues. DM patients have higher-than-normal quantities of IsoPs in their bodies [39]. LPO produces the highly reactive and deadly chemical acrolein. It has the ability to react with DNA and proteins, causing damage to DNA and malfunctioning of cells. DM and other disorders linked to OS have been linked to elevated levels of acrolein [40]. Lipid hydroperoxides are the early phases of the oxidative destruction of lipids and are intermediate products of LPO. They have the ability to interact further with other parts of the cell, intensifying the harmful consequences of OS [41]. The oxidized counterparts of cholesterol are called oxysterols. They are created when interactions with ROS occur, and they can negatively impact cell membranes by changing their permeability and fluidity [42].

3.3 Antioxidant Enzymes

The body's defense against OS depends on antioxidant enzymes because they neutralize ROS and prevent them from damaging cells. Changes in these enzymes' activity can have a major effect on the equilibrium between antioxidant defense mechanisms and OS in the context of DM [6]. Superoxide dismutase (SOD) is in charge of converting superoxide radicals into less dangerous compounds like oxygen (O₂) and H₂O₂. SOD comes in a variety of isoforms, including extracellular SOD (Ec-SOD), mitochondrial SOD (Mn-SOD), and cytosolic SOD (Cyt-SOD). They are dispersed throughout different cellular compartments to focus on particular sources of superoxide generation [27]. One enzyme that helps break down H₂O₂ into water (H₂O) and O₂ is called catalase, and it is present in peroxisomes. Particularly in areas with high metabolic activity, it acts as an essential defense against the detrimental effects of accumulating H₂O₂ [43]. Reduced glutathione (GSH) is used as a cofactor by glutathione peroxidase (GPx) enzymes to lower lipid hydroperoxides and H₂O₂. This process changes GSH into its oxidized form glutathione-disulfide (GSSG), which is then regenerable by other enzymes, such as glutathione reductase. GPx enzymes play a critical role in limiting the buildup of hazardous peroxides [44]. The thioredoxin framework, which is made up of the essential enzyme thioredoxin reductase, is responsible for preserving the redox balance of cells. It does this by facilitating the reduction of oxidized

thioredoxin, which allows it to participate in antioxidant defense mechanisms [45]. Heme oxygenase-1 (HO-1) is an essential component of cellular defense against OS, despite not being a traditional antioxidant enzyme. It breaks down heme to release iron, carbon monoxide, and biliverdin. These metabolites exhibit strong cytoprotective and antioxidant properties [46]. One important peroxidase family that is involved in neutralizing peroxides, such as H₂O₂, is called peroxiredoxin. They are extensively dispersed throughout different cellular compartments and are essential elements of the antioxidant defense mechanism [47].

4. OS AND INSULIN RESISTANCE

Insulin resistance is significantly influenced by OS. By preventing the insulin receptor substrate (IRS) from being phosphorylated, which is essential for subsequent signaling processes, it interferes with insulin signaling pathways. It also obstructs insulin-sensitive tissues' ability to absorb glucose, worsening hyperglycemia. All of these complex consequences of OS play a part in the emergence and advancement of insulin

resistance, a feature of type 2 DM (T2DM). Comprehending these nuances is essential for focused treatments in the management of DM [48]. Once insulin binds to the alpha sections, a number of substrate proteins are phosphorylated, with the IRS proteins proving to be the most significant. This results in the phosphorylation of the beta sections. Phosphorylated IRS proteins are initiated at tyrosine residues and can subsequently initiate two significant signaling cascades. Involved in both gene expression and cell proliferation, the first pathway produces mitogen-activated protein kinase (MAPK). The second is the insulin metabolic pathway, which is controlled by the phosphatidylinositol-3 kinase (PI3K) pathway. Pyruvate dehydrogenase kinase (PDK) is activated when phosphatidyl inositol 3,4,5 trisphosphate (PIP₃) is produced as a result of PI3K's phosphorylation of phosphatidyl inositol 4,5 bisphosphate (PIP₂). PDK phosphorylates and activates Protein Kinase-B (PKB; often referred to as AKT), which causes the translocation of the glucose transporter 4 (GLUT4) to the cell membrane, which results in a subsequent glucose influx [49] (Fig. 3).




Fig. 3. Insulin signaling pathways affected by OS: Tyrosine residues are the site of phosphorylation, and after that, phosphorylated IRS proteins can activate two important signaling pathways. First pathway leads to MAPK, being involved in cell growth and gene expression. The second is the PI3K pathway, that is in charge of insulin's metabolic function. OS disrupts both insulin signaling pathways by impeding tyrosine phosphorylation of the IRS, crucial for downstream signaling events. Additionally, it interferes with glucose uptake in insulin-sensitive tissues, further exacerbating hyperglycemia

4.1 Impact on Insulin Signaling Pathways

One of the main characteristics of T2DM is insulin resistance, which is characterized by a decrease in the cellular reactivity to the insulin signal. Insulin resistance is made worse by OS, which has a significant impact on the insulin signaling pathways. Insulin typically binds to its receptor and sets off a cascade of phosphorylation events that lead to the activation of signaling molecules downstream. These early phosphorylation activities are susceptible to disruption by OS, which hinders the activation of later signaling pathway components [50]. Insulin receptor signals are transmitted to downstream effectors through the critical intermediates, IRS proteins, in the insulin signaling cascade. Insulin signaling is inhibited when serine phosphorylation of IRS proteins occurs due to OS. This occurs because the IRS proteins are unable to bind to the insulin receptor. AKT, often referred to as PKB, is activated by the essential enzyme PI3K in the insulin signaling cascade. For most of the metabolic effects of insulin, AKT is an essential mediator. Reduction of AKT activation and decreased glucose absorption can result from PI3K inhibition caused by OS [51,52].

Muscle and adipose tissue in particular are insulin-sensitive cells, and GLUT4 is in charge of promoting glucose transport into these cells. The normal translocation of GLUT4 to the cell membrane is disrupted by OS, which lowers glucose absorption and increases the risk of hyperglycemia. Activation of specific PKC isoforms can result from elevated OS levels. By encouraging the phosphorylation of IRS proteins in an inhibitory manner and interfering with downstream signaling events, this can obstruct insulin signaling [53]. Chronic inflammation is a key component of insulin resistance and is closely associated with oxidative damage. Insulin signaling pathways can be directly disrupted by inflammatory cytokines and messenger molecules that are triggered by OS. Diacylglycerol (DAG), one lipid intermediary that can accumulate as a result of OS-induced mitochondrial malfunction, can activate PKC theta and disrupt insulin signaling [53,54].

4.2 Disruption of Glucose Uptake

Interfering with glucose uptake into insulin-sensitive cells is one of the main ways that OS leads to insulin resistance in DM. The two main sites of insulin-mediated glucose uptake, skeletal muscle and adipose tissue, are the primary

targets of this disturbance. The major glucose transporter in muscle and adipose tissue that facilitates insulin-stimulated glucose absorption is called GLUT4 [55]. Normally, insulin causes GLUT4 to move from intracellular vesicles to the cell membrane, which opens the door for glucose to enter the cell. This translocation pathway is hampered by OS, which lowers GLUT4's availability at the cell surface. OS disrupts essential elements of the insulin signaling cascade, as was previously mentioned. This disturbance directly impacts the proteins involved in vesicle trafficking and GLUT4 translocation, in addition to reducing AKT activation and impairing downstream effector phosphorylation [55,56].

Lipid intermediates like DAG and ceramides can build up as a result of mitochondrial malfunction brought on by OS. These lipid compounds cause insulin resistance by interfering with insulin signaling pathways. They may also cause the activation of protein kinase C isoforms, which further suppresses the effects of insulin. Chronic inflammation is linked to elevated levels of OS and is a major contributor to insulin resistance. Adipose tissue releases adipokines and inflammatory cytokines that can obstruct insulin signaling and impair the absorption of glucose [57]. ER stress, which is defined by the build-up of misfolded proteins in the ER, can be brought on by OS. GLUT4 translocation can be decreased, and insulin signaling is hampered by ER stress. The synthesis of ATP can be hindered by damage to mitochondria caused by OS. Insulin resistance may worsen as a result of the decreased energy available for the transport and metabolism of glucose. Insulin sensitivity and glucose metabolism are significantly influenced by adipose tissue. Dysfunctional adipocytes, which release adipokines and cytokines that obstruct insulin signaling and glucose absorption, can be brought on by OS [58].

5. COMPLICATIONS OF DM MEDIATED BY OS

One major factor in problems with DM is OS. It is the underlying cause of cardiomyopathy, altering the heart's structure and function. Deficits in sensation and motor function result from oxidative damage to nerve cells, which causes neuropathy. Increased OS causes renal tissue damage in nephropathy. Cerebral OS causes encephalopathy, which affects cognitive function. The oxidative damage to blood vessels that results in peripheral vascular disease impairs blood flow. In conclusion, retinopathy presents as

oxidative damage to the retina, endangering one's vision [59]. It is essential to comprehend these complexities in order to mitigate issues related to DM (Fig. 4).

5.1 Cardiomyopathy

DM significantly increases the risk of cardiovascular problems, which are mostly brought on by OS. Stress reduces the availability of NO, which prevents arterial dilatation and inhibits endothelial function. OS-induced LDL oxidation speeds up the course of atherosclerosis, which is characterized by the accumulation of arterial plaque. This process promotes inflammation, the recruitment of monocytes, and the production of foam cells. Increased vascular resistance, decreased vasodilation, and renin-angiotensin-aldosterone

system (RAAS) activation are the main causes of hypertension, which is frequent in people with DM [60,61]. The heart is prone to oxidative damage, which can result in remodelling and fibrosis, as well as impaired contractility and relaxation. The activation of platelets increases the risk of thrombosis. ROS-induced vascular inflammation attracts leukocytes, worsening the condition. Oxidative nerve damage causes autonomic neuropathy, a frequent DM consequence that impairs heart rate regulation. This increases the risk of arrhythmia and causes irregular heart rate variability. Additionally, OS reduces coronary flow reserve, which makes it more difficult for the heart to get enough blood while under stress or exercising. These results highlight the critical role that OS plays in DM-related cardiomyopathy and its associated consequences [61].




Fig. 4. Diabetic complications caused by OS: OS underlies retinopathy, causing oxidative injury to the retina, threatening vision. Cardiomyopathy manifests as structural and functional changes in the heart. Neuropathy arises from oxidative damage to nerve cells, leading to sensory and motor deficits. Nephropathy involves renal tissue injury due to increased oxidative load. Encephalopathy results from cerebral OS, impacting cognitive function. Peripheral arteriopathy arises from oxidative damage to blood vessels, compromising circulation

5.2 Neuropathy

OS plays a major role in the development of diabetic neuropathy, a condition marked by nerve damage. Particularly vulnerable to OS-induced mitochondrial dysfunction, which impairs synaptic function and energy production, are nerve cells. An accumulation of dangerous oxidative molecules brought on by overzealous antioxidant defenses directly damages cellular components and impairs neuronal function [62,63]. Furthermore, endothelial dysfunction brought on by OS exacerbates nerve injury by reducing blood flow. The development of neuropathy is significantly influenced by OS-induced inflammation, which is exacerbated by immune cells and inflammatory mediators. High blood sugar causes AGEs to develop, which worsen nerve function even further. Neural signal transmission is disrupted by OS-induced structural changes such as demyelination and axonal degeneration, leading to impairments in motor and sensory functions. Inequalities in neurotrophic factors, which are essential for the survival of nerve cells, exacerbate nerve injury. Furthermore, neuropathic pain—a prevalent sign of diabetic neuropathy—is exacerbated by OS, which alters pain perception pathways. Pain sensitization is caused by changes in neurotransmitters and increased inflammation [62].

5.3 Nephropathy

OS has a major role in diabetic nephropathy, a condition marked by kidney damage and decreased function. Under these conditions, pro-oxidant and antioxidant components in the glomeruli are out of balance, which causes oxidative damage and poor filtration. The progression of nephropathy and structural alterations are associated with activation of the fibrotic pathway and inflammatory responses [64]. Moreover, RAAS activation brought on by OS increases blood pressure and exacerbates renal impairment. The damage caused by OS impairs the function of podocytes, which are essential for filtration. Glomerular enlargement and the onset of nephropathy are caused by mesangial cell dysfunction. Damage to intracellular signaling networks results in malfunctioning cells and compromised repair systems. The AGEs produced by elevated glucose levels worsen inflammation and oxidative damage. Damaged components accumulate, and kidney damage gets worse as a result of OS's disruption of autophagy.

Nephropathy is made worse by microvascular injury, which reduces oxygen and blood supply to the kidneys. These complex processes highlight how important OS is to diabetic nephropathy [65].

5.4 Retinopathy

OS is a major factor in diabetic retinopathy, which is defined by retinal blood vessel destruction and possible visual loss. Microvascular injury and fluid leaking into surrounding tissue are caused by this stress, which impairs endothelial cell function. High blood sugar causes AGEs to accumulate in the retinal tissues, which furthers the retinopathy's progression by causing OS, inflammation, and cellular damage [66]. Retinal edema is a result of blood and fluid leaking from damaged blood-retinal barriers, which increases vascular permeability. Damage to the retina is exacerbated by OS, which triggers inflammatory reactions that draw immune cells and release cytokines that promote inflammation. It is possible for ischemia to cause neovascularization; however, the retinopathy may worsen since these new arteries are frequently brittle and can burst. Contributing to visual impairment are retinal ganglion cell dysfunction and apoptosis, which are essential for the transmission of visual signals. Because OS causes an increase in vascular endothelial development factor, this may exacerbate damage by encouraging the creation of new blood vessels. Further damage to retinal tissue results from OS's disruption of the regulation of retinal blood flow, which exacerbates ischemia-reperfusion injury. These results highlight how important OS is to the onset and course of diabetic retinopathy [67].

5.5 Encephalopathy

OS is intimately associated with encephalopathy, a disorder characterized by malfunction of the brain, especially in circumstances such as DM. The brain is more vulnerable to oxidative injury when glucose levels are high and antioxidant defenses are weakened. Because of their lipid-rich makeup and high metabolic activity, neurons are especially susceptible to the effects of ROS and RNS. This oxidative assault compromises cellular homeostasis, which results in decreased energy production, mitochondrial malfunction, and an ongoing cycle of ROS formation [68]. Furthermore, inflammation brought on by OS plays a major role in the onset and development

of encephalopathy. Neuroinflammation is sparked by the upregulation of inflammatory mediators, including chemokines and cytokines, which exacerbates damage to neurons. OS can also weaken the blood-brain barrier, which is an important protective interface, and can let potentially hazardous compounds into the brain environment, which can lead to encephalopathy. Acknowledging the complex interactions between OS and encephalopathy is essential to designing focused treatment approaches to reduce brain dysfunction in both DM and other neurological conditions with increased oxidative load [68,69].

5.6 Peripheral arteriopathy

OS and peripheral arteriopathy are closely related, especially in the case of lower limb wounds. Long-term high glucose levels and weakened antioxidant defenses in diseases like DM foster an atmosphere that encourages oxidative damage to the peripheral vasculature. ROS and RNS have a particularly negative effect on endothelial cells, which are essential for vascular function. This oxidative attack compromises vasodilation, creates atherosclerotic plaque, and interferes with endothelial function [70]. Furthermore, inflammation brought on by OS plays a major role in the advancement of peripheral arteriopathy. Inflammatory mediators worsen endothelial dysfunction and encourage a milieu that is pro-inflammatory and favours atherogenesis. Additionally, soft muscle tissues in the vessel walls are directly impacted by OS, which promotes their growth and narrows the vessels. Furthermore, LPO brought on by OS intensifies the development of atherosclerotic plaque. Understanding the critical role that OS plays in the aetiology of peripheral arteriopathy is essential for creating focused treatment plans to reduce vascular consequences, particularly in patients with diseases like DM that have an elevated oxidative load [71].

6. ANTIOXIDANT DEFENSE SYSTEM

Protection against oxidative damage is provided by antioxidant defense systems in cells. Protecting the body from damaging ROS are endogenous antioxidants, such as SOD. Fruits and vegetables, which are rich in dietary

antioxidants, help to protect the body from OS by restocking the antioxidant reservoir in the body [2] (Table 1).

6.1 Endogenous antioxidants

The body spontaneously produces endogenous antioxidants, which are vital for preserving redox balance and counteracting damaging ROS. The defense of tissues and cells against oxidative damage depends on these antioxidants [72]. Superoxide radicals are transformed into less dangerous molecules like H_2O_2 and O_2 by the catalyst action of SOD, a major endogenous antioxidant enzyme. SOD is found in multiple forms in different parts of the cell. These forms include extracellular SOD (Ec-SOD), mitochondrial SOD (Mn-SOD), and cytosolic SOD (Cyt-SOD) [26]. The essential enzyme catalase also functions in cellular peroxisomes. It helps to break down H_2O_2 into O_2 and H_2O , avoiding the buildup of this potentially dangerous molecule [43]. The GPx enzymes are a class of selenium-dependent enzymes that decrease H_2O_2 and lipid hydroperoxides by using GSH as a cofactor. This enzymatic process shields cells from oxidative harm [44]. This enzyme is necessary for the recycling and regeneration of GSH from GSSG, which is the oxidized form of this amino acid. This recycling mechanism helps glutathione keep on acting as an effective cellular antioxidant [73]. Thioredoxin, thioredoxin reductase, and NADPH make up the thioredoxin system. It is essential for preserving the redox equilibrium within cells. By catalyzing the reduction of oxidized thioredoxin, thioredoxin reductase enables thioredoxin to take part in antioxidant defense mechanisms [74]. These proteins play a role in controlling the amount of iron that cells contain. They aid in preventing the production of hazardous ROS via Fenton reactions, which might end up in oxidative damage, by sequestering iron [75]. Heme, which is created when red blood cells break down, is broken down into bilirubin. It functions as a strong natural antioxidant that has the ability to counteract free radicals [76]. The natural result of purine metabolism is uric acid. Uric acid possesses antioxidant qualities and can aid in the neutralization of specific ROS forms, regardless of its link to gout [77].

Table 1. Types of antioxidants in defense system: (a) endogenous antioxidants which are produced within the body and neutralize harmful ROS, and (b) dietary antioxidants which bolster its ability to counteract OS [89]

Types of antioxidants in defense system	
Endogenous Antioxidants	Dietary Antioxidants
Superoxide dismutase	Vitamin C & E
Catalase	Zinc
Glutathione peroxidase	Beta-carotene
Glutathione reductase	Selenium
Thioredoxin system	Flavonoids
Ferritin and heme oxygenase	Polyphenols
Bilirubin	Lycopene
Uric acid	Coenzyme Q10

6.2 Dietary Antioxidants

Dietary antioxidants are substances present in a wide range of foods that, by scavenging dangerous ROS and free radicals, can aid in shielding cells from oxidative damage. A person's general health and well-being can be enhanced by consuming a range of foods high in these antioxidants [78]. As a strong water-soluble antioxidant, vitamin C (ascorbic acid) can be found in citrus fruits (lemons, oranges), berries (strawberries, kiwi), leafy greens (kale, spinach), and bell peppers. It replenishes vitamin E, scavenges free radicals, and boosts immunity [79]. Rich sources of vitamin E (tocopherols and tocotrienols) include nuts (almonds, hazelnuts), seeds (pumpkin, sunflower), spinach, and vegetable oils (safflower, sunflower). This lipid-soluble antioxidant shields cell membranes from LPO [80]. Generally, orange fruits and vegetables, including carrots, sweet potatoes, and mangoes, are rich in beta-carotene, a precursor of vitamin A. It is an effective antioxidant that supports cell damage prevention [81]. Whole grains, chicken, fish (tuna, halibut), Brazil nuts, and fowl are good sources of selenium, a trace mineral. It is an important part of selenoproteins (SEPs), which are necessary for antioxidant defense and include glutathione peroxidases [82]. Plant chemicals known as flavonoids are a varied class that feature antioxidant capabilities. Foods such as citrus fruits, onions, apples, tea, and berries (strawberries, blueberries) contain flavonoids. There are several health advantages, and flavonoids assist in scavenging free radicals [83]. Foods high in polyphenols include red wine, berries, grapes, dark chocolate, and green tea. They support general health since they have strong anti-inflammatory and antioxidant effects [84]. Lycopene, a red pigment, is abundant in

watermelon, pink grapefruit, and tomatoes. It is a potent antioxidant that is well-known for helping to lower the risk of a number of chronic ailments, including heart disease and several types of cancer [85]. Fish, whole grains, and organ meats are good sources of coenzyme Q10 (ubiquinone). It is an essential part of the mitochondria's electron transport chain and functions as an antioxidant to shield cells from oxidative damage [86]. Meat, shellfish, nuts, and seeds are among the foods that contain zinc, a trace mineral. It is a crucial part of the enzymes that defend against free radicals and repair DNA [87]. Anthocyanins are pigments that can be found in purple, blue, and red fruits and vegetables, such as grapes, eggplants, and berries. They give these meals their vivid colours and have potent antioxidant qualities [88].

7. THERAPEUTIC INTERVENTIONS

Numerous treatment options, including antioxidant supplements, lifestyle adjustments, and pharmaceutical therapies, have been investigated in the quest to mitigate OS in DM.

7.1 Antioxidant Supplementation

In order to strengthen the body's natural defenses against OS, antioxidant supplementation refers to the use of exogenous antioxidants through food, supplements, or pharmaceuticals. Coenzyme Q10, beta-carotene, selenium, vitamins C and E, and other nutrients are common antioxidant supplements. These supplements come in a number of formats, including liquid extracts, pills, and capsules. Supplementing with antioxidants may help lessen oxidative damage and lower the chance of developing some chronic illnesses linked to high OS levels [90]. Conditions including age-related

macular degeneration, some malignancies, and cardiovascular disease fall under this category. Researchers are still studying the efficacy of antioxidant supplements. While several studies point to possible dangers or no discernible effects at all in some populations, others indicate potential advantages. It's crucial to speak with a healthcare professional before beginning any antioxidant supplementation program because taking too much of some antioxidants can have negative consequences [91].

7.2 Lifestyle Modifications

The management of OS and the mitigation of related problems are greatly aided by changes in lifestyle. Antioxidants and vital nutrients can be obtained by eating a balanced diet full of whole grains, fruits, vegetables, and lean meats. It also promotes general health by reducing OS. Active living can help lower OS and improve general well-being. Its benefits include improved cardiovascular health and strengthened antioxidant defense mechanisms. Both OS and chronic stress are related [92]. Its effects can be lessened by putting stress-reduction strategies like deep breathing exercises, mindfulness training, and meditation into practice. The two main causes of OS are smoking and binge drinking. The amount of OS that the body experiences can be greatly decreased by giving up smoking and consuming alcohol in moderation. The body's natural healing processes, which include those involving OS, depend on getting enough good sleep. To promote general health, try to get between seven and nine hours of sleep each night [92,93].

7.3 Pharmaceutical Approaches

Medications or other treatments created specially to address OS and its related problems are used in pharmaceutical techniques. Certain pharmaceuticals either increase the body's natural antioxidant defenses or function as direct antioxidants. These include alpha-lipoic acid, N-acetylcysteine (NAC), and specific metal chelators. OS and inflammation are strongly connected. Medications that reduce inflammation, such as anti-inflammatory medicines or nonsteroidal anti-inflammatory drugs (NSAIDs), may be able to lessen the effects of OS. It is necessary to use pharmaceutical therapies designed to target the underlying pathology, depending on the particular OS-related consequence (such as cardiovascular disease or neurodegenerative

illnesses). Pharmaceutical strategies must be customized based on each patient's unique medical history, risk factors, and treatment response. Finding the right pharmaceutical interventions requires consulting with a healthcare professional [94].

8. EMERGING RESEARCH AND FUTURE DIRECTIONS

To better understand the intricate connection between OS and DM and to develop targeted, individualized approaches for managing and treating OS-related complications linked to the disease, researchers are focusing on the following areas:

8.1 Novel Therapies Targeting OS

Novel treatment approaches are being intensively investigated in the quickly developing field of OS research. This involves the investigation of novel antioxidant chemicals that exhibit enhanced bioavailability and specificity in addressing particular forms of ROS, or OS, in discrete cellular divisions. Additionally, efforts are being directed toward creating treatments that target OS and mitochondrial malfunction directly, with possible drugs that protect and improve mitochondrial DNA [95]. To further boost the therapeutic efficiency of antioxidants, research is being done on how well nanoparticles and nanocarriers can carry antioxidants to specific cellular compartments or tissues, enhance their bioavailability, and stabilize them. Promising advances in gene therapy and editing technologies have the ability to directly modify genes implicated in antioxidant defense systems, which could raise endogenous antioxidant enzyme levels or change pathways connected to OS [96,97]. Furthermore, studies explore bioactive substances derived from plants and other natural sources that possess strong antioxidant capabilities, offering possible treatments for disorders associated with OS. Research on metalloenzymes and redox-active metals aims to regulate OS by means of their therapeutic potential; they may function as modulators of redox signaling pathways or as catalysts for antioxidant processes. Optimizing therapy outcomes may involve customizing interventions based on unique genetic and environmental characteristics. The possible synergistic benefits of combining various antioxidants or antioxidant-based therapies with conventional treatments are being investigated.

Developments in biomarker research allow for accurate, personalized evaluation of OS levels, which helps select the right therapies and enable real-time therapy monitoring. Clinical trials assessing new treatments aimed at OS show promise for being incorporated into clinical practice, highlighting the ever-changing field of OS management research and its potential to completely transform the treatment of related illnesses [95,96].

8.2 Personalized Medicine in DM Management

Precision medicine, also known as personalized medicine, transforms healthcare by considering individual differences in illness susceptibility, progression, and response to therapy. It orients therapies for DM control to the unique features of each patient. Recent advances in genomic and molecular research have identified genetic variants and molecular markers associated with DM risks, subtypes, and treatment outcomes. This information is used in personalized medicine to classify people according to their molecular and genetic profiles [98,99]. Employing genetic and medical information, it determines a person's risk of developing DM. This allows for focused prevention through dietary changes, medication, and close monitoring of individuals who are more vulnerable. DM involves a variety of pathophysiological pathways, which is why customized therapy divides patients into several categories. This method customizes therapy regimens by taking into consideration lifestyle factors, comorbidities, metabolic profiles, and genetic markers. It simplifies the process of choosing a course of therapy for the highest level of tolerance and effectiveness by predicting a person's reaction to particular drugs based on their genetic and metabolic composition [99,100]. Exercise prescriptions and individualized nutrition regimens maximize the suggested intake of food and exercise, respectively. Digital health technologies offer real-time data for dynamic therapy modifications, such as wearables and smartphone apps. Early intervention options are made possible by considering a person's lifetime risk of complications associated with DM. Careful attention is required to ensure informed consent, data security, and ethical adherence in light of ethical and privacy considerations. Personalized medicine promises to improve patient satisfaction, results, and the burden of complications when it is used in DM care, which is a major improvement in clinical practice [100,101].

9. CONCLUSION

DM is often associated with OS, which is defined as an imbalance between the body's antioxidant defense systems and the generation of ROS. Its prevalence is not only associated with hyperglycemia but also affects cellular processes, insulin signaling, and the functioning of key organs through a variety of mechanisms. This imbalance is the starting point for a series of events that eventually result in inflammation, cellular damage, and the emergence of problems like retinopathy, neuropathy, nephropathy, encephalopathy, and cardiomyopathy. The field of DM and OS is constantly developing, and early management and treatment are becoming more and more essential. This entails developing novel treatments that deal with the underlying causes of OS in addition to the specialized utilization of antioxidants and changes in lifestyle. In conclusion, OS and DM are related through a dynamic and intricate interaction involving genetic, molecular, and environmental variables. An understanding of this link lays the groundwork for the creation of focused therapies that could revolutionize the way DM is managed. With research continuing to advance and customized treatments being more widely incorporated into clinical practice, there is hope that the impact of DM-related complications will be significantly mitigated in the future, ultimately improving the quality of life for millions of people worldwide.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Azzi A. Oxidative stress: What is it? Can it be measured? Where is it located? Can it be good or bad? Can it be prevented? Can it be cured? Antioxidants. 2022;11:1431. DOI: 10.3390/antiox11081431
2. Sharifi-Rad M, Anil Kumar NV, Zucca P, et al. Lifestyle, oxidative stress, and antioxidants: Back and forth in the

- pathophysiology of chronic diseases. *Front Physiol.* 2020;11:694.
DOI: 10.3389/fphys.2020.00694
3. González P, Lozano P, Ros G, et al. Hyperglycemia and oxidative stress: An Integral, updated and critical overview of their metabolic interconnections. *Int J Mol Sci.* 2023;24:9352.
DOI: 10.3390/ijms24119352
 4. Bhatti JS, Sehrawat A, Mishra J, et al. Oxidative stress in the pathophysiology of type 2 diabetes and related complications: Current therapeutics strategies and future perspectives. *Free Radic Biol Med.* 2022; 184:114–34.
DOI: 10.1016/j.freeradbiomed.2022.03.019
 5. Masenga SK, Kabwe LS, Chakulya M, et al. Mechanisms of oxidative stress in metabolic syndrome. *Int J Mol Sci.* 2023;24:7898.
DOI: 10.3390/ijms24097898
 6. Caturano A, D'Angelo M, Mormone A, et al. Oxidative stress in type 2 diabetes: Impacts from pathogenesis to lifestyle modifications. *Curr Issues Mol Biol.* 2023;45:6651–66.
DOI: 10.3390/cimb45080420
 7. Ly LD, Xu S, Choi S-K, et al. Oxidative stress and calcium dysregulation by palmitate in type 2 diabetes. *Exp Mol Med.* 2017;49:e291–e291.
DOI: 10.1038/emm.2016.157
 8. Rehman K, Akash MSH. Mechanism of generation of oxidative stress and pathophysiology of type 2 diabetes mellitus: How Are they interlinked? *J Cell Biochem.* 2017;118:3577–85.
DOI: 10.1002/jcb.26097
 9. Chianese R, Pierantoni R. Mitochondrial Reactive Oxygen Species (ROS) production alters sperm quality. *Antioxidants.* 2021;10:92.
DOI: 10.3390/antiox10010092
 10. Twarda-Clapa A, Olczak A, Białkowska AM, et al. Advanced Glycation End-Products (AGEs): Formation, chemistry, classification, receptors, and diseases related to AGEs. *Cells.* 2022;11: 1312.
DOI: 10.3390/cells11081312
 11. Njeim R, Alkhansa S, Fornoni A. Unraveling the crosstalk between lipids and NADPH oxidases in diabetic kidney disease. *Pharmaceutics.* 2023;15: 1360.
DOI: 10.3390/pharmaceutics15051360
 12. Aldosari DI, Malik A, Alhomida AS, et al. Implications of Diabetes-induced altered metabolites on retinal neurodegeneration. *Front Neurosci.* 2022;16.
DOI: 10.3389/fnins.2022.938029
DOI: 10.3389/fnins.2022.938029
 13. Black HS. A synopsis of the associations of oxidative stress, ROS, and antioxidants with diabetes mellitus. *Antioxidants.* 2022;11:2003.
DOI: 10.3390/antiox11102003
 14. Abdal Dayem A, Hossain M, Lee S, et al. The role of reactive oxygen species (ROS) in the biological activities of metallic nanoparticles. *Int J Mol Sci.* 2017;18:120.
DOI: 10.3390/ijms18010120
 15. Kang Q, Yang C. Oxidative stress and diabetic retinopathy: Molecular mechanisms, pathogenetic role and therapeutic implications. *Redox Biol.* 2020; 37:101799.
DOI: 10.1016/j.redox.2020.101799
 16. Asmat U, Abad K, Ismail K. Diabetes mellitus and oxidative stress—A concise review. *Saudi Pharm J.* 2016;24:547–53.
DOI: 10.1016/j.jsps.2015.03.013
 17. Su L-J, Zhang J-H, Gomez H, et al. Reactive oxygen species-induced lipid peroxidation in apoptosis, autophagy, and ferroptosis. *Oxid Med Cell Longev.* 2019;2019:1–13.
DOI: 10.1155/2019/5080843
 18. Andrés CMC, Pérez de la Lastra JM, Andrés Juan C, et al. reactive species on amino acids—biological relevance in proteins and induced pathologies. *Int J Mol Sci.* 2022;23:14049.
DOI: 10.3390/ijms232214049
 19. Juan CA, Pérez de la Lastra JM, Plou FJ, et al. The chemistry of reactive oxygen species (ros) revisited: Outlining their role in biological macromolecules (DNA, Lipids and Proteins) and Induced Pathologies. *Int J Mol Sci.* 2021;22:4642.
DOI: 10.3390/ijms22094642
 20. Rowe LA, Degtyareva N, Doetsch PW. DNA damage-induced reactive oxygen species (ROS) stress response in *Saccharomyces cerevisiae*. *Free Radic Biol Med.* 2008;45:1167–77.
DOI: 10.1016/j.freeradbiomed.2008.07.018
 21. Kowalczyk P, Sulejczak D, Kleczkowska P, et al. Mitochondrial Oxidative Stress—A causative factor and therapeutic target in many diseases. *Int J Mol Sci.* 2021;22: 13384.
DOI: 10.3390/ijms222413384
 22. Amen OM, Sarker SD, Ghildyal R, et al.

- Endoplasmic reticulum stress activates unfolded protein response signaling and mediates inflammation, obesity, and cardiac dysfunction: Therapeutic and molecular approach. *Front Pharmacol.* 2019;10:977.
DOI: 10.3389/fphar.2019.00977
23. Checa J, Aran JM. Reactive Oxygen species: Drivers of physiological and pathological processes. *J Inflamm Res.* 2020;13:1057–73.
DOI: 10.2147/JIR.S275595
 24. Miller MA, Zachary JF. Mechanisms and morphology of cellular injury, adaptation, and death. *Pathologic Basis of Veterinary Disease.* Elsevier. 2017;2-43.e19.
Available:<https://doi.org/10.1016/B978-0-323-35775-3.00001-1>
DOI: 10.1016/B978-0-323-35775-3.00001-1
 25. Goycheva P, Petkova-Parlapanska K, Georgieva E, et al. Biomarkers of oxidative stress in diabetes mellitus with diabetic nephropathy complications. *Int J Mol Sci.* 2023;24:13541.
DOI: 10.3390/ijms241713541
 26. Fujii J, Homma T, Osaki T. Superoxide radicals in the execution of cell death. *Antioxidants.* 2022;11:501.
DOI: 10.3390/antiox11030501
 27. Wang Y, Branicky R, Noë A, et al. Superoxide dismutases: Dual roles in controlling ROS damage and regulating ROS signaling. *J Cell Biol.* 2018;217:1915–28.
DOI: 10.1083/jcb.201708007
 28. Zhao Z. Iron and oxidizing species in oxidative stress and Alzheimer's disease. *AGING Med.* 2019;2:82–7.
DOI: 10.1002/agm2.12074
 29. Forstermann U, Sessa WC. Nitric oxide synthases: Regulation and function. *Eur Heart J.* 2012;33:829–37.
DOI: 10.1093/eurheartj/ehr304
 30. Andrés CMC, Pérez de la Lastra JM, Andrés Juan C, et al. Superoxide anion chemistry—Its role at the core of the innate immunity. *Int J Mol Sci.* 2023;24:1841.
DOI: 10.3390/ijms24031841
 31. Tarafdardar A, Pula G. The role of NADPH Oxidases and oxidative stress in neurodegenerative disorders. *Int J Mol Sci.* 2018;19:3824.
DOI: 10.3390/ijms19123824
 32. Tenopoulou M, Doulias P-T. Endothelial nitric oxide synthase-derived nitric oxide in the regulation of metabolism. *F1000Research.* 2020;9:1190.
DOI: 10.12688/f1000research.19998.1
 33. Król M, Kepinska M. Human nitric oxide synthase—Its functions, polymorphisms, and inhibitors in the context of inflammation, diabetes and cardiovascular diseases. *Int J Mol Sci.* 2020;22:56.
DOI: 10.3390/ijms22010056
 34. Zorov DB, Juhaszova M, Sollott SJ. Mitochondrial Reactive Oxygen Species (ROS) and ROS-Induced ROS Release. *Physiol Rev.* 2014;94:909–50.
DOI: 10.1152/physrev.00026.2013
 35. Mittal M, Siddiqui MR, Tran K, et al. Reactive oxygen species in inflammation and tissue injury. *Antioxid Redox Signal.* 2014;20:1126–67.
DOI: 10.1089/ars.2012.5149
 36. Ayala A, Muñoz MF, Argüelles S. Lipid Peroxidation: Production, metabolism, and signaling mechanisms of malondialdehyde and 4-Hydroxy-2-Nonenal. *Oxid Med Cell Longev.* 2014;2014:1–31.
DOI: 10.1155/2014/360438
 37. Cherian D, Peter T, Narayanan A, et al. Malondialdehyde as a marker of oxidative stress in periodontitis patients. *J Pharm Bioallied Sci.* 2019;11:297.
DOI: 10.4103/JPBS.JPBS_17_19
 38. Li Y, Zhao T, Li J, et al. Oxidative Stress and 4-hydroxy-2-nonenal (4-HNE): Implications in the pathogenesis and treatment of aging-related diseases. *J Immunol Res.* 2022;2022:1–12.
DOI: 10.1155/2022/2233906
 39. Milne GL, Yin H, Hardy KD, et al. Isoprostane generation and function. *Chem Rev.* 2011;111:5973–96.
DOI: 10.1021/cr200160h
 40. Chang X, Wang Y, Zheng B, et al. The Role of acrolein in neurodegenerative diseases and its protective strategy. *Foods.* 2022;11:3203.
DOI: 10.3390/foods11203203
 41. Barrera G. Oxidative stress and lipid peroxidation products in cancer progression and therapy. *ISRN Oncol.* 2012;2012:1–21.
DOI: 10.5402/2012/137289
 42. De Freitas FA, Levy D, Zarrouk A, et al. Impact of oxysterols on cell death, proliferation, and differentiation induction: current status. *Cells.* 2021;10:2301.
DOI: 10.3390/cells10092301
 43. Nandi A, Yan L-J, Jana CK, et al. Role of catalase in oxidative stress- and age-associated degenerative diseases. *Oxid*

- Med Cell Longev. 2019;2019:1–19.
DOI: 10.1155/2019/9613090
44. Lubos E, Loscalzo J, Handy DE. Glutathione peroxidase-1 in health and disease: from molecular mechanisms to therapeutic opportunities. *Antioxid Redox Signal*. 2011;15:1957–97.
DOI: 10.1089/ars.2010.3586
 45. Lee S, Kim SM, Lee RT. Thioredoxin and thioredoxin target proteins: from molecular mechanisms to functional significance. *Antioxid Redox Signal*. 2013;18:1165–207.
DOI: 10.1089/ars.2011.4322
 46. Puentes-Pardo JD, Moreno-SanJuan S, Carazo Á, et al. Heme oxygenase-1 in gastrointestinal tract health and disease. *Antioxidants*. 2020;9:1214.
DOI: 10.3390/antiox9121214
 47. Karpenko IL, Valuev-Elliston VT, Ivanova ON, et al. Peroxiredoxins—The underrated actors during virus-induced oxidative stress. *Antioxidants*. 2021;10:977.
DOI: 10.3390/antiox10060977
 48. Silva Rosa SC, Nayak N, Caymo AM, et al. Mechanisms of muscle insulin resistance and the cross-talk with liver and adipose tissue. *Physiol Rep*. 2020;8:e14607.
DOI: 10.14814/phy2.14607
 49. Boucher J, Kleinridders A, Kahn CR. Insulin Receptor Signaling in Normal and Insulin-Resistant States. *Cold Spring Harb Perspect Biol*. 2014;6:a009191–a009191.
DOI: 10.1101/cshperspect.a009191
 50. Petersen MC, Shulman GI. Mechanisms of insulin action and insulin resistance. *Physiol Rev*. 2018;98:2133–223.
DOI: 10.1152/physrev.00063.2017
 51. White MF. The insulin signalling system and the IRS proteins. *Diabetologia*. 1997;40:S2–17.
DOI: 10.1007/s001250051387
 52. White MF, Kahn CR. Insulin action at a molecular level – 100 years of progress. *Mol Metab*. 2021;52:101304.
DOI: 10.1016/j.molmet.2021.101304
 53. Chadt A, Al-Hasani H. Glucose transporters in adipose tissue, liver, and skeletal muscle in metabolic health and disease. *Pflügers Arch - Eur J Physiol*. 2020;472:1273–98.
DOI: 10.1007/s00424-020-02417-x
 54. Fazakerley DJ, Krycer JR, Kearney AL, et al. Muscle and adipose tissue insulin resistance: malady without mechanism? *J Lipid Res*. 2019;60:1720–32.
DOI: 10.1194/jlr.R087510
 55. Tangvarasittichai S. Oxidative stress, insulin resistance, dyslipidemia and type 2 diabetes mellitus. *World J Diabetes*. 2015;6:456.
DOI: 10.4239/wjd.v6.i3.456
 56. Rains JL, Jain SK. Oxidative stress, insulin signaling, and diabetes. *Free Radic Biol Med*. 2011;50:567–75.
DOI: 10.1016/j.freeradbiomed.2010.12.006
 57. Park SS, Seo Y-K. Excess accumulation of lipid impairs insulin sensitivity in skeletal muscle. *Int J Mol Sci*. 2020;21:1949.
DOI: 10.3390/ijms21061949
 58. Cao SS, Kaufman RJ. Endoplasmic reticulum stress and oxidative stress in cell fate decision and human disease. *Antioxid Redox Signal*. 2014;21:396–413.
DOI: 10.1089/ars.2014.5851
 59. Oyenihi AB, Ayeleso AO, Mukwevho E, et al. Antioxidant strategies in the management of diabetic neuropathy. *Biomed Res Int*. 2015;2015:1–15.
DOI: 10.1155/2015/515042
 60. Kayama Y, Raaz U, Jagger A, et al. Diabetic cardiovascular disease induced by oxidative stress. *Int J Mol Sci*. 2015;16:25234–63.
DOI: 10.3390/ijms161025234
 61. Petrie JR, Guzik TJ, Touyz RM. Diabetes, hypertension, and cardiovascular disease: clinical insights and vascular mechanisms. *Can J Cardiol*. 2018;34:575–84.
DOI: 10.1016/j.cjca.2017.12.005
 62. Sytze van Dam P. Oxidative stress and diabetic neuropathy: pathophysiological mechanisms and treatment perspectives. *Diabetes Metab Res Rev*. 2002;18:176–84.
DOI: 10.1002/dmrr.287
 63. Pang L, Lian X, Liu H, et al. Understanding diabetic neuropathy: Focus on oxidative stress. *Oxid Med Cell Longev*. 2020;2020:1–13.
DOI: 10.1155/2020/9524635
 64. Jha JC, Banal C, Chow BSM, et al. Diabetes and kidney disease: role of oxidative stress. *Antioxid Redox Signal*. 2016;25:657–84.
DOI: 10.1089/ars.2016.6664
 65. Darenskaya M, Kolesnikov S, Semenova N, et al. Diabetic nephropathy: Significance of determining oxidative stress and opportunities for antioxidant therapies. *Int J Mol Sci*. 2023;24:12378.
DOI: 10.3390/ijms241512378
 66. Kowluru RA, Chan P-S. Oxidative stress and diabetic retinopathy. *Exp Diabetes*

- Res. 2007;2007:1–12.
DOI: 10.1155/2007/43603
67. Zhou J, Chen B. Retinal cell damage in diabetic retinopathy. *Cells*. 2023;12:1342. DOI: 10.3390/cells12091342.
68. Jelinek M, Jurajda M, Duris K. Oxidative Stress in the Brain: Basic concepts and treatment strategies in stroke. *Antioxidants*. 2021;10:1886. DOI: 10.3390/antiox10121886
69. Muriach M, Flores-Bellver M, Romero FJ, et al. Diabetes and the brain: Oxidative stress, inflammation, and autophagy. *Oxid Med Cell Longev*. 2014;2014:1–9. DOI: 10.1155/2014/102158
70. Signorelli SS, Scuto S, Marino E, et al. oxidative stress in peripheral arterial disease (pad) mechanism and biomarkers. *Antioxidants*. 2019;8:367. DOI: 10.3390/antiox8090367
71. Soyoye DO, Abiodun OO, Ikem RT, et al. Diabetes and peripheral artery disease: A review. *World J Diabetes*. 2021;12:827–38. DOI: 10.4239/wjd.v12.i6.827
72. Korczowska-Łącka I, Słowikowski B, Piekut T, et al. Disorders of endogenous and exogenous antioxidants in neurological diseases. *Antioxidants*. 2023;12:1811. DOI: 10.3390/antiox12101811
73. Averill-Bates DA. The antioxidant glutathione. *Vitam Horm*. 2023;121:109–41. DOI: 10.1016/bs.vh.2022.09.002
74. Hasan AA, Kalinina E, Tatarskiy V, et al. The Thioredoxin system of mammalian cells and its modulators. *Biomedicines*. 2022;10:1757. DOI: 10.3390/biomedicines10071757
75. ORINO K, LEHMAN L, TSUJI Y, et al. Ferritin and the response to oxidative stress. *Biochem J*. 2001;357:241. DOI: 10.1042/0264-6021:3570241
76. Ryter SW. Significance of heme and heme degradation in the pathogenesis of acute lung and inflammatory disorders. *Int J Mol Sci*. 2021;22:5509. DOI: 10.3390/ijms22115509
77. Kushiyama A, Nakatsu Y, Matsunaga Y, et al. Role of uric acid metabolism-related inflammation in the pathogenesis of metabolic syndrome components such as atherosclerosis and nonalcoholic steatohepatitis. *Mediators Inflamm*. 2016; 1–15. DOI: 10.1155/2016/8603164
78. Ponnampalam EN, Kiani A, Santhiravel S, et al. The importance of dietary antioxidants on oxidative stress, meat and milk production, and their preservative aspects in farm animals: antioxidant action, animal health, and product quality—invited review. *Animals*. 2022;12:3279. DOI: 10.3390/ani12233279
79. Susa F, Pisano R. Advances in ascorbic acid (Vitamin C) manufacturing: Green extraction techniques from natural sources. *Processes*. 2023;11:3167. DOI: 10.3390/pr11113167
80. Michalak M, Pierzak M, Kręcisz B, et al. Bioactive compounds for skin health: A review. *Nutrients*. 2021;13:203. DOI: 10.3390/nu13010203
81. Elvira-Torales LI, García-Alonso J, Periago-Castón MJ. Nutritional importance of carotenoids and their effect on liver health: A review. *Antioxidants*. 2019;8:229. DOI: 10.3390/antiox8070229
82. Tinggi U. Selenium: its role as antioxidant in human health. *Environ Health Prev Med*. 2008;13:102–8. DOI: 10.1007/s12199-007-0019-4
83. Khan J, Deb PK, Priya S, et al. Dietary flavonoids: Cardioprotective potential with antioxidant effects and their pharmacokinetic, toxicological and therapeutic concerns. *Molecules*. 2021;26:4021. DOI: 10.3390/molecules26134021
84. Rudrapal M, Khairnar SJ, Khan J, et al. Dietary polyphenols and their role in oxidative stress-induced human diseases: Insights into protective effects, antioxidant potentials and mechanism(s) of action. *Front Pharmacol*. 2022;13: 806470. DOI: 10.3389/fphar.2022.806470
85. Imran M, Ghorat F, Ul-Haq I, et al. Lycopene as a Natural antioxidant used to prevent human health disorders. *Antioxidants*. 2020;9:706. DOI: 10.3390/antiox9080706
86. Pallotti F, Bergamini C, Lamperti C, et al. The roles of coenzyme q in disease: Direct and indirect involvement in cellular functions. *Int J Mol Sci*. 2021;23:128. DOI: 10.3390/ijms23010128
87. Souyoul SA, Saussy KP, Lupo MP. Nutraceuticals: A review. *Dermatol Ther (Heidelb)*. 2018;8:5–16. DOI: 10.1007/s13555-018-0221-x
88. Gonçalves AC, Nunes AR, Falcão A, et al. Dietary effects of anthocyanins in human health: A comprehensive review.

- Pharmaceuticals. 2021;14:690.
DOI: 10.3390/ph14070690
89. Janciauskiene S. The beneficial effects of antioxidants in health and diseases. *Chronic Obstr Pulm Dis J COPD Found.* 2020;7:182–202.
DOI: 10.15326/jcopdf.7.3.2019.0152
90. Bouayed J, Bohn T. Exogenous antioxidants—double-edged swords in cellular redox state: health beneficial effects at physiologic doses versus deleterious effects at high doses. *Oxid Med Cell Longev.* 2010;3:228–37.
DOI: 10.4161/oxim.3.4.12858
91. Meulmeester FL, Luo J, Martens LG, *et al.* Antioxidant supplementation in oxidative stress-related diseases: What have we learned from studies on alpha-tocopherol? *Antioxidants.* 2022;11:2322.
DOI: 10.3390/antiox11122322
92. Ghodeswar GK, Dube A, Khobragade D. Impact of lifestyle modifications on cardiovascular health: A narrative review. *Cureus.* 2023;15:e42616.
DOI: 10.7759/cureus.42616
93. Leonidis A, Korozi M, Sykianaki E, *et al.* Improving stress management and sleep hygiene in intelligent homes. *Sensors.* 2021;21:2398.
DOI: 10.3390/s21072398
94. Tyuryaeva I, Lyublinskaya O. Expected and unexpected effects of pharmacological antioxidants. *Int J Mol Sci.* 2023;24:9303.
DOI: 10.3390/ijms24119303
95. Vassalle C, Maltinti M, Sabatino L. Targeting oxidative stress for disease prevention and therapy: Where do we stand, and where do we go from here. *Molecules.* 2020;25:2653.
DOI: 10.3390/molecules25112653
96. Curpan AS, Luca A-C, Ciobica A. Potential novel therapies for neurodevelopmental diseases targeting oxidative stress. *Oxid Med Cell Longev.* 2021;2021:1–13.
DOI: 10.1155/2021/6640206
97. Forman HJ, Zhang H. Targeting oxidative stress in disease: Promise and limitations of antioxidant therapy. *Nat Rev Drug Discov.* 2021;20: 689–709.
DOI: 10.1038/s41573-021-00233-1
98. Scheen AJ. Precision medicine: The future in diabetes care? *Diabetes Res Clin Pract.* 2016;117:12–21.
DOI: 10.1016/j.diabres.2016.04.033
99. Williams DM, Jones H, Stephens JW. Personalized Type 2 Diabetes Management: An Update on Recent Advances and Recommendations. *Diabetes, Metab Syndr Obes Targets Ther.* 2022;15:281–95.
DOI: 10.2147/DMSO.S331654.
100. Darenskaya MA, Kolesnikova LI, Kolesnikov SI. Oxidative stress: Pathogenetic role in diabetes mellitus and its complications and therapeutic approaches to correction. *Bull Exp Biol Med.* 2021;171:179–89.
DOI: 10.1007/s10517-021-05191-7
101. Van den Brink WJ, van den Broek TJ, Palmisano S, *et al.* Digital biomarkers for personalized nutrition: Predicting meal moments and interstitial glucose with non-invasive, wearable technologies. *Nutrients.* 2022; 14:4465.
DOI: 10.3390/nu14214465

© 2024 Rais et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/112366>