



# **Phytochemical Screening and Evaluation of the Larvicidal Activities of Three Organic Extracts Derived from the Leaves of *Cassia sieberiana* to Control *Anopheles gambiae*, the Vector of Malaria**

**Mamadou Kébé <sup>a</sup>, El Hadji Gorgui Diouf <sup>a\*</sup>,  
Mamadou Latyr Ndour <sup>a</sup>, Adama Faye <sup>a</sup>, Abdou Sarr <sup>b</sup>,  
Talibouya Ndior <sup>a</sup>, Ndeye Bineta Camara <sup>a</sup>,  
Mamadou Ngom <sup>a</sup>, Baboucar Sarr <sup>a</sup> and Aboubacry Sène <sup>a</sup>**

<sup>a</sup> *Laboratory of Natural Products, Department of Chemistry, Faculty of Science and Technology, Cheikh Anta Diop University of Dakar, Senegal.*

<sup>b</sup> *Department of Pharmacy, Pharmacognosy and Botanic Laboratory, UCAD, Senegal.*

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/CSJI/2024/v33i11884

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/112651>

**Original Research Article**

**Received: 27/11/2023**

**Accepted: 03/02/2024**

**Published: 06/02/2024**

\*Corresponding author: E-mail: elhadjigorgui1.diouf@ucad.edu.sn;

## ABSTRACT

Senegal is one of the African countries where malaria remains a major public health problem. In the fight against mosquitoes, the vector of this disease, the significant accumulation of active ingredients in ecosystems treated with synthetic insecticides poses an increased environmental problem linked to the very high persistence time.

The aim of this study is to propose an alternative method of vector control using natural products of plant origin.

Three organic extracts (cyclohexanic, chloroformic and methanolic) from the leaves of *Cassia sieberiana* were tested by contact on mosquito larvae of the genus *Anopheles gambiae*. Four doses of each extract were prepared and monitored over 48 hours.

The families of chemical compounds in the various organic extracts are identified using coloring and precipitation tests.

The results obtained show a variable sensitivity of the larvae, reflected in low to very high mortality rates when moving from one extract to another. The chloroform extract was the most effective, with a mortality rate of just under 90% after 48 hours' exposure.

Phytochemical screening reveals that this plant is rich in secondary metabolites such as polyphenols and alkaloids, which could explain its larvicidal activity.

**Keywords:** *Cassia sieberiana*; *Anopheles gambiae*; larvicidal activity; organic extracts.

## 1. INTRODUCTION

Malaria is a parasitic endemic to which 2 billion people worldwide are exposed. In 2022, there are an estimated 247 million cases and 619,000 deaths from malaria worldwide [1]. Most of these deaths will occur in sub-Saharan Africa, with children under five being the worst affected. Senegal is one of the African countries where malaria remains a public health problem, and despite efforts at various levels to reduce its burden, the statistics remain almost unchanged. An average of 300,000 to 350,000 confirmed cases are recorded each year [2].

Mosquitoes are the exclusive vectors of malaria. There are currently 3,500 species of mosquitoes in the world [3]. Only around sixty species, belonging to the genus *Anopheles*, are capable of transmitting plasmodium, the parasite responsible for human malaria.

A number of resources have been made available to combat vector-borne diseases.

These measures range from building villages away from the tides to draining collections of water where mosquito larvae develop.

The advent of insecticides raised hopes of eradicating malaria from areas where the disease was endemic. Thus, on the initiative of the WHO, a vast malaria eradication program had been launched in the years since the 1950s. The active ingredients of the insecticides used in mosquito control campaigns belong to the

following families: organophosphates, synthetic pyrethroids, organochlorines and synthetic carbamates [4].

When cases of resistance appeared, all hope of eradication was dashed. Although these preparations have proved highly effective against Culicidae mosquitoes, they have a number of drawbacks. The significant accumulation of active ingredients in treated aquatic and terrestrial ecosystems poses a pollution problem.

Furthermore, the active substances in the products used have a broad spectrum of action and do not spare non-target organisms.

In addition to all these disadvantages, there is also the major problem of the development of resistance to chemical insecticides in mosquitoes [5]. As a result, the traditional method of chemical mosquito control is increasingly giving way to biological control.

Furthermore, a systematic examination of phytochemical discoveries listed using the NAPRALERT (Natural Products Alert database) reveals that only less than 30% of plant species have been examined in detail. Given that 65% of plant biodiversity is tropical, it is clear that a large number of molecules of plant origin remain to be discovered.

As part of this vector control program, research has been carried out in our laboratory into the toxicity of a number of native plant extracts.

Three *Cassia sieberiana* leaf extracts were tested on *Anopheles gambiae* mosquito larvae.

## 2. MATERIALS AND METHODS

### 2.1 Equipment

#### 2.1.1 Plant material

*Cassia sieberiana* leaves are harvested in May 2022 in the THIES region located 70 km from DAKAR (SENEGAL). In order to preserve the efficacy of the active ingredients, drying is carried out immediately after harvesting, in a well-ventilated place and away from light. They are then ground using an electric grinder.

#### 2.1.2 Biological material

The *Anopheles gambiae* larvae tested for toxicity came from stagnant water at the Lycée Blaise Diagne (Dakar/Senegal).

The species collected were sorted and isolated from each other to avoid interspecific interactions. Only stage 3 larvae (L3) were selected for testing.

### 2.2 Methods

#### 2.2.1 Extraction

To extract the active ingredients, 100 g of *Cassia sieberiana* leaf powder was macerated in 1 liter of solvent for 72 h in the dark at room temperature.

Three extraction solvents were used (cyclohexane, chloroform and methanol).

Using a rotary evaporator, all the extracts were concentrated to dryness and then stored at -18°C to prevent any deterioration.

#### 2.2.2 Phytochemical screening

The various extracts were subjected to various phytochemical tests to identify the main chemical groups contained in them, using the standard method based on staining and precipitation reactions. Several types of reagents were used for this purpose [6].

**Characterization of polyphenols:** An infusion was prepared by dissolving 1mg in 20 mL of distilled water. The mixture was left to stand for 25 min and then filtered. A 2 mL sample of the filtrate was taken and placed in test tubes. A few drops of an alcoholic solution of FeCl<sub>3</sub> were

added. The appearance of a blackish-blue color indicates the presence of polyphenols. The same filtrate is also used to characterize flavonoids and tannins.

**Characterization of flavonoids:** First, 1 ml of the previous infusion is added, followed by 1 ml of hydrochloric acid. A few shavings of magnesium are added to this mixture. The appearance of an orange-pink color indicates the presence of flavones, a purplish-pink color indicates the presence of flavanones and a red color indicates the presence of flavanols.

**Catechic tannins or non-hydrolysable tannins:** 1 mL of the infusion is added to 0.5 mL of Stiasny's reagent (a mixture of methanal and hydrochloric acid). The resulting mixture was heated to 90° C for 15 min. The appearance of a precipitate indicates the presence of catechic tannins.

**Gallic or condensed tannins:** To reveal the gallic tannins, the contents of the tubes after the catechin tannin test were first filtered, then 0.5 ml of the filtrate was saturated with sodium acetate and finally 0.2 mL of a 2% solution of FeCl<sub>3</sub> was added to this mixture. The appearance of a blue-green color indicates the presence of gallic tannins.

**Alkaloids:** 1 mg of each dry extract was dissolved in 20 mL of 10% concentrated sulphuric acid. A few drops of Dragendoff's reagent were added to this mixture. The appearance of an orange-red precipitate indicates the presence of alkaloids.

### 2.3 Toxicity Tests

The methodology of our tests is inspired by the sensitivity test technique standardized by the World Health Organization [7].

The doses were prepared by dissolving dry alcoholic extracts (such as methanolic extract), chloroform and cyclohexane in dimethyl sulphoxide (DMSO). Four doses were prepared for each extract: dose 1 (50 mg/mL), dose 2 (25 mg/mL), dose 3 (12.5 mg/mL) and dose4 (6.25 mg/mL). For the tests, 10 stage 3 larvae were placed in glass beakers in 20 mL of water from the gites. Using a pipette, 0.5 mL of each of the previously prepared doses was poured into the beaker, left at room temperature and the number of deaths determined at the following times: 24 h, 48 h of exposure.

At each concentration, the test is repeated 4 times. For each test, we use a control tube containing 10 larvae and 20 mL of gite water.

Dead larvae were considered to be those that were immobile and had fallen to the bottom of the beaker. The observation was made with the naked eye. The larval mortality rate was corrected using Abbott's formula (1925).

Statistical analysis of the means was carried out using the ANOVA analysis of variance test with Minitab version 19 software.

### 3. RESULTS AND DISCUSSION

#### 3.1 Phytochemical Screening

The results of the phytochemical tests are given in the Table 1.

From the results obtained in the Table 1, it should be noted that the methanolic extract of *Cassia sieberiana* leaves is very rich in polyphenols, flavonoids and tannins (condensed and hydrolysable).

Condensed tannins are only slightly present in the cyclohexane extract and completely absent in the chloroform extract.

Alkaloids are also present in small quantities in the chloroform extract of the leaves.

The work of K. S. EVENAMEDE et al. [8] also showed that alcoholic extracts of leaves of the same species from Togo also contain flavonoids, polyphenols and tannins. However, alkaloids are absent in the Togo species. This can be explained by a difference in several parameters either geographical, physicochemical or biological such as: the difference in the harvesting site including the environment of the plant, the light, the season, but also and above all the harvesting period and the extraction procedure used [9].

The results obtained in this study are similar to those of Eliasse Zongo et al. [10] whose work was recently carried out (January 2023) in the Hauts-Bassins region of Burkina Faso. They found virtually the same composition of secondary metabolites in the species harvested.

#### 3.2 Biological Tests

The results of the larvicidal activity of the different extracts tested on *Anopheles gambiae* are presented in Figs. 2 and 3 and Table 2.

Table 1. Results of phytochemical screening

Extraction solvents	Polyphenols	Flavonoids	Tannins Condense	Tannins Hydrolysable	Alkaloids
Cyclohexane	-	-	+	-	-
Chloroform	-	-	-	-	+
Methanol	+	+	+	+	-

+ present    - Absent



Fig. 1. Phytochemical testing equipment in the laboratory





Fig. 2. Mortality curve as a function of extract, dose and time



Fig. 3. Interaction between extracts and doses, extracts and time, doses and time

Table 2. Analysis of variance

Source	DL	SomCar séq	Contribution	SomCar ajust	CM ajust	F value
extracts	2	482,33	41,75%	482,33	241,167	78,42
doses	3	290,92	25,18%	290,92	96,972	31,53
temp	1	108,38	9,38%	108,38	108,375	35,24
Error	89	273,71	23,69%	273,71	3,075	
Inadequate fit	17	243,71	21,09%	243,71	14,336	34,41
Pure error	72	30,00	2,60%	30,00	0,417	
<b>Total</b>	<b>95</b>	<b>1155,33</b>	<b>100,00%</b>			
Source	Value of p					
extracts	0,000					
doses	0,000					
temp	0,000					
Error						
Inadequate fit	0,000					
Pure error						
<b>Total</b>						



**Fig. 4. Biological testing equipment in the laboratory**

After exposing *Anopheles gambiae* larvae to different concentrations of the three extracts for 48 hours, the mortality rate varied according to the concentration. The higher the concentration, the greater the mortality.

The analysis of variance of the mean mortality of the larvae shows that there is a significant difference between the 4 doses of extracts used ( $P < 0.05$ ).

The results obtained show a variable sensitivity of the larvae, reflected in low to very high mortality rates when moving from one extract to another. The chloroform extract proved to be the most effective, with a mortality rate of just under 90% after 48 hours' exposure. This activity could be explained by the presence of alkaloids in the chloroform extract, detected during phytochemical tests. Matasyoh et al. [11] showed a high larvicidal efficacy of alkaloids extracted from the leaves of *Zanthoxylum leprieurii*.

These results also show that larvicidal activity depends on the length of time the larvae are exposed to the different extracts. In fact, an increase in mortality was recorded as a function of exposure time.

Analysis of the interaction diagrams shows that all the extracts are equally sensitive to dose1 (100% mortality after 48 hours' exposure). In the short term, the chloroform extract was slightly more effective than the alcoholic extract. However, in the long term, the opposite was observed.

The results obtained for the main effects of the extracts are perfectly consistent with the observations reported by Issoufou et al. [12] who state in their study on Phytochemical Characterization and Larvicidal Activity of Raw Plant Extracts from the Traditional Pharmacopoeia of Niger on *Anopheles gambiae*

Larvae that lethality increases as a function of the concentration and duration of exposure of the larvae.

Research conducted by Aziz Bouchelta [13] on the biocidal effects of alkaloids, saponins and flavonoids extracted from *Capsicum frutescens* L but also the work of Ndiaye. A [14] on phytochemical testing and larvicidal activity of three organic extracts of *Indigofera pilosa* stems on mosquito larvae revealed that alkaloid-containing extracts were among the most effective. These previous studies corroborate the results of the present study.

#### **4. CONCLUSION**

Mosquitoes have always been considered a source of nuisance for humans, mainly because they can be vectors of disease.

In order to solve the problems linked to the use of chemical insecticides and their harmful impact on health and the environment, the use of natural alternatives that fulfil the same role and offer ecological advantages is necessary, hence the purpose of this study.

These results show good larvicidal activity of the three *Cassia sieberiana* leaf extracts. All the extracts tested constitute promising larvicides for the control of mosquitoes and by ricochet against malaria even if the lethality rate varies from one extract to another.

In the future, we plan to fractionate and purify the most active extracts in order to isolate the molecule(s) responsible for the biological activity.

#### **COMPETING INTERESTS**

Authors have declared that no competing interests exist.



## REFERENCES

1. WHO. Treatment of uncomplicated *P. falciparum* malaria in the first trimester of pregnancy: Implementation of revised WHO guidelines (25 November 2022); 2023. Consulted in November 2023. Available: <https://www.who.int/publications/i/item/guidelines-for-malaria>.
2. Seddoh A, Onyeze A, Gyapong JO, Holt J, Bundy D. Towards an investment case for neglected tropical diseases. Lancet. 2013;9908:1898-1955.
3. Heu K, Gendrin M. Mosquito microbiota and its influences on disease vectorial transmission. Biol Aujourd'hui. 2018;212(3-4):119-136.
4. Maifi Radhia, Salmi Meriem. Study of the larvicidal activity of aqueous extract of *Ruta graveolens* against a species of mosquito *Culesita longiareoleta*. University of larbi tébessi -tébessa; 2017.
5. Mouchet J, Pichon G, Gayral P, Hamo J. Sensitivity and resistance to insecticides of *Aedes aegypti* in West Africa and methods of controlling this vector. Bull. Org. world. Health. 1963; 1971(145):394-404.
6. Houmènou V, Adjatin A, Assogba F, Gbénou J, Akoègninou A. Phytochemical and cytotoxicity study of some plants used in the treatment of female infertility in South Benin. European Scientific Journal. ESJ. 2018;14(6):156.
7. World Health Organization, et al. Test procedures for insecticide resistance monitoring in malaria vector mosquitoes; 2016.
8. Evenamede KS, Kpegba K, Idoh K, Amegnona A, Simalou O, Boyode P, Oke OE, Gbeassor M. Comparative study of the toxicity of hydroethanolic extracts of the root and stem barks of *Cassia sieberiana* D.C. on wistar rats. JApp Biol Biotech. 2019;7(03):47-52.
9. Malik, Farnaz, Hussain, Shahzad, Sadiq, Alia, et al. Phyto-chemical analysis, anti-allergic and anti-inflammatory activity of *Mentha arvensis* in animals. Afr J Pharm Pharmacol. 2012;6(9):613-619.
10. Zongo E, Busuioc A, Meda RN, Botezatu AV, Mihaila MD, Mocanu AM, Avramescu SM, Koama BK, Kam SE, Belem H, Somda FLS, Ouedraogo C, Ouedraogo GA, Dinica RM. Exploration of the antioxidant and anti-inflammatory potential of *Cassia sieberiana* DC and *Piliostigma thonningii* (Schumach.) Milne-Redh, traditionally Used in the treatment of hepatitis in the hauts-bassins region of Burkina Faso. Pharmaceuticals (Basel). 2023Jan 16;16(1):133.
11. Matasyoh JC, Dittrich B, Schueffler A, Laatsch H. Larvicidal activity of metabolites from the endophytic podospora sp. against the malaria vector *Anopheles gambiae*. Parasitol Res. 2011, Mar;108(3):561-6.
12. Yolidjé I, Keita DA, Moussa I, Toumane A, Maarouhi IM, Saley K, Pirat J-L, Much T, Ouamba JM. Phytochemical characterization and larvicidal activity of raw plant extracts iss of the traditional pharmacopoeia of niger on the larvae of *Anopheles gambiae* S.L. ESJ. 2019; 15(12):30.
13. Bouchelta A, Boughdad A, Blenzar A. Effets biocides des alcaloïdes, des saponines et des flavonoïdes extraits de *Capsicum frutescens* L. (Solanaceae) sur *Bemisia tabaci* (Gennadius) (Homoptera : Aleyrodidae). Biotechnologie, Agronomie, Société et Environnement. 2005;(9):259-269.
14. Ndiaye A, Diop M, Diouf El HG. Phytochemical test and larvicidal activity of three organic extracts exits of the stems of *Indigofera pilosa* on larvae of mosquitoes. Open Access Library Journal. 2017; (4):e3203.

© 2024 Kébé et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/112651>