



Coordination of N, O-donor Appended Schiff Base Ligand *N, N'*-bis (3-Methoxy Salicylideneimino-1,3-diaminopropane) towards Nickel (II), Copper (II) and Zinc (II) : Synthèses, Crystal Structure

Moussa Faye ^a, Mariama Sarr ^a, Mbossé Ndiaye Guèye ^b,
Farba Bouyagui Tamboura ^a, Ibrahima Elhadj Thiam ^b,
Simon Coles ^c, James Orton ^c, Moussa Dieng ^{a*}
and Mohamed Gaye ^b

^a Department of Chemistry, UFR SATIC, Alioune DIOP University, Bambey 21400, Senegal.

^b Department of Chemistry, Cheikh Anta Diop University, Dakar 10700, Senegal.

^c UK National Crystallography Service, School of Chemistry, Faculty of Engineering and Physical Sciences, University of Southampton, Southampton SO17 1BJ, United Kingdoms.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IRJPAC/2024/v25i3853

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/116100>

Original Research Article

Received: 15/02/2024

Accepted: 19/04/2024

Published: 23/04/2024

*Corresponding author: E-mail: moussa.dieng@uadb.edu.sn;

ABSTRACT

New complexes prepared by the reaction of *N,N'*-bis(3-methoxysalicylideneimino-1,3-diaminopropane) (H_2L) with Ni(II), Zn(II) and Cu(II) ions are reported in this paper. The H_2L ligand is structurally characterized by elemental analysis, NMR, infrared, UV-Vis spectroscopies, conductance and magnetic room temperature measurement. The mononuclear complex of Ni(II) ($C_{19}H_{24}N_2NiO_6$) is characterized by single X-ray diffraction. The compound crystallizes in the orthorhombic system in the space group *Pnma* with the unit cell parameters $a = 7.3754(5)$ Å, $b = 21.9576(13)$ Å, $c = 11.5546(6)$ Å, $\alpha = 90^\circ$, $\beta = 90^\circ$, $\gamma = 90^\circ$. The ligand H_2L acts in tetradentate fashion in its di-deprotonated form. Two coordinated water molecules complete the coordination sphere. The environment around the Ni(II) center is best described as an octahedral geometry.

Keywords: Schiff base; X-ray; mononuclear complex; tetra dentate.

1. INTRODUCTION

“The use of Schiff bases in coordination chemistry, to develop from molecular units with specific properties has undergone spectacular development in recent years thanks to very elaborate synthesis methods which make it possible to control the synthesis of complexes with original structures. Polyfunctional acyclic, symmetric, or asymmetric ligands and their complexes represent an essential component of this chemistry” [1–6]. “These ligands are widely used for the synthesis of complexes with very interesting biochemical and physical properties” [7–10]. “The formation of acyclic ligand complexes depends on the flexibility of the arms bearing the groups having donor sites, the nature of the donor atoms and the complexation properties of the cations and anions playing the role of charge balance” [11–14]. “These Schiff bases have made it possible to develop molecular materials with specific properties” [15–19]. “This article describes the synthesis of a tetradentate Schiff base obtained by the reaction between 2-hydroxy-3-methoxybenzaldehyde and 1,3-diamine propane. The flexibility and numerous coordination sites of the resulting acyclic ligands of this type offer several possible structures with lanthanide ions” [20–25].

2. METHODOLOGY

2.1 Synthesis of *N,N'*-bis(3-methoxysalicylideneimino-1,3-diaminopropane) (H_2L)

The Schiff base *N,N'*-bis(3-methoxysalicylideneimino-1,3-diaminopropane) (H_2L) was prepared with slight modification of literature procedures. 1,3-diamine propane dissolved in methanol was refluxed with an methanolic solution of 2-hydroxy-3-

méthoxybenzaldéhyde, in the presence of few drops of glacial acetic acid. In a 250 ml flask, dissolve 3.04 g (20 mmol) of *o*-vanillin or (2-hydroxy-3-methoxybenzaldehyde) in methanol. Add 0.83g (10 mmol) of 1,3-diamine propane suspended in methanol. A yellow solution was obtained. Then heated to reflux temperature and was maintained for 4 hours. On cooling the reaction mass, yellow precipitate was separated. The compound was thoroughly washed with ether and dried over P_2O_{10} .

Yield: 77.15%; Melting temperature: 140 - 141°C


RMN 1H : (500 MHz, DMSO- d_6 , δ , ppm) : 3.71 (s, 6H, CH_3), 6.8-6.62 (m, 6H, Ar-H), 2.53 (t, 2H, CH), 1.63 (s, 2H, OH), 3.79 (t, 4H, CH_2).

RMN ^{13}C : (125 MHz, DMSO- d_6 , δ , ppm): 55,47 (C_2 ; C_2'), 117,88 (C_6 ; C_6'), 120,041 (C_4 ; C_4'), 120,47 (C_5 ; C_5'), 147,38 (C_3 ; C_3'), 146,82 (C_9 ; C_9'), 46,11 (C_{10}), 111,00 (C_7 ; C_7'), 28,82 (C_{11}), 147,38 (C_8 ; C_8').

IR: ν (cm^{-1}) : 1628 ($C=N$)_{imine} ; 3078 (H-C=N) ; 3189 (OH), (1466 - 1358) (C=C) ; 1245 (C-O), 2946 (C-H).

2.2 Synthesis of the Complexes

In a 100 mL flask containing 20 mL of ethanol, dissolve 0.15 g (0.44 mmol) of the H_2L ligand. Add (0.44 mmol) metal nitrate x hydrated $M(NO_3)_2 \cdot xH_2O$ with ($M = Ni, Cu, Zn$ and $x = 3; 4; 6$) previously dissolved in 10 mL of ethanol. A precipitate immediately forms for all complexes. The mixture obtained is stirred and heated to reflux for three hours. Collect the precipitates by filtration, wash with ether and dry in the open air. Leave the filtrates to slowly evaporate for a few days; green crystals form for the nickel (II) complex filtrate. Yields range from 78.7% to 97.77%.



Scheme 1. Synthesis of the ligand H₂L





Fig. 1. H₂L ligand



Scheme 2. Synthesis of nickel (II) complex

3. RESULTS AND DISCUSSION

3.1 General Studies

The infrared spectrum of the ligand reveals the disappearance of the intense band pointed at 1637 cm^{-1} attributed to the $\nu(\text{C}=\text{O})$ vibration of o-vanillin and the appearance of a new strong band at 1628 cm^{-1} corresponding to the vibration $\nu(\text{C}=\text{N})$. This proves that condensation has indeed taken place. The bands attributed to the C-H vibrations of the aromatic skeleton are noted at 3078 , 1076 , 958 and 833 cm^{-1} [26]. The valence vibration of the phenolic (C-O) bond is pointed at 1245 cm^{-1} and those of the CH₂ groups at 2843 cm^{-1} . The very strong band at

730 cm^{-1} accompanied by a shoulder justifies the presence of an aromatic core. We observe a broad band at 3189 cm^{-1} attributable to the $\nu(\text{OH})$ vibrations of the ligand.

The analysis of infrared data of the H₂L ligand complexes reveals a decrease towards low frequencies of the vibration bands characteristic of the C=N bonds of the C-O imine group of the phenol group. This reduction ranging from 1628 to 1612 cm^{-1} for the imine group and 1245 to 1216 cm^{-1} testifies to the coordination of the electron donor atoms of these groups to the metal [27]. However, despite the slight reduction in the vibration band characteristic of the C-O bond of the methoxy group observed at 1076 cm^{-1}

¹ on the spectrum of the ligand, its donor atom does not participate in the coordination of the metal. On the spectra of all complexes, we note the presence of new vibration bands in the regions (1470 – 1762), (1298 – 1292) and (1040 – 1008) cm⁻¹. These triplet bands indicate the presence of coordinated nitrate in the complexes. The magnitude values estimated respectively at 169, 172 and 167 cm⁻¹ for the copper, nickel and zinc complexes prove that the nitrates are coordinated to the metal in a bidentate chelate manner. On the spectrum of the zinc complex, we also observe a characteristic vibration band of free nitrate group at 1367 cm⁻¹.

The study of the molar conductivity of these complexes in a millimolaire solution of DMF confirms the position of the nitrate groups in the complexes. Indeed, the newly found values for the copper and nickel complexes indicate that their electrolytes are of neutral type while that of the zinc complex gives an electrolyte of type 1: 1. Fifteen days later the values found show that the complexes are all stable in DMF [28].

The UV-visible spectra of the complexes are recorded in a millimolaire solution of DMF. They present bands attributed to the $\pi \rightarrow \pi^*$, $n \rightarrow \pi^*$ transitions of the chromophore groups and charge transfer bands from the ligand to the metal (CTLM). These transitions are observed in the range 410 – 460 nm. In addition to these bands, we note on the spectrum of the copper complex a broad band pointed at 630 nm which characterizes the $d \rightarrow d$ transitions of a copper ion in an octahedral environment [29]. This environment of the metal is confirmed by the magnetic moment which is equal to 1.77 μ_B , a value which is typical of an octahedral copper mononuclear complex [30].

On the spectrum of the nickel (II) complex, the broad band observed at 638 nm is attributed to the sum of the transitions ${}^3A_{2g} \rightarrow {}^3T_{2g}$ and ${}^3A_{2g} \rightarrow {}^3T_{1g}$ of an octahedral nickel (II) complex [30]. The magnetic moment found is 2.76 μ_B . This value is that of a mononuclear nickel complex in an octahedral environment.

On the spectrum of the zinc (II) complex the charge transfer band from the ligand to the metal is observed at 460 nm. It suggests the presence of zinc in the product.

X-ray Crystallography: Details of the crystal structure solution and refinement are given in Table 2. Single-crystal X-ray diffraction data were measured on a Rigaku Synergy S diffractometer at the MoK α radiation ($\lambda = 0.71073 \text{ \AA}$).

Crystallographic analysis of the [Ni(HL)(H₂O)₂] complex reveals that it crystallizes in an Orthorhombic system with a Pnma space group. The unit cell parameters recorded in Table 2 are $a = 7.3754 (5) \text{ \AA}$; $b = 21.9576 (13) \text{ \AA}$; $c = 11.5546 (6) \text{ \AA}$ and $\alpha = \beta = \gamma = 90^\circ$. The asymmetric unit consists of a mononuclear complex in which a deprotonated ligand (L⁻) coordinates through its two iminic nitrogen atoms and its two phenolate oxygen atoms forming an ONNO equatorial plane around the metal. However, it should be noted that the axial plane of the octahedral environment of nickel Ni²⁺ is formed by two water molecules. Analysis of crystallographic data reveals that the angles around nickel (II) in the equatorial plane O2-Ni-N1, N1-Ni-N1i and O2-Ni-O2i are respectively 91.93(7)°, 97.38(11)° and 85.98(10)° while those of the axial plane O2i-Ni-O3 and N1i-Ni-O4 are 91.93(7)° and 89.04(6)°. These values suggest that the octahedral geometry around the nickel (II) is slightly distorted. This observation is confirmed by the lengths of the equatorial and axial connections. Indeed the lengths of these bonds are 2.073(2) \AA for Ni—N1 and 2.0141(17) \AA Ni—O2 all equatorial. For the bond lengths with nickel in the axial plane, the values are 2.112(3) \AA for Ni—O3 and 2.095(2) \AA for Ni—O4, respectively comparable to the axial bond lengths [Ni1—O1W = 2.0955 (19) \AA and Ni1—O2W = 2.1071 (18) \AA with Barr and al. [30]. The crystallographic structure of the complex is stabilized by intermolecular hydrogen bonds of the O-phenol-H---O water type.

Table 1. UV-visible and magnetic data of metal nitrate complexes with H₂L ligand

Complexes	UV-Visible		μ_{eff} (μ_B)
	λ (nm)	Attributions	
[Cu(HL)(NO ₃)]·xH ₂ O	344 ; 392 ; 410 ; 630	$\pi \rightarrow \pi^*$, $n \rightarrow \pi^*$, CTLM, $d \rightarrow d$	1.77
[Ni(HL)(NO ₃)]·xH ₂ O	348 ; 388 ; 408 ; 638	$\pi \rightarrow \pi^*$, $n \rightarrow \pi^*$, CTLM M, $d \rightarrow d$	2.77
[Zn(H ₂ L)(NO ₃)]·(NO ₃)·xH ₂ O	332 ; 385 ; 460	$\pi \rightarrow \pi^*$, $n \rightarrow \pi^*$, CTLM, $d \rightarrow d$	Diam

Table 2. Crystal data and details of the structure determination

Chemical formula	$C_{19}H_{24}N_2NiO_6$
Mr (g/mol)	431.11
Température (K)	100 K
Radiation (Å)	0.71075
Crystal size (mm^3)	0.22 × 0.1 × 0.06
Crystal system	Orthorhombique
Space group	Pnma
Z	4
a ; b ; c (Å)	7.3754 (5) ; 21.9576 (13) ; 11.5546 (6)
α ; β ; γ ($^\circ$)	90
V (Å^3)	1871.22 (19)
Calculated density (Mg.m^{-3})	1.544
Radiation type	Mo K α
μ (mm^{-1})	1.08
F(000)	912
θ ($^\circ$)	3.3 – 31.8
h, k, l	$-9 \leq h \leq 9$; $-24 \leq k \leq 29$; $-15 \leq l \leq 15$
No. of measured reflections, independent and observed [$I \geq 2\sigma(I)$]	2468 ; 1639 ; 2166
Final Indices R [$I > 2\sigma(I)$]	
GOF	1.06
$R[F^2 > 2\sigma(F^2)]$; $wR(F^2)$; R_{int}	0.045 ; 0.120 ; 0.027
No. of parameters	66
$\Delta\rho_{\text{max}}$, $\Delta\rho_{\text{min}}$ ($e \text{ Å}^{-3}$)	0.81 ; -0.68

Table 3. Bond lengths (Å) and angles around Ni (°) in the complex [Ni(H₂L)(H₂O)₂]

Bond	Bond lengths (Å)	Corner	Value (°)
Ni2-N1 ⁱ	2.073(2)	O2-Ni-N1	91.93(7)
Ni2-N1 ⁱ	2.073(2)	N1-Ni-N1 ⁱ	97.38(11)
Ni2-O1	2.0141(17)	O2-Ni-O2 ⁱ	85.98(10)
Ni2-O2 ⁱ	2.0141(17)	O2 ⁱ -Ni-O3	91.93(7)
Ni2-O4	2.095(2)	N1 ⁱ -Ni-O4	89.04(6)
Ni2-O3	2.112(3)	O2-Ni-N1 ⁱ	174.29(7)

Symmetry code : (i) x ; -y-1/2, z

Table 4. Intermolecular hydrogen bond lengths in (Å) for the complex $[\text{Ni}(\text{H}_2\text{L})(\text{H}_2\text{O})_2]$

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{O4}-\text{H4A} \cdots \text{O2}^i$	0.88	1.99	2.75(3)	148.6

Symmetry code : (i) $x-1/2, -y-1/2, -z+3/2$

3.2 Structure Description of the Complex of Nickel (II)




Fig. 2. Crystallographic structure of the complex $[\text{Ni}(\text{L})(\text{H}_2\text{O})_2]$




Fig. 3. Three-dimensional structure of the complex $[\text{Ni}(\text{L})(\text{H}_2\text{O})_2]$




Fig. 4. Proposed structure of copper, nickel and zinc (II) complexes with the H₂L ligand

3.3 Proposed Other Structures

Based on spectroscopic, magnetic and conductometric analyses, we propose the following structures:

- The complexes are mononuclear.
- In the structure of the complexes, the ligand acts in a tetradentate manner through its two phenolic oxygen atoms and two iminic nitrogen atoms.
- The nitrate ions are coordinated to the metal in a bidentate chelate manner and a free nitrate which plays the stabilizing role in the zinc (II) complex.
- The complexes contain free water molecules.

4. CONCLUSION

The synthesis and characterization of the Schiff base ligand *N,N'*-bis(3-methoxysalicylideneimino-1,3-diaminopropane), three metal transition complexes were synthesized and studied using spectroscopic and spectrophotometric methods as well as X-ray diffraction technique for nickel (II) complex. The dianionic ligand acts in tetradentate mode via two azomethine nitrogen atoms and two oxygen phenolate atoms. Infrared spectroscopy confirms the coordination between the Schiff base ligand and the cations Ni(II), Cu(II) and Zn(II). The structure of nickel (II) complex is determined by single crystal x-ray diffraction. The environment around each cation is octahedral. UV-Vis spectrophotometry indicates internal transition in the ligand and metal. This paper describes a mononuclear nickel (II) complex. The structure reveals tetradentate coordination of the ligand via two nitrogen atoms and two deprotonated phenolic oxygen atoms. In this complex the central metal center is hexacoordinated. These complexes

were characterized by spectroscopic studies (IR, UV-Vis), molar conductivity and room temperature magnetic moment measurements. Suitable X-ray crystal diffraction were obtained for the Nickel (II) complex.

Nitrate ion complete the sphere of coordination forming Ni(II), Zn (II) and Zn(II) complexes with a coordination number of six. Sometime later two water molecules replaced the nitrate ion in Nickel (II) complex. The environment around the metal center is best described as a as a octahedral geometry. The ligands show similar fashion of coordination mode for each metal ion, acting as tetradentate. Each metal center is eight coordinated with two nitrogen atom from the ligand and two oxygen atom from two water molecules. The environment around each metal for both complexes is best described as a octahedral geometry.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES


1. Ibers JA, Holm RH. Synthetic metal complexes can closely approach the properties of metalions in proteins and yield useful information concerning biological structure and function. Modeling coordination sites in mettallobiomolecules. 1980;209:4453. DOI:10. 1126/science.7384796
2. Caki N, Gündüz S, Rengarasu R, Angelovski G. Synthetic strategies for preparation of cyclen-based MRI contrast agents. Tetrahedron Lett. 2015;56:759–765. Available:https://doi.org/10.1016/j.tetlet.2014.12.087

3. Fenton DE, Casellato U, Vigato PA, Vidali M. Acyclic and macrocyclic schiff base complexes of lanthanides and actinides. *Inorg. Chim. Acta.* 1984;95:187. Available:[https://doi.org/10.1016/S0020-1693\(00\)87465-4](https://doi.org/10.1016/S0020-1693(00)87465-4)
4. Dieng M, Thiam I, Gaye M, Sall AS, Barry AH. Synthesis, Crystal Structures and Spectroscopic Properties of a Trinuclear [Cu₃(HL)₂(NO₃)₂](H₂O)(CH₂CH₂OH) Complex and a [Mn(HL)(CH₃COO)]_n Polymer with H₃L = N, N'-(2-hydroxypropane-1,3-diyl)-bis-(salicylaldimine). *Acta chim. Slov.* 2006;53:417-423 Available:<http://acta-arhiv.chem-soc.si/53/53-4-417.pdf> Available:<http://cat.inist.fr/?aModele=afficheN&cpsid=18358989>
5. Diallo M, Dieng M, Gaye M, Sall AS, Barry AH, Chahrazed B. Aquamalonato (1,10-phenanthroline) copper (II) sesquihydrate. *Acta. Cryst.* 2007;63:1810-m 1811 DOI:10.1107/S160053680702332X
6. Dieng M, Barry AH, Gaye M, Sall AS, Pérez-Lourido P, Valencia-Matarranz L. (2-[[2-(2-Aminoethylamino)ethylimino]-methyl]phenolato-k⁴O,N',N'',N''')-copper(II) perchlorate. *Acta. Cryst.* 2011;67:830-m 831 DOI:10.1107/S1600536811020095
7. Winkler ME, Spira DJ, LuBein CD, Thamann TJ, Solomon EI. Anion binding to oxidized type 2 depleted and native laccase: A spectroscopically effective model for exogenous ligand binding to the type 3 - type 2 active site. 1982;107(2):727-734. Available:[https://doi.org/10.1016/0006-291X\(82\)91551-0](https://doi.org/10.1016/0006-291X(82)91551-0)
8. Gueye MN, Dieng M, Thiam IE, Lo D, Barry AH, Gaye M, Retailleau P. Lanthanide (III) Complexes with Tridentate Schiff Base Ligand, Antioxidant Activity and X-Ray Crystal Structures of the Nd(III) and Sm(III) Complexes, *South African Journal of Chemistry.* 2017;70:8-15. Available:<http://dx.doi.org/10.17159/0379-4350/2017/v70a2>
9. Gueye MN, Dieng M, Lo D, Thiam I, Barry AH, Gaye M, Sall AS, Retailleau, P. Synthesis, physical studies and crystal structure determination of Y(III) and Er(III) complexes of 1-(pyridin-2-yl)-2-(pyridine-2-ylmethylene)hydrazine. *European Journal of Chemistry.* 2017;8(2): 137-143. Available:<https://doi.org/10.5155/eurjchem.8.2.137-143.1557>
10. Majumdar D, Biswas JK, Mondal M, Babu MSS, Metre RK, Das S, Bankura K, Mishra D. Coordination of N,O-donor appended Schiff base ligand (H₂L₁) towards Zinc antimicrobial activities and Hirshfeld surfaces. *Journal of Molecular Structure.* 2018;1155:745-757. Available:<https://doi.org/10.1016/j.molstruc.2017.11.052>
11. Tinguiano D, Thiam I, Dieng M, Gaye M, Retailleau P. Dichlorido[2-pyridin-2-yl]-N-(pyridine-2-ylmethyl-idene)ethanamine-k³N,N',N''] manganese (II) monohydrate. *Acta. Cryst.* 2012;68:1250 DOI:10.1107/S1600536812037877
12. Ishak NNM, Jamsari J, Ismail AZ, Tahir MIM, Tiekink ERT, Veerakumarasivam A, Ravooof TBSA. Synthesis, characterisation and biological studies of mixed-ligand nickel (II) complexes containing imidazole derivatives and thiosemicarbazide Schiff bases, *J. Mol. Struct.* 2019;1198:126888 DOI:10.1016/j.molstruc.2019.126888
13. Deun RV, Binnemans K. Lanthanide containing Schiff's base complexes with chloride counter-ions: Mesomorphic properties. *Materials Science and Engineering: C.* 2001;18(1,2):211-215 Available:[https://doi.org/10.1016/S0928-4931\(01\)00360-5](https://doi.org/10.1016/S0928-4931(01)00360-5)
14. Vashistha N, Chandra A, Singh M. Influence of rhodamine B on interaction behavior of lanthanide nitrates with 1st tier dendrimer in aqueous DMSO: A physicochemical, critical aggregation concentration and antioxidant activity study. *Journal of Molecular Liquids.* 2018;260:323-341 Available:<https://doi.org/10.1016/j.molliq.2018.03.056>
15. Refat MS, El-Metwaly NM. Spectral, thermal and biological studies of Mn(II) and Cu(II) complexes with two thiosemicarbazide derivatives, *Spectrochim. Acta Part A.* 2012;92:336-346. Available:<https://doi.org/10.1016/j.saa.2012.02.041>
16. Gueye MN, Dieng M, Thiam IE, Lo D, Barry AH, Gaye M, Retailleau P. Lanthanide (III) Complexes with Tridentate Schiff Base Ligand, Antioxidant Activity and X-Ray Crystal Structures of the Nd (III) and Sm (III) Complexes. *South African Chemical Institute.* 2017;70:8-15.


- DOI:<http://dx.doi.org/10.17159/03794350/2017/v70a2>
17. Lo D, Diaw M, Gaye PA, Dieng M, Diouf O, Barry AH, Sall AS, Gaye M. Synthesis and Crystal Structure of Nickel (II) and Zinc (II) Complex From 1,5-Bis [(2-Thiophenyl) Methylidene] Thiocarbonohydrazide. IOSR Journal of Applied Chemistry (IOSR-JAC). 2017;10:83-89.
DOI:10.9790/5736-1001018389
 18. Gueye MN, Dieng M, Lo D, Thiam IE, Barry AH, Gaye M, Sall AS, Retailleau P. Synthesis, physical studies and crystal structure determination of Y (III) and Er (III) complexes of 1-(pyridin-2-yl)-2-(pyridine-2-ylmethylene) hydrazine. European Journal of Chemistry. 2017;8:137-143
Available:<http://dx.doi.org/10.5155/eurjchem.8.2.137-143.1557>
 19. Gueye MN, Dieng M, Thiam IE, Barry AH, Gaye M, Retailleau P. Crystal structure of bis (acetato-k²O,O')diaqua[1(pyridin-2-ylmethylidene-kN)-2-(pyridin-2-yl-kN)hydrazine-kN¹] terbium (III) nitrate monohydrate. Acta Cryst. 2017;73:1121-1124.
Available:<https://doi.org/10.1107/S2056989017009653>
 20. Faye M, Gaye PA, Sow MM, Dieng M, Tamboura FB, Gruber N and Gaye M. Synthesis, Characterization and Single Crystal X-ray Crystallography of Nd (III) and Pr (III) Complexes with the Tridentate Schiff Base Ligand N'-(1-(pyridin-2-yl)ethylidene) nicotinothiazide. Earthline Journal of Chemical Sciences. 2021;6(1):99-117.
Available:<https://doi.org/10.34198/ejcs.6121.99117>
 21. Faye M, Kama AB, Gueye MN, Dieng M, Tamboura FB, Gautier R, Gaye M. Synthesis and characterization of a coordination polymer [Ce(HL)(NO₃)₂(H₂O)]_n with H₂L = N-[(2-hydroxyphenyl) methylidene] pyridine-3-carbohydrazide. Science Journal of chemistry. 2022;10(5): 161-169.
DOI:10.11648/j.sjc.20221005.14
 22. Barr F, Camara PS, Gueye A, Zazouli S, Gruber N, Tamboura FB, Dieng M, Gaye M. Spectroscopic Studies and X-ray Structural of Dinuclear Lanthanide (III) Complexes Derived from N'-(2-hydroxy-3-methoxybenzylidene) Nicotinothiazide. Science Journal of Chemistry. 2023;11(2):56-63.
DOI:10.11648/j.sjc.20231102.13
 23. Faye M, Gueye MN, Dieng M, Tamboura FB, Gaye M. Crystal structure of Y(III) Complex with the Tridentate Schiff Base Ligand N'-(1-(pyridin-2-yl)ethylidene) nicotinothiazide. International Journal of Novel Research in Physics Chemistry and Mathematics. 2023;10(1):60-66.
DOI:<https://doi.org/10.5281/zenodo.7876199>
 24. Camara PS, Guéye A, Traoré B, Zazouli S, Gruber N, Thiam IE, Tamboura FB, Dieng M and Gaye M. Preparation, Characterization of Some Lanthanide(III) Complexes of N'-(phenyl (pyridin-2-yl)methylene)nicotinothiazide. Earthline Journal of Chemical Sciences. 2023;10(1):136-146
Available:<https://doi.org/10.34198/ejcs.10123.131146>
 25. Baraniak E, Bruce RSL, Freeman HC, Hair NJ, James J. Structure, and properties of a dimeric hydroxo-bridged ytterbium(III) complex, di-μ₂-hydroxo-bis[triaquo(pyridine-2-carboxaldehyde-2'-pyridylhydrazone)ytterbium(III)] tetrachloride tetrahydrate. Inorganic Chemistry. 1976;15(9):2226-2228.
Available:<https://doi.org/10.1021/ic50163a046>
 26. Nakamoto, K. Infrared Spectra of Inorganic and Coordination Compounds second edition; 1970.
Available:<https://doi.org/10.1002/bbpc.19710750622>
 27. Tamboura FB, Haba P, Gaye M, Sall AS, Barry AH, Jouini T. Structural studies of bis-(2,6-diacetylpyridine-bis-(phenylhydrazone)) and X-ray structure of its Y(III), Pr(III), Sm(III) and Er(III) complex. Polyhedron. 2004;23:1191–1197.
Available:<https://doi.org/10.1016/j.poly.2004.01.014>
 28. Geary WJ. The use of conductivity measurements in organic solvents for the characterisation of coordination compounds, Coordination Chemistry Reviews. 1971;7:81-122
Available:[https://doi.org/10.1016/S0010-8545\(00\)80009-0](https://doi.org/10.1016/S0010-8545(00)80009-0)
 29. Roztocki K, Matoga D et Nitek W. Cobalt(II) compounds with acetone isonicotinoyl hydrazone tautomers : Syntheses and crystal structures of complexes with free donor atoms. Inorg. Chim. Acta. 2016;448:86–92.

- Available:<https://doi.org/10.1016/j.ica.2016.03.045>
30. Barr F, Ndiaye NF, Kama AB, Tamboura FB, Dieng M, Alamedji RB, Thiam IE, Barry AH, Claiser N, Souhassou, M and Gaye ML. Synthesis, Characterization, Fluorescence and Antibacterial Activities Studies of Transition Metals Complexes with a Macro Acyclic Schiff Base Ligand. *Chemical Science International Journal*. 2024;33(1):22-35. Article no.CSIJ.111342 DOI:10.9734/CSJI/2024/v33i1882


SUPPLEMENTARY MATERIALS




Spectrum IR of ligand H₂L




Spectrum RMN ¹H




SPECTRUM RMN ¹³C




Spectrum IR of complex $[Cu(L)(NO_3)] \cdot nH_2O$




Spectrum IR of complex $[Ni(L)(NO_3)] \cdot nH_2O$




Spectrum IR of complex $[Zn(L)(NO_3)] \cdot nH_2O$



Spectrum UV-Vis of complex [Cu(L)(NO₃)]·nH₂O



Spectrum UV-Vis of complex [Ni(L)(NO₃)]·nH₂O



Spectrum UV-Vis of crystal [Ni(L)]·2H₂O

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/116100>