



# **Micronutrients in the Plant-Soil-Environment Continuum: A Comprehensive Review**

**Himanshu Tiwari <sup>a++</sup>, Rajan Bhatt <sup>b</sup>, P. K. Singh <sup>c#</sup>,  
Dhritiman Das <sup>d†</sup>, Shivam Vihan <sup>e‡</sup>, Tarun Tomar <sup>a^\*</sup>,  
Abhishek Kumar Singh <sup>a^</sup> and Suneel Kumar <sup>f^</sup>**

<sup>a</sup> Department of Agronomy, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut, U.P., India.

<sup>b</sup> Krishi Vigyan Kendra, Amritsar, Punjab Agricultural University, Ludhiana, Punjab, India.

<sup>c</sup> Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut, U.P., India.

<sup>d</sup> School of Smart Agriculture, Adamas University, Kolkata, West Bengal, India.

<sup>e</sup> College of Agriculture, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur, U.P., India.

<sup>f</sup> Department of Soil Science and Agricultural Chemistry, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut, U.P., India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: <https://doi.org/10.9734/ejnf/2024/v16i71465>

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/118164>

**Review Article**

**Received: 05/04/2024**

**Accepted: 07/06/2024**

**Published: 17/06/2024**

<sup>++</sup> Guest Faculty;

<sup>#</sup> Director Extension;

<sup>†</sup> Assistant Professor;

<sup>‡</sup> M.Sc. (Agronomy);

<sup>^</sup> Ph.D. Research Scholar;

<sup>\*</sup>Corresponding author: Email: taruntomar33@gmail.com;

**Cite as:** Tiwari, Himanshu, Rajan Bhatt, P. K. Singh, Dhritiman Das, Shivam Vihan, Tarun Tomar, Abhishek Kumar Singh, and Suneel Kumar. 2024. "Micronutrients in the Plant-Soil-Environment Continuum: A Comprehensive Review". *European Journal of Nutrition & Food Safety* 16 (7):171-78. <https://doi.org/10.9734/ejnf/2024/v16i71465>.

## ABSTRACT

Though mostly ignored, micronutrients are most important. Reducing the negative effects of micronutrient shortages requires the use of effective techniques like vitamin supplements and dietary diversification. It is known that when agricultural yields are reduced because of inadequate soil micronutrient concentration, malnutrition may result. Deficits in soil micronutrients can result from a number of variables, such as the use of intensive cropping practices, micronutrient leaching, soil characteristics, and decreased use of farmyard waste. The availability of micronutrients in the soil is determined by a number of factors, including soil pH, and status of the soil organic matters. Numerous factors, including the structure of root hairs, the release of organic acids, sugar, and several root enzymes, as well as interactions between microorganisms and plants, affect how well plants absorb and use micronutrients. Numerous studies have been carried out to investigate the soil's micronutrient absorption and bioavailability. In order to improve crop utilisation of soil nutrients, the discussion also included an analysis of the relationships between plant functions and soil nutrients.

*Keywords: Soils; micronutrients; plants; grains; ecosystem; malnutrition.*

## 1. INTRODUCTION

Crop production and quality have decreased throughout the years as a result of inadequate management of soil resources, which includes intensive crop cultivation without sufficient nutrient replenishment, limited crop rotations, and little to no addition of organic matter. Globally, there has been a rise in micronutrient deficiencies in soils, crops, animals, and humans [1,2]. The evidence that is currently available is adequate to establish the role that micronutrients play in the growth and reproduction of plants, animals, and humans. Vital micronutrients for plant development include zinc (Zn), copper (Cu), iron (Fe), manganese (Mn), boron (B), molybdenum (Mo), chlorine (Cl), and nickel (Ni). Because it keeps the body from losing calcium (Ca) and magnesium (Mg), boron (B) is a trace element that is thought to be advantageous for both humans and other animals [3]. Every micronutrient has a distinct function in the metabolism of plants, animals, and humans; therefore, a shortfall in one nutrient cannot be compensated for by an excess of another.

In several states across the nation, crop productivity is being hindered by micronutrient deficiencies, particularly in Zn, Fe, and B [4]. The relatively low fertility of Indian soils is made worse by the slow development of micronutrient deficits caused by their quicker removal during agricultural growth. According to the latest estimations, out of the 263 million tonnes (Mt) of food grains produced, 188.4 thousand tonnes (Tt) of micronutrients were lost. Out of these, the following were eliminated: 23.9 Tt of Zn, 110.6 Tt

of Fe, 37.4 Tt of Cu, 63.3 Tt of Mn, 9.2 Tt of B, and 0.99 Tt of Mo. Zn, Fe, Mn, Cu, and B deficiencies were found in 36.5%, 12.8%, 7.1, 4.2, and 23.4% of the over 2 lakh soil samples examined in order to evaluate the micronutrient condition of the soil [5]. Micronutrient deficiencies in crops and soils have a detrimental effect on yield, animal and human health, and associated variables [6]. Micronutrient deficiencies in animals lead to a range of physiological effects because these nutrients are needed for metabolic processes that affect growth, reproduction, and overall health. Although subclinical micronutrient insufficiency may not impact growth and feed efficiency, it can impact immunity and reproduction. Agroclimatic circumstances and increasing population pressure have forced the world system to prioritise food security and safety over nutrition by increasing agricultural productivity [7-9]. Recent years have seen an increase in the prevalence of micronutrient deficiencies due to the introduction of contemporary, high-yielding crop varieties, increased soil erosion, and unbalanced fertilizer use. Significant research is required to completely comprehend micronutrients, their agroecological complexity in the system, and the best strategies to use them. Crops must be biofortified using various techniques to boost their micronutrient content [10]. Future crops will significantly degrade due to the ongoing rise in micronutrient deficiencies. Given the significance of micronutrients in plant sciences, agronomic solutions for enhancing micronutrient use efficiency in crops are required to ensure the nutritional security of food production.

**Table 1. Roles of micronutrients viz-a-viz plants (Bhatt et al. [24])**

<b>Elements</b>	<b>Taken Form</b>	<b>Roles / Functional Contributions</b>
Iron	Fe <sup>2+</sup> / Fe <sup>3+</sup>	<ul style="list-style-type: none"> <li>❖ Promotes formation of chlorophyll</li> <li>❖ Enzyme mechanism which operates the respiratory system of cells</li> <li>❖ Reactions involving cell division and growth</li> </ul>
Manganese	Mn <sup>2+</sup>	<ul style="list-style-type: none"> <li>❖ Predominant in metabolism of organic acids</li> <li>❖ Activates the reduction of nitrite and hydroxylamine to ammonia</li> <li>❖ Role in important enzymes involved in respiration and enzyme synthesis</li> <li>❖ Activator of enzyme reactions such as oxidation / reduction, hydrolysis</li> <li>❖ Direct influence on sunlight conversion in the chloroplast</li> </ul>
Zinc	Zn <sup>2+</sup>	<ul style="list-style-type: none"> <li>❖ Formation of growth hormones (Auxin)</li> <li>❖ Seed and grain formation</li> <li>❖ Promotes maturity</li> <li>❖ Plant height</li> <li>❖ Protein synthesis</li> <li>❖ Transformation and consumption of carbohydrate</li> </ul>
Boron	B[OH] <sub>3</sub> Or as H <sub>2</sub> BO <sub>3</sub> (Depends on pH)	<ul style="list-style-type: none"> <li>❖ Protein synthesis</li> <li>❖ Formation of plant hormones</li> <li>❖ Promotes maturity</li> <li>❖ Increase in flowering set</li> <li>❖ Affects nitrogen and carbohydrate metabolism</li> <li>❖ Water relation in plant</li> </ul>
Copper	Cu <sup>2+</sup>	<ul style="list-style-type: none"> <li>❖ Enzyme activator</li> <li>❖ Major function in photosynthesis and reproductive stage</li> <li>❖ Indirect role in chlorophyll production</li> <li>❖ Increase sugar content</li> <li>❖ Function in respiratory system</li> <li>❖ Intensifies colour</li> <li>❖ Improves flavour in fruits and vegetables</li> </ul>
Molybdenum	MoO <sub>4</sub>	<ul style="list-style-type: none"> <li>❖ Essential for nitrogen fixation, nitrate reduction and plant growth</li> </ul>
Chlorine	Cl <sup>-</sup>	<ul style="list-style-type: none"> <li>❖ Activator for the enzyme that releases oxygen from water during photosynthesis</li> <li>❖ Regulating turgor pressure and growth of cells and is important in drought resistance</li> <li>❖ Counter ion to the positively charged “cations” in the cells</li> </ul>

*\*Ni will not be reviewed since it has minor importance in agricultural practices*

## 2. MICRONUTRIENTS viz-a-viz ECOSYSTEM

In order to use micronutrients in agricultural soil, they are frequently extracted from the earth's crust. Add mined minerals to agricultural soils to compensate for deficiencies in micronutrients. The weathering and decomposition of soil minerals and organic components produce the majority of micronutrients. There are complexes

in the fluid that attach to organic matter and clay as well as free, soluble nutrient ions. Micronutrient availability is influenced by a number of soil parameters, including ion interaction, redox potential, organic matter, acidity, and alkalinity. Soils with low fertility can receive micronutrients from animal feed additives and mined minerals. Micronutrients and organic amendments can increase crop yields and water efficiency when there is a water shortage [11].




Fig. 1. Shows the essential macro and micro-nutrients [both minerals and non-minerals] for plant growth [25]




Fig. 2. Micronutrients are essential in all stages of a plant's growth and development (<https://imma.co.in/micronutrients-for-plants/>)

Plant availability and Zn solubility are controlled by the adsorption-desorption process [12]. According to Hussain et al. [13], Zn fertiliser

increased grain absorption and mobilisation. Agronomic biofortification reduces the quantity and quality of agricultural productivity, which is

impacted by a lack of soil micronutrients. Mineral shortages result from insufficient soil replenishment of parent material and adsorbed complexed fractions. Nutrient deficiencies in agricultural soils were caused by natural processes, soil pH, human activities, and intensive farming without fertilizer [14]. Although micronutrient deficiencies in soil are common, their severity varies by geography and nutrient [15]. Micronutrient deficits are common in crops such as grains, oilseeds, pulses, and vegetables that are cultivated extensively. The micronutrient deficiencies of zinc (Zn) 40%, iron (Fe) 12.6%, copper (Cu) 4.5%, manganese (Mn) 6.0%, and boron (B) 22.8% in soils have been reported across the country. The manganese deficiency is emerging extremely fast, particularly in wheat crops grown after rice in Haryana (12%) and Punjab (18%) due to leaching of Mn from the upper surface of the coarse-textured soils [16]. Application to the soil and the leaves is advised for crops lacking Zn, B, and Mo; for Fe and Mn, only application to the leaves is advised [16].

### 3. MICRONUTRIENT *viz-a-viz.* PLANT SYSTEM

The average amounts of chlorine, iron, manganese, boron, zinc, copper, molybdenum, and nickel needed per kilogramme of dry matter are 100, 50, 20, 20, 6, 0.1, and 0.1 mg, respectively. When vitamin concentrations fall below critical levels or when they become malnourished, plants exhibit symptoms of deficiencies. Deficits in micronutrients affect the physiological and biochemical functions of plants. Understanding each micronutrient's function within the plant system is necessary to comprehend the mechanisms governing their availability in soil. The majority of soil micronutrient cations are divalent (trivalent iron is possible), and negatively charged humus and clay effectively absorb them. The life cycle, environment, and genetics of the plant determine its nutritional ratio. Through a beneficial interaction, micronutrients increase the nutritional content of crops [17].

Specific nutrients are absorbed by plants. As a result, their nutritional ratios are different from the soil's. Without micronutrients, the metabolism of other macro and micronutrients by plants would not function as efficiently. To comprehend plant micronutrient transport, precise membrane micronutrient flux measurements are required. Compared to macronutrients, these elements have lower internal demands. It is important to

comprehend how minerals are absorbed by plant roots in their available forms, as well as the limits of their supply and phytoavailability. Plant mineral element accumulation is determined by the supply of mineral elements and phytoavailability [18].

### 4. MICRONUTRIENT ACCESSIBILITY DECIDING FACTORS

Soil pH affects micronutrient availability to plants as well as their ionic form, mobility, and solubility [19]. A decrease in micronutrient availability, i.e., Zn, Fe, Cu, Mn, B, etc., and Mo become more available when soil pH rises. These micronutrients are often adsorbed onto soil surfaces as sesquioxide. Sharma et al. [20] report that for every unit of soil pH increase, the solubility of Mn, Cu, and Zn decreases by approximately 100 and 1000 times, respectively. Soil pH has a direct effect on zinc mobility and availability; when pH rises, zinc availability falls [21]. Zn adsorption as hydrous oxides of Fe, Al, and Mn happens when soil pH rises over 5.5 (Moraghan and Mascagni 1991). Zn content in soil solution is increased by OM solubilization; nevertheless, Zn (OH)<sup>+</sup> develops in the soil at pH values over 7 (Barber 1995). Zn concentration in acidic soils decreases thirty times for every unit increase in soil pH between 5.0 and 7.0, according to McCride and Blasiak (1979).

Soil Organic Matter (SOM) also effected the micronutrient availability. Compounds that are categorised as either water-soluble or water-insoluble make up soil organic matter. Fulvic acids are soluble in water and have a comparatively large molecular weight to humic acids. On the other hand, humic compounds, or humic acids, are insoluble in water because they include anionic oxygen groups like phenolic hydroxyl, alcoholic hydroxyl, carboxyl, and aliphatic carboxyl [22]. It has been shown that humic acids and metals can undergo ionic bonding or complexation reactions [23]. Organic acids of low molecular weight, such malic, citric, and acetic acids, have strong ionic interactions or metal complexes. Because organic matter forms soluble complexes with organic molecules like fulvic acids or amino acids, it increases zinc availability in soil [24,25]. The formation of insoluble zinc-organic complexes with soil organic matter (SOM) affects zinc's solubility. Exudation from roots and mineralizing bacteria promote complex formation in the rhizosphere, which increases Zn availability to plants [26].

## 5. MICRONUTRIENTS IMPROVEMENT METHODS AND MITIGATE FACTORS AFFECTING MICRO-NUTRIENTS ACCESSIBILITY

Micronutrient improvement approaches include soil amendments, foliar treatments, and the use of micronutrient-efficient crop cultivars. To reduce obstacles impacting micronutrient accessibility, major initiatives include:

- ❖ **Soil pH Management:** Adjusting soil pH to the right amount can improve micronutrient availability. For example, acidifying alkaline soils can improve the availability of iron and manganese.
- ❖ **Organic Matter Addition:** Incorporating organic matter, such as compost or manure, can improve soil structure while also increasing micronutrient availability by chelating and limiting leaching.
- ❖ **Balanced Fertilization:** Using balanced fertilization procedures ensures that the presence of one nutrient does not interfere with the uptake of others, resulting in overall nutritional balance.
- ❖ **Crop Rotation and Diversity:** Crop rotation and the use of different crop species can help to promote soil health and micronutrient cycling.
- ❖ **Use of Chelates:** Applying chelated micronutrients can improve their stability and availability to plants.
- ❖ **Irrigation Management:** Proper irrigation procedures can help keep soil moisture levels stable, which is essential for micronutrient uptake.

## 6. CONCLUSIONS

Micronutrient deficits in sustainable agriculture have received a lot of attention lately, particularly when intensive cropping techniques are used. Numerous factors contribute to this significant issue, including the high demand for nutrients, nutrient loss via leaching and topsoil erosion, the use of acid soil additives, the scarcity of farmyard manure, fertiliser impurities, and the farming of marginal lands. The pH levels, organic matter content, and microbial activity all influence the availability of minerals to plants. The range of their application rates is 0.2 kg ha<sup>-1</sup> to 100 kg ha<sup>-1</sup>, contingent upon the soil type, crop requirements, and method of application. Micronutrients are typically better absorbed from the soil than from foliar sprays, despite the fact that they are required in lesser quantities and are

frequently added to fertilisers containing macronutrients. It is crucial to cultivate genotypes that use micronutrients more effectively if farming is to produce more in the future. The specificity of this review is on their critical function in plant growth and development, interactions with soil characteristics, and bioavailability as impacted by environmental variables, all of which contribute to plant health and production.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Bhatt R, Kukal SS, Busari MA, Arora S, Yadav M. Sustainability issues on rice–wheat cropping system. *International Soil and Water Conservation Research*. 2016; 4(1):64-74.
2. Bhatt R, Singh P, Hossain A, Timsina J. Rice–wheat system in the northwest Indo-Gangetic plains of South Asia: issues and technological interventions for increasing productivity and sustainability. *Paddy and Water Environment*. 2021;19(3):345-365. DOI: 10.1007/s10333-021-00846-7
3. Dhaliwal SS, Sharma V, Shukla AK. Impact of micronutrients in mitigation of abiotic stresses in soils and plants—A progressive step toward crop security and nutritional quality. *Advances in Agronomy*. 2022;173:1-78. Available: <https://doi.org/10.1016/bs.agron.2022.02.001>
4. Kumar D, Patel KC, Ramani VP, Shukla AK, Behera SK, Patel RA. influence of different rates and frequencies of Zn application to maize–wheat cropping on crop productivity and Zn use efficiency. *Sustainability*. 2022;14(22): 15091. Available: <https://doi.org/10.3390/su142215091>

5. Shukla AK, Behera SK. Micronutrients research in India: Retrospect and prospects. In Preprint, FAI Annual Seminar. The Fertiliser Association of India, New Delhi. 2017;10:SII-4/1-SII-4/17
6. Shukla AK. Understanding the mechanism of variation in status of a few nutritionally important micronutrients in some important food crops and the mechanism of micronutrient enrichment in plant parts, NAIP Funded Research Project. AICRP on Micronutrients, IISS, Nabibagh, Berasia Road, Bhopal. 2014;1-97.
7. Afrin S, Islam N, Mustaki S, Araf T, Choudhury S. Impact of Micronutrients and Plant Growth Regulators on Brinjal (*Solanum melongena* L.) Growth, Yield and Quality. Asian Journal of Soil Science and Plant Nutrition. 2024;10(2):72–79. Available:<https://doi.org/10.9734/ajsspn/2024/v10i2262>
8. Jayakumar S, Abhangrao AK, Sarje RA, Gupta R, Pathania S, Sree BV. Critical Analysis on Effect of Micronutrients on Flowering Plants: A Review. International Journal of Plant & Soil Science. 2024;36(6):776–782. Available:<https://doi.org/10.9734/ijpss/2024/v36i64683>
9. Fageria NK, Baligar VC, Clark RB. Micronutrients in crop production. Advances in agronomy. 2002;77: 185-268.
10. Sandhu A, Dhaliwal SS, Shukla AK, Sharma V, Singh R. Fodder quality improvement and enrichment of oats with Cu through biofortification: A technique to reduce animal malnutrition. Journal of Plant Nutrition. 2020;43(10): 1378-1389.
11. Molden D, Oweis T, Steduto P, Bindraban P, Hanjra MA, Kijne J. Improving agricultural water productivity: Between optimism and caution. Agricultural water management. 2010;97(4):528-535.
12. Alloway BJ. 'Zn in soils and crop nutrition' IZA Publications. International fertilizer Industry Association, Brussels. 2004:1–116.
13. Husain FM, Ahmad I, Baig MH, Khan MS, Khan MS, Hassan I, Al-Shabib NA. Broad-spectrum inhibition of AHL-regulated virulence factors and biofilms by sub-inhibitory concentrations of ceftazidime. RSC advances. 2016;6(33): 27952-27962.
14. Dhaliwal SS, Dhaliwal MK, Shukla AK, Manchanda JS. Long-term effect of integrated nutrient management on distribution of DTPA-extractable micronutrients in Typic Haplustept under rice-wheat system. Indian Journal of Fertilisers. 2017;13(8):54-57.
15. Voortman RL, Bindraban PS. Beyond N and P: towards a land resource ecology perspective and impactful fertilizer interventions in Sub-Sahara Africa. VFRC Report 2015/1. Virtual Fertilizer Research Centre, Washington, DC, USA. 2015:49
16. Kumar D, Patel KP, Ramani VP, Shukla AK, Meena RS. Management of micronutrients in soil for the nutritional security. Nutrient Dynamics for Sustainable Crop Production. 2020;103-134.
17. Dimkpa CO, Hansen T, Stewart J, McLean JE, Britt DW, Anderson AJ. ZnO nanoparticles and root colonization by a beneficial pseudomonad influence essential metal responses in bean (*Phaseolus vulgaris*). Nanotoxicology. 2015;9(3):271-278.
18. Ahmed N, Zhang B, Chachar Z, Li J, Xiao G, Wang Q, Tu P. Micronutrients and their effects on horticultural crop quality, productivity and sustainability. Scientia Horticulturae. 2024;323:112512.
19. Fageria NK, Baligar VC, Jones CA. Growth and Mineral Nutrition of Field Crops. 2nd edition. Dekker, New York; 1997.
20. Sharma V, Kanwar BB, Verma TS. Iron status in pea growing soils of dry temperate Zone of Himachal Pradesh. Journal of Soils and Crops. 2007; 17(1):7-13.
21. Saeed M, Fox RL. Relations between suspension ph and zinc solubility in acid and calcareous soils. Soil Science. 1977; 124(4):199-204.
22. Tate III RL. Soil Organic Matter: Biological and Ecological Effects. Wiley, New York; 1987.
23. Stevenson FJ. Cycles of Soil Carbon, Nitrogen, Phosphorous, Sulfur, and Micronutrients. Wiley, New York; 1986.
24. Bhat BA, Islam ST, Ali A, Sheikh BA, Tariq L, Islam SU, Hassan Dar TU. Role of micronutrients in secondary metabolism of plants. Plant Micronutrients: Deficiency and Toxicity Management. 2020;311-329.
25. Kerry RG, Mahapatra GP, Patra S, Sahoo SL, Pradhan C, Padhi BK, Rout JR.

Proteomic and genomic responses of plants to nutritional stress. *Biometals*. 2018;31:161-187. 26. Lindsay WL. Zinc in soils and plant nutrition. *Adv. Agron.*, T. 1972;24:147.

---

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/118164>