



Impact of Molybdenum Foliar Application on Black Bean Development and Seed Production

Fabio da Silva Ribeiro ^a
and Maximiliano Kawahata Pagliarini ^{b*}

^a Grand Dourados University Center, Brazil.

^b Experimental Farm, Grand Dourados Federal University, Brazil.

Authors' contributions

This work was carried out in collaboration between both authors. Author FSR was the fellow responsible for executing the project, writing the first version in Portuguese, and collecting data. Author MKP was the project coordinator and advisor to the fellow. He conceived the project, wrote the research project, assisted in collecting data in the field, finalized the version of the article in Portuguese, and translated it into English. Both authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i111719>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/127255>

Original Research Article

Received: 18/09/2024

Accepted: 22/11/2024

Published: 26/11/2024

ABSTRACT

Bean is a crop of great importance to Brazilian agriculture, due to their relevance for population diet and also as it is one of the countries that produces and consumes more beans. However, their production remains low and one of the reasons may be the failure in absorbing the most of the fertilization applied. Therefore, research is needed to find ways to ensure maximum efficiency and absorption of nutrients by plants, such as the use of Molybdenum application via foliar. The objective of the work was to evaluate the development and yield of beans with foliar application of Molybdenum. The experiment was conducted in randomized blocks in a 2 x 4 x 3 factorial scheme

*Corresponding author: E-mail: mpagliarini@ufgd.edu.br;

Cite as: Ribeiro, Fabio da Silva, and Maximiliano Kawahata Pagliarini. 2024. "Impact of Molybdenum Foliar Application on Black Bean Development and Seed Production". *Journal of Advances in Biology & Biotechnology* 27 (11):1322-29. <https://doi.org/10.9734/jabb/2024/v27i111719>.

(sowing season x doses x application time), with treatments consisting of two sowing season: autumn/winter (sowing in February - dry season) and spring/summer (sowing in October – rainy season), four Mo doses (0, 50, 150 and 300 ml ha⁻¹ of commercial Mo based product) and three application times, starting the count after 50% plant emergence (10 days after emergence – DAE, 20 DAE and 30 DAE). The experiment was performed at the Experimental Farm of the Grand Dourados Federal University, at the municipality of Dourados, MS, Brazil. The Mo source was a commercial product with a concentration of 216 g l⁻¹ Mo. Black beans were sown using a conventional procedure, with soil preparation by subsoiling, harrowing and leveling. A pneumatic seeder was used with a spacing of 0.45 m, 12 seeds per meter of furrow, in plots of 5 x 5 m, totaling 10 plants per meter, a population of 222,222 plants ha⁻¹ and depth of 0.02 m. For the agronomic evaluations, five plants per plot were collected at the end, when the crop reached physiological maturity, in the central line, for the following determinations: number of pods per plant; number of seeds per plant; number of seeds per pod; mass of a thousand seeds and yield. Positive Results in yield components and effective yield of black-beans were found when 50 ml ha⁻¹ of molybdenum was applied via foliar application. Although it was not one of the objectives of the study, it was concluded that the amount of rainfall in the crop cycle affects the action of molybdenum on bean plants.

Keywords: Sowing season; bean yield; foliar fertilizer.

1. INTRODUCTION

Global agricultural production has been growing exponentially and this is due to research that guarantees greater yield in less time as well as the emergence of new technologies (Kuzma et al., 2023).

Brazil occupies a prominent place in global agricultural production as its unique soil and climate characteristics guarantee year-round production of various plant species.

Among them, beans, a species belonging to the Fabaceae family, stand out due to their great importance for Brazilian agriculture, as the relevance for population eating habits and for being one of the countries that produces and consumes more (Oliveira et al., 2023).

Despite this, bean production has not varied much over time and remains stagnant, as shown by data published by the Brazilian Supply Company (CONAB), in which the average production in season 2011/2012 harvest was 890 kg ha⁻¹ and in 2023/2024 harvest it was 1,141 kg ha⁻¹, an increment of just 211 kg in twelve years (Conab, 2024).

Low crop yield occurs, among other factors, due to inadequate mineral nutrition based on chemical fertilization, commonly by nitrogen, phosphorus and potassium.

This type of fertilization is advantageous by rapid availability of nutrients for plants, since they are

soluble sources. However, this high solubility may bring disadvantages, as its application must be done annually, and therefore, what is not assimilated by the crops may be lost through leaching and may contaminate surface and groundwater.

Therefore, it is necessary that different management methods be tested in order to guarantee maximum efficiency and absorption of these nutrients by plants, such as the use of molybdenum (Mo) application via foliar (Reddy et al., 2020).

The importance of Mo for the biological nitrogen fixation (BNF) process was first described by Bortels (Bortels, 1930), who demonstrated that *Azotobacter vinelandii*, when inoculated in a culture medium without N, required Mo to grow, a fact that did not occur if the source of N in the culture medium was ammonium, that is, in a medium where N was present, the bacteria did not need Mo to synthesize assimilates (Dhaliwal et al., 2022).

In soil, in general, the amount of Mo is in the range of 0.5 to 5.0 mg kg⁻¹, occurring in soluble form at soil solution, adsorbed in colloidal fraction, retained in the crystalline network of primary minerals and chelated to organic matter (Marcondes, 2001).

The Mo participates in two factors of nitrogen metabolism that are of utmost importance for plants of the Fabaceae family: it participates in nitrate reductase (NO⁻³), in which this element is

reduced to nitrite (NO^-2), as well as nitrogenase, a process in which nitrogen-fixing bacteria convert dinitrogen gas (N_2) into ammonia (NH_3) (Epstein and Bloom, 2006), Mo synthesizes enzymes for the fixation of atmospheric nitrogen.

As with other nutrients, soil pH may also influence the availability of Mo for plants, being low at acidic pH (5.0 to 5.5) and high at pH levels greater than 7.0 (Abreu et al., 2007).

Taking into account the importance of Mo in BNF and all the possibilities of this micronutrient not being available to plants, it may be seen that at region where the experiment was conducted, the Red Latosol presents in majority of agricultural areas is rich in iron and/or aluminum oxides with low pH (4.0 to 5.5) witch means ideal conditions for unavailability of Mo, thus being necessary to add at system by seed treatment and/or via foliar application (Mahadule and Sale, 2018; Khan et al., 2024; Karthikeyan et al., 2020).

Thus, the objective of the work was to evaluate the development and yield of beans with foliar application of Molybdenum.

2. MATERIALS AND METHODS

The experiment was performed at the Experimental Farm of the Grand Dourados

Federal University, at the municipality of Dourados, MS, Brazil. The experimental location is located at latitude $22^\circ14'08''$ S, longitude $54^\circ59'13''$ W and altitude of 434 m. The climate, according to the Koppen classification, is Cwa and the soil is a Dystroferric Red Latosol (Embrapa, 2013). Average temperature and precipitation data during the experiment period are described in Fig. 1.

The experiment was conducted in randomized blocks in a $2 \times 4 \times 3$ factorial scheme (sowing season \times doses \times application time), with treatments consisting of two sowing times: autumn/winter (sowing in February - dry season) and spring/summer (sowing in October - rainy season), four Mo doses (0, 50, 150 and 300 ml ha^{-1} of commercial Mo based product) and three application times, starting the count after 50% plant emergence (10 days after emergence - DAE, 20 DAE and 30 DAE).

The Mo source was a commercial product with a concentration of 216 g L^{-1} Mo. Black beans were sown using a conventional procedure, with soil preparation by subsoiling, harrowing and leveling. A pneumatic seeder was used with a spacing of 0.45 m, 12 seeds per meter of furrow, in plots of 5 x 5 m, totaling 10 plants per meter, a population of 222,222 plants ha^{-1} and depth of 0.02 m.




Fig. 1. Monthly average temperature and precipitation accumulation during the growing seasons; A: 2022 dry season; B: 2022 rainy season; C: 2023 dry season

Fertilization was performed in the seeding furrow in all plots, regardless of treatment with 250 kg ha⁻¹ of 08-28-16 formulation. For topdressing, 60 kg of N ha⁻¹ was applied (urea will be used as N source). Molybdenum was applied using a CO₂ backpack sprayer with a “fan” spray tip, applying a spray volume equivalent to 200 L ha⁻¹. Molybdenum application times were repeated each season for comparison.

During the crop cycle, phytotechnical practices were carried out according to necessity, such as: control of weeds, diseases and insect pests. For this, specific products were used for each case, aiming to obtain the best crop development.

For the agronomic evaluations, five plants per plot were collected at the end, when the crop reached physiological maturity, in the central line, for the following determinations: number of pods per plant: all pods with seeds were selected and divided by the number of plants evaluated; number of seeds per plant: the pods obtained in the previous evaluation were manually threshed and counted. The value obtained was divided by the number of plants evaluated to obtain the number of seeds per plant; number of seeds per pod: was calculated by dividing the number of seeds per plant by the number of pods per plant; mass of a thousand seeds: 8 samples of 100 seeds per plot were selected and weighed on a precision scale (0.01 g) and yield (kg ha⁻¹): two central lines of 1 m in length were harvested, from which the pods were removed, threshed, and these grains were weighed, extrapolating in kilograms per hectare. Both the mass of a

thousand seeds and yield had their moisture corrected to 13%.

The results were subjected to analysis of variance using the F test (P<0.05) and the means were compared using the Tukey test, at 5% probability. The molybdenum doses and application times were analyzed using polynomial regression.

3. RESULTS AND DISCUSSION

The statistical analysis showed significant effects for evaluated characteristics, both for Molybdenum doses, application times and seasons.

Regarding the number of seeds per pod (Fig. 2), it was observed that there was a significant difference, only for 2022 dry season, for Mo application times, with a quadratic trend for regression analysis in which at 10 days after emergence (DAE) an average of 4.7 seeds per pod was observed; there was a reduction at 20 DAE and a subsequent increase at 30 DAE.

Oliveira et al. (2017), when working with beans, obtained a greater number of pods and consequently greater yield with the application of molybdenum. According to Kubota et al. (2008), the effect generated by Mo doses is due to the fact that this element is a constituent of the nitrogenase enzymes, responsible for the conversion of atmospheric nitrogen into ammonia, and nitrate reductase, which acts in the assimilation of N, essentially in the reduction of nitrate to nitrite.




Fig. 2. Average number of seeds per black-bean pod with foliar application of Molybdenum at different application times at 2022 dry season

The justification for the increase in N accumulation in the aerial part may be justified by the increase in the efficiency of biological nitrogen fixation (BNF) due to the application of Mo. Epstein and Bloom (Epstein and Bloom, 2006) mention that N is the nutrient that is directly related to the increase in plant biomass production, as observed by Souza et al. (2008), for soybean and Xavier et al. (2008) for cowpea.

Therefore, the accumulation of biomass that generates a greater number of pods is directly related to the greater production of grains per pod, which will result in greater production due to the greater allocation of nitrogen for grain production. As nitrogen is absorbed in greater quantities by bean plants, approximately 50% of the total N absorbed is exported to seeds (2006).

The weight of a thousand seeds (WTS) showed significant results with the application of Mo in the three sowing seasons. In the 2022 dry season (A) at a dose of 0 mL ha⁻¹ (control), the WTS was 143.43 g and increased to 145.72 g at a dose of 50 mL ha⁻¹ and subsequently there was reduction in mass (Fig. 3A).

In Fig. 3B and 3C, the regression curves showed similarities with an increase in mass gain at a dose of 150 mL ha⁻¹ and decline with higher doses, demonstrating that even in different harvests, the application of Mo provides mass accumulation up to a certain applied concentration.

As the rainfall levels and average temperature of each harvest were different (Fig. 1), it was possible to observe that in the dry harvest (C) with less precipitation distributed between February to May, the masses were greater compared to the others.

Some authors state that even in small concentrations, the presence of Mo in this legume promotes better expressivity of agronomic variables (1986).

The process of acetylene reduction and N remobilization at the time of grain development occurs due to the participation of Mo, which results in greater mass accumulation in grains (1998). The dose of 150 mL ha⁻¹ promoted greater WTS due to NBF in which the plant obtained greater use of atmospheric N.

Therefore, smaller masses was observed in the 2022 dry harvest and even smaller in the 2022 rainy harvest (Fig. 3A and 3B) due to the heavy rainfall in the period close to harvest (Fig. 1), which is the time of physiological maturation. The excess water in this period may impair the maturation, harvesting and pods drying, significantly affecting the grain yield. According to Conceição et al. (2018), excess water in the maturation period may generate intense enzymatic degradation and consequently reduction in grain mass.




Fig. 3. Weight of a thousand seeds (WTS - g) of black-beans with Molybdenum via foliar application at different application times. A: 2022 dry season; B: 2022 rainy season; C: 2023 dry season




Fig. 4. Black-bean yield (kg ha⁻¹) at different doses of Molybdenum via foliar application. A: 2022 dry season; B: 2022 rainy season; C: 2023 dry season

Regarding the yield (Fig. 4), the molybdenum doses showed significant results in the three sowing seasons, demonstrating that there was nonconformity between harvests due to climatic factors. Therefore, the effects of the doses were also different in the harvests, as in the two dry harvests the highest averages were 1,150.29 kg ha⁻¹ (A) and 613.89 kg ha⁻¹ (C) at control, however, compared to the rainy harvest (B) they were 918.88 kg ha⁻¹ at the dose of 150 mL ha⁻¹, thus, the effects of the doses were greater in crops with better rainfall distribution.

According to the precipitation data (Fig. 1), it may be seen that the availability of water during the cycle did affect yield between harvests, but it was also what most influenced the results of the doses (Fig. 4), demonstrating that in the harvest that had better rainfall distribution, molybdenum was more active in the plants, generating a difference in grain production.

This may have possibly occurred due to N stress caused by the reduced amount of water in the soil, making it difficult to absorb the nutrient made available in fertilization, and because there is no synchronization between the period of nitrogen depletion in the cotyledons and the beginning of N₂ fixation, as in bean plants this

period is around 15 to 20 days after emergence (Barradas and Hungria, 1989).

Due to the delay in N absorption, it can directly affect the flowering and grain filling period, as the demand for this nutrient is very high (Portes, 1996). According to Rosolem (1987), good plant nutrition should be carried out at a time that can increase the number of pods per plant, which would be until the beginning of flowering.

As with the weight of a thousand seeds (Fig. 3) and the number of seed per pods (Fig. 2), the accumulation of mass is directly related to yield, making plants have greater availability during the reproductive period and grain formation, as 60% of the accumulation of N during the bean cycle is absorbed and metabolized between the flowering and grain filling stages (Hungria et al., 1985).

In relation to the average yield of the Central West region of Brazil, the three harvests were lower, considering that they were cultivated in the dry harvests of 2022 and 2023; and in the rainy harvest of 2022, where the region's productions were 992.00 kg ha⁻¹, 1,239.00 kg ha⁻¹, 2,263.00 kg ha⁻¹ respectively (Conab, 2024).

4. CONCLUSION

Positive results in yield components and effective yield of black beans were found when 50 ml ha⁻¹ of molybdenum was applied via foliar application.

Although it was not one of the objectives of the study, it was concluded that the amount of rainfall in the crop cycle affects the action of molybdenum on bean plants.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

ACKNOWLEDGEMENTS

The authors would like to thank the Foundation for Support to Teaching, Research and Extension (FUNAEP) for managing the resources obtained to develop the research (Contract 13/2021 - Process No. 23005.003601/2020-81). They would also like to thank the outsourced employees who were essential for the implementation and conduct of the experiments.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Abreu, C. A., Lopes, A. S., & Santos, G. C. G. (2007). Micronutrients. In R. F. Novais, V. V. H. Alvarez, N. F. Barros, R. L. F. Fontes, R. B. Cantarutti, & J. C. Neves (Eds.), *Soil fertility* (pp. 333–342). Viçosa: Brazilian Society of Soil Science.
- Barradas, C. A., & Hungria, M. (1989). Selection of Rhizobium strains for common bean. I. Precocity for nodulation and nitrogen fixation. *Turrialba*, 39(2), 236–242.
- Bortels, H. (1930). Molybdenum as a catalyst for biological stick binding. *Archives of Microbiology*, 1, 333–342.
- Companhia Nacional De Abastecimento. (2024). *Harvest 2023/24: Twelfth survey August 2024* (Vol. 6, No. 12). Retrieved August 20, 2024, from <https://www.conab.gov.br/info-agro/safras/graos>
- Conceição, C. G., Robaina, A. D., Peiter, M. X., Parizi, A. C., Conceição, J. A., & Bruning, J. (2018). Economically optimal water depth and grain yield of common bean subjected to different irrigation depths. *Brazilian Journal of Agricultural and Environmental Engineering*, 22(7), 482–487.
- Dhaliwal, S., Sharma, V., Shukla, A., Kaur, J., Verma, V., Kaur, M., ... & Hossain, A. (2022). Interactive effects of molybdenum, zinc, and iron on the grain yield, quality, and nodulation of cowpea (*Vigna unguiculata* (L.) Walp.) in North-Western India. *Molecules*, 27(11), 3622.
- Embrapa. (2013). *Brazilian soil classification system* (3rd ed.). Brasília: Embrapa.
- Epstein, E., & Bloom, A. J. (2006). *Mineral nutrition of plants* (2nd ed.). Londrina: Planta.
- Hungria, M., Neves, M. C. P., & Victoria, R. L. (1985). Nitrogen assimilation by common bean. II. Uptake and translocation of mineral N and fixed N₂. *Brazilian Journal of Soil Science*, 9(1), 201–209.
- Kamicker, B. J., & Brill, W. J. (1986). Identification of *Bradyrhizobium japonicum* nodule isolates from Wisconsin soybean farms. *Applied and Environmental Microbiology*, 51(3), 487–492.
- Karthikeyan, A., Vanathi, J., Babu, S., & Ravikumar, C. (2020). Studies on the effect of foliar application of organic and inorganic nutrients on the phenotypic enhancement of black gram cv. Vamban-6. *Plant Archives*, 20(2), 1161–1164.
- Khan, A. Q., Khan, A. Q., Ullah, H., Iqbal, N., Junaid, M., Murad, I., ... & Ahmed, U. (2024). Maximizing growth and yield of black seed (*Nigella sativa* L.) through optimized phosphorus and boron levels. *Journal of Bioresource Management*, 11(3), 17.
- Kubota, F. Y., Andrade Neto, A. C., Araújo, A. P., & Teixeira, M. G. (2008). Growth and nitrogen accumulation of common bean plants from seeds with high molybdenum content. *Brazilian Journal of Soil Science*, 32, 1635–1641.
- Kuzma, S., Bierkens, M. F. P., Lakshman, S., Luo, T., Saccoccia, L., Sutanudjaja, E. H., & Beek, R. V. (2023). *Aqueduct 4.0: Updated decision-relevant global water risk indicators*, Technical Note. World Resources Institute.
- Mahadule, P. A., & Sale, R. B. (2018). Effect of foliar sprays of boron on growth,

- yield, nutrient uptake, and quality of French bean (*Phaseolus vulgaris* L.) in Entisol: A review. *Journal of Pharmacognosy and Phytochemistry*, 7(5), 74–78.
- Marcondes, J. A. P. (2001). *Nodulation and nitrogen uptake by soybean in response to cobalt and molybdenum application* (Master's thesis, Universidade Federal do Paraná, Curitiba).
- Oliveira, C. A. B., Pelá, G. M., & Pelá, A. (2017). Inoculation with *Rhizobium tropici* and foliar fertilization with molybdenum in common bean crops. *Journal of Neotropical Agriculture*, 4(1), 43–50.
- Oliveira, G., & Wander, A. (2023). Mapping the common bean production chain in Brazil. *Political Economy of Development Journal*, 14(32), 96–132.
- Portes, T. A. (1996). Ecophysiology. In R. S. Araújo et al. (Coords.), *Common bean culture in Brazil* (pp. 101–137). Piracicaba: Potafós.
- Reddy, K., Bulle, M., Wany, A., & Gupta, K. (2020). An overview of important enzymes involved in nitrogen assimilation in plants. *Methods in Molecular Biology*, 2057, 1–13.
- Rosalem, C. A. (1987). *Nutrition and fertilization of common bean* (p. 93). Piracicaba.
- Soratto, R. P., Carvalho, M. A. C., & Arf, O. (2006). Nitrogen topdressing in common beans grown under no-tillage. *Brazilian Journal of Soil Science*, 30(2), 259–265.
- Souza, R. A., Hungria, M., Franchini, J. C., Chueire, L. M. O., Barcellos, F. G., & Campo, R. J. (2008). Minimum set of parameters for evaluating soil microbiota and biological nitrogen fixation by soybean. *Brazilian Agricultural Research*, 43(1), 83–91.
- Vieira, R. F., Cardoso, E. J. B. N., Vieira, C., & Cassini, S. T. A. (1998). Foliar application of molybdenum in common beans. I. Nitrogenase and reductase activities in a soil of high fertility. *Journal of Plant Nutrition*, 21(1), 169–180.
- Xavier, T. F., Araújo, A. S. F., Santos, V. B., & Campos, F. L. (2008). Influence of inoculation and nitrogen fertilization on nodulation and grain yield of cowpea. *Rural Science*, 38(7), 25–32.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/127255>