



Effect of Integrated Nutrient Management (INM) on Growth Parameters and Yield of Wheat (*Triticum aestivum* L.)

**Anjali Patyal^a, Chandra Shekhar^{a*}, Ravindra Sachan^b, Deepak Kumar^c,
Ankit Yadav^b and Gyanendra Kumar^d**

^a Department of Agronomy, Dolphin PG College of Science and Agriculture, Chunni Kalan Fatehgarh Sahib, Punjab-140407, India.

^b Department of Soil Science and Agricultural Chemistry, Chandra Shekhar Azad University of Agriculture & Technology, Kanpur (U.P.), 208002, India.

^c Department of Soil Science and Agricultural Chemistry, Eternal University, Baru Sahib, Himanchal Pradesh, 173101, India.

^d Department of Agronomy, AN.D.UA & T, Ayodhya (U.P.), 224229, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i2231457

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/90985>

Original Research Article

Received 16 June 2022
Accepted 20 August 2022
Published 24 August 2022

ABSTRACT

An experiment was conducted during *spring* season of 2021-22 at Agronomy Research Farm, Dolphin PG college of Science and Agriculture Chunni Kalan, Fatehgarh Sahib Punjab, India. The experiment consists of fourteen treatments were laid out in Randomized Block Design (RBD) with three replication. The obtained results revealed that the growth parameters were significantly influenced by different integrated nutrient management at successive growth stages of wheat crop except 30 days. The growth rate of the experimented plants were increased progressively with increase in duration of wheat crop. Growth rate was maximized up to 90 days and followed by a slightly increase in growth was obtained up to harvest. Among the treatment evaluated, 100% RDF +25% N through (vermicompost) + ZnSO₄ @ 25 kg ha⁻¹ proved to found better with respect to plant height (92.25 cm), dry matter accumulation (274.65 g m⁻²) and number tillers m⁻² (92.43 m⁻²) at harvest stage in respective years. Leaf area index was measured tended to maximum at 60 days.

*Corresponding author: E-mail: csnduat6425@gmail.com;

The minimum growth parameters were observed under control. Maximum grain yield (55.32 q ha⁻¹), straw yield (81.14 q ha⁻¹), biological yield (136.46 q ha⁻¹) and harvest index (40.53 %) was associated with the plot fertilized with 100% RDF +25% N through (vermicompost)+ ZnSO₄ @ 25 kg ha⁻¹

Keywords: Growth; plant height; dry matter; LAI; tillers and yield.

1. INTRODUCTION

Wheat (*Triticum aestivum* L.) had a major cereal crop cultivated in India and abroad. It belongs to Poaceae family. Wheat generally grown in spring season and it can be successful grown in the tropical and sub-tropical climates as well as in the temperate zone. Wheat can tolerate to severe cold and snow and resume growth with the setting in warm weather in spring. It could be cultivated from sea level to an altitude of 3300 meters. The most favorable climatic condition for wheat cultivation is cool and moist season during the vegetative growth period followed by dry, moderate season for the grain to mature and ripening. The optimum temperature range for ideal germination of wheat seed is 20-25^o C. Warm and dry climatic conditions were not suited for wheat during the heading and flowering stage, excessively high or low temperatures and drought were harmful to wheat growth. Cloudy weather with high humidity and low temperatures is conducive for rust attack. It requires about 25-30^oC optimum average temperature at the time of ripening. The temperature at the time of grain filling and development were very crucial for yield. Temperatures above 25^oC tend to depress grain weight. Wheat grain contains more protein in the form of gluten than other cereals crop .Wheat occupied an acreages about 215.29 million ha with production of 730.84 million metric tons with productivity of 3390 kg ha⁻¹ [1]. In India, the wheat crop were grown on 29.55 million ha⁻¹ (13.43% global area) with the production of 101.20 million tonnes (1.3% rises previous year) and productivity about 3424 kg ha⁻¹. In Uttar Pradesh wheat occupied about 9.78 million ha⁻¹ achieved rank first in production (31.99 million tons) with productivity (3269 kg ha⁻¹). Comparatively, the wheat productivity was comparatively lower than that the Punjab (5030 kg ha⁻¹) and Haryana (4410 kg ha⁻¹)

Adoption of intensive cropping system would meet the food as per the demands of increasing population, it requires high input energy, which were not only responsible for environment changes but also increased the cost of cultivation. After the green revolution, production

of crops had increased to a great extent due to the use of chemical fertilizers but their indiscriminate use had led to soil sickness, ecological hazards and depletion of other sources of energy. The recent energy crisis, high fertilizer cost and low purchasing power of the farming community have made it necessary to re-think alternatives. Under these situations, INM was a good option in food grain security as well as to maintain soil health. Integrated nutrient management (INM) was the combined use of mineral fertilizers with organic resources such as cattle manures, crop residues, urban/rural wastes, composts, green manures etc. It contributes in attaining agronomical feasible, economically viable, environmentally sound and sustainable high crop yields in cropping systems by enhancing nutrient use efficiency and soil fertility, reducing nitrogen losses due to nitrate leaching and emission of greenhouse gases. Kaur et al. [2] reported that the plant height (cm), dry matter accumulation (g), leaf area index and number of tillers (m⁻²) of wheat improved significantly by the application of FYM @10t ha⁻¹ and phosphorus @ 80 kg P₂O₅ ha⁻¹.

FYM was a great source of supplements and contributed towards build-up of organic matter in soil. Nitrogen is a crucial component for ideal working of crops. The increment in eco-friendly production of wheat can be made conceivable by far reaching selection of progressed innovations of which fertilizer administration especially that of nitrogen though organic manure could play a key part. Among the micronutrients, zinc deficiency appears to be the most widespread owing to intensive agricultural practices, use of high analysis NPK fertilizers and limited or no application of Zinc by farmers. At present about 48.1% of Indian soils are rated as very low in available zinc [3]. Hence, present investigation was carried out to study the growth, yield and nutrient uptake behaviour of wheat to define optimum dose under integrated use of FYM [4].

2. MATERIALS AND METHODS

An experiment was conducted in Effect of Integrated Nutrient Management (INM) on growth

parameters of wheat (*Triticum aestivum* L.) during *spring* season of 2021-22 at the Agronomy Research Farm, Dolphin PG college of Science and Agriculture Chunni Kalan Fatehgarh sahib Punjab, India. The experiment consists of fourteen treatments *i.e.* T₁-100% RDF (Recommended dose of fertilizer), T₂. 75% RDF, T₃. 75% RDF + ZnSO₄ @25kg ha⁻¹, T₄. 75% RDF + sulphur @40kg/ha⁻¹, T₅. 75%RDF +25% N through FYM + *Azotobacter* T₆. 75% RDF + ZnSO₄ @25kg ha⁻¹ + Sulphur @40kg ha⁻¹, T₇. 100% RDF +25% N through (vermicompost) + ZnSO₄ @25kg ha⁻¹ and Control (T₈) treatments were laid out in randomized block design (RBD) with three replications. The experimental field was well drained and leveled in order to determine the fertility status and soil class. The soil samples were collected randomly from different places of the experimental field by using of soil auger at the depth of 0-15 cm for physico-chemical analysis before execution of the fertility treatments. The soil texture was silty loam with the pH of (7.8) and EC (0.14,dS/m). The organic carbon content of soil was (0.44%) with nitrogen (189 kg ha⁻¹), phosphorous (15 kg ha⁻¹), and potassium (178 kg ha⁻¹). The experimental location was related to the sub-tropical climate of Indo-Gangatic alluvial plains zone (IGP) had alluvial calcareous soil and located at 26°47' N latitude and 82°12' E longitude with an altitude of 113 m above the mean sea level. Variety PBW 154 was shown proper moisture stage on 18 Nov, 2021-22. Seed was treated with @ 1 packet *Azotobacter* with 10 kg seed of wheat by making the pest with the help of 500 ml water and 50 g jaggery heated and cools down and mix with seed. After dry seeds in shade and use for sowing. The seeds were placed in manually opened furrows at the depth of 5 cm below with the row spacing of 20 cm apart. The seeds was applied informally in each and every plots @ 100 kg ha⁻¹. Nitrogen, phosphorus and potassium fertilizers were applied in the forms of Urea, DAP and Muriatic of potash @ 120, 60 and 40 kg ha⁻¹, respectively. Full dose of phosphorus, potassium and half dose of nitrogen were applied as basal dressing at the time of sowing and rest half dose of nitrogen was applied as two split doses at the first and second irrigation.

Leaf area index was calculated by the following equation according to Mckee, [5]

$$\text{Leaf area index} = \frac{\text{Total leaf area (cm}^2\text{)}}{\text{Total land area (cm}^2\text{)}}$$

3. RESULTS AND DISCUSSION

3.1 Growth Parameters

3.1.1 Plant height (cm)

Plant height increased progressively with increase in duration of wheat crop (Table 1). Crop growth rate was maximized up to 90 DAS followed by a slow increase in growth. The plant height was significantly influenced by the various integrated nutrient management practices. The taller plants were recorded with the application of 100% RDF +25% N through (vermicompost) + ZnSO₄ @ 25 kg ha⁻¹ (T₇), which was statistically at par with T₁, T₅, and T₆ treatments, respectively at all the successive growth stages of crop. This might be due to increment of additional nutrient applied as the form of different component of INM. Probably the increase in auxin, cell division and cell enlargement by the supply of higher level of INM brought about increased in the plant height Pandey et al. [6]. Whereas the minimum plant height was recorded in control at all the growth stages this might be due to lack of the nutrient available in soils Khattak et al. [7].

3.1.2 Dry matter accumulation

The dry matter accumulation was increased with increasing the rate of photosynthesis (Table 2). The highest dry matter accumulation was recorded under 100% RDF +25% N through (vermicompost) + ZnSO₄ @ 25 kg ha⁻¹ (T₇) treatments, which was statistically at par with T₁, T₅ and T₆ treatment, respectively. While it was significantly superior over rest of treatments at all successive growth stages of plant. The significant improvement dry matter accumulation was observed when increased the different nutrients sources. This was not only provides nutrients but also improved physical condition of soil in respect of granulation, friability and porosity in addition to increased cation exchange capacity, Phosphate and sulphate availability of soil and low nitrogen loses due to slow release of nutrient from different components of integration. Thus these mechanism improved the plant height, number of leaves, leaf area index and number of tillers which were enhanced the dry matter accumulation [8].

3.1.3 Leaf area index

Integrated nutrient management practices significantly influenced on the leaf area index at 30, 60 and 90 days in both the years

experimentation (Table 3). Among the integrated nutrient management, 100% RDF +25% N through (vermicompost) + ZnSO₄ @ 25 kg ha⁻¹ was exerted maximum leaf area index. The leaf area index gradually increased upto 90 DAS and tended to maximum value (2.36) at 90 DAS. This might be due to maximum availability of nutrient in soil cause more nutrient uptake by in plant, which increased although growth parameters *i.e.* plant height, number of leaf plant⁻¹, number of tillers m⁻² which has ultimately enhanced the leaf area index Bachhao et al. [9].

3.1.4 Number of tillers (m⁻²)

Obviously the data summarized in (Table 4) revealed that the number of tillers significantly increased with the increasing of different components of plant nutrients. The number of tillers were increased progressively up to 90 DAS and thereafter decreased. The maximum number

of tillers (m⁻²) were observed with the application of 100% RDF +25% N through (vermicompost) + ZnSO₄ @ 25 kg ha⁻¹, which was significantly higher over control. This might be due to increased the application of different components of INM in soil and improved the nutrient status of the soil as well as increased the microbial activity of soil. This indicated that, the plant get favorable condition for growth and development as well as chlorophyll content by canopy of plants caused more photosynthesis which was ultimately increased number of tillers m⁻². Similar findings was reported by Akhtar et al. [10].

3.2 Productivity Parameters

Observed that the data given in Table 5 clearly shows that among the productivity parameters viz. grain yield, straw yield, biological yield and harvest index weresignificantly affected by different treatment combinations of organic

Table 1. Plant height (cm) at subsequent growth stages of wheat as influenced by various integrated nutrient management (INM)

S. no.	Treatments	Plant height (cm)		
		30 DAS	60 DAS	90 DAS
1	100% RDF (Recommended dose of fertilizer)	28.40	51.40	88.75
2	75% RDF	28.41	49.12	88.10
3	75% RDF + ZnSO ₄ @25 kg ha ⁻¹	28.31	50.49	88.50
4	75% RDF + Sulphur @40 kg ha ⁻¹	28.13	50.15	88.10
5	75%RDF +25% N through FYM + <i>Azotobacter</i>	28.14	53.15	90.47
6	75% RDF + ZnSO ₄ @25 kg ha ⁻¹ + Sulphur @40 kg ha ⁻¹	28.32	50.46	89.12
7	100% RDF +25% N through (vermicompost)+ ZnSO ₄ @25kg ha ⁻¹	28.42	54.41	92.25
8	Control	28.50	49.00	87.15
	C.D. at 5 %	N/A	2.89	3.15
	SEm±	0.767	1.36	1.39

Table 2. Dry matter accumulation (g m⁻²) at successive growth stages of wheat crop as influenced by various integrated nutrient management (INM)

S. no.	Treatments	Dry matter accumulation (g m ⁻²)			
		30 DAS	60 DAS	90 DAS	Harvest
1	100% RDF (Recommended dose of fertilizer)	29.14	97.10	165.48	264.72
2	75% RDF	29.52	94.15	159.13	260.14
3	75% RDF + ZnSO ₄ @25 kg ha ⁻¹	29.46	96.15	164.36	262.35
4	75% RDF + Sulphur @40 kg ha ⁻¹	29.48	95.90	161.46	259.35
5	75%RDF +25% N through FYM + <i>Azotobacter</i>	29.50	100.12	168.98	267.65
6	75% RDF + ZnSO ₄ @25 kg ha ⁻¹ + Sulphur @40 kg ha ⁻¹	29.41	98.56	167.55	260.14
7	100% RDF +25% N through (vermicompost)+ ZnSO ₄ @25kg ha ⁻¹	28.41	108.35	170.63	274.65
8	Control	28.51	93.16	158.19	258.10
	C.D. at 5 %	N/A	7.98	3.64	6.56
	SEm±	0.79	2.60	2.40	3.50

Table 3. Leaf area index at successive growth stages of wheat crop as influenced by various integrated nutrient management (INM)

S. no.	Treatments	Leaf area index (LAI)		
		30 DAS	60 DAS	90 DAS
1	100% RDF (Recommended dose of fertilizer)	1.62	1.84	1.92
2	75% RDF	1.65	1.79	1.90
3	75% RDF + ZnSO ₄ @25 kg ha ⁻¹	1.63	1.80	1.92
4	75% RDF + Sulphur @40 kg ha ⁻¹	1.62	1.77	1.80
5	75%RDF +25% N through FYM + <i>Azotobacter</i>	1.64	1.87	2.10
6	75% RDF + ZnSO ₄ @25 kg ha ⁻¹ + Sulphur @40 kg ha ⁻¹	1.65	1.85	1.95
7	100% RDF +25% N through (vermicompost)+ ZnSO ₄ @25kg ha ⁻¹	1.63	1.96	2.36
8	Control	1.64	1.75	1.78
	C.D. at 5 %	N/A	0.24	0.15
	SEm ±	0.04	0.04	0.05

Table 4. Number of tillers per m⁻² of wheat as influence by various integrated nutrient management (INM)

S. no.	Treatments	Number of tillers (m ⁻²)			
		30 DAS	60 DAS	90 DAS	Harvest
1	100% RDF (Recommended dose of fertilizer)	45.10	90.15	89.14	85.13
2	75% RDF	45.12	86.42	85.12	84.13
3	75% RDF + ZnSO ₄ @25 kg ha ⁻¹	45.12	89.50	88.41	82.61
4	75% RDF + Sulphur @40 kg ha ⁻¹	45.14	82.14	82.13	81.12
5	75%RDF +25% N through FYM + <i>Azotobacter</i>	46.41	92.16	91.14	91.13
6	75% RDF + ZnSO ₄ @25 kg ha ⁻¹ + Sulphur @40 kg ha ⁻¹	46.31	90.45	90.14	89.10
7	100% RDF +25% N through (vermicompost)+ ZnSO ₄ @25kg ha ⁻¹	47.50	93.52	92.43	91.53
8	Control	41.47	85.16	84.10	83.12
	C.D. at 5 %	N/A	3.35	3.45	3.67
	SEm ±	1.21	2.36	2.33	2.31

Table 5. Effect of integrated nutrient management (INM) on productivity parameters

S. no.	Treatments	Grain yield (q ha ⁻¹)	Straw yield (q ha ⁻¹)	Biological yield (q ha ⁻¹)	Harvesting index (%)
1	100% RDF (Recommended dose of fertilizer)	49.39	77.73	127.12	38.85
2	75% RDF	45.70	77.46	123.16	37.10
3	75% RDF + ZnSO ₄ @25 kg ha ⁻¹	48.35	78.50	126.85	38.11
4	75% RDF + Sulphur @40 kg ha ⁻¹	47.85	77.65	125.50	38.12
5	75%RDF +25% N through FYM + <i>Azotobacter</i>	52.45	80.29	132.74	39.51
6	75% RDF + ZnSO ₄ @25 kg ha ⁻¹ + Sulphur @40 kg ha ⁻¹	50.65	79.20	129.85	39.00
7	100% RDF +25% N through (vermicompost)+ ZnSO ₄ @25kg ha ⁻¹	55.32	81.14	136.46	40.53
8	Control	46.10	78.65	124.75	36.95
	C.D. at 5 %	4.00	3.54	7.34	2.35
	SE m±	1.30	2.12	3.42	1.03

manure and inorganic fertilizers. Maximum grain yield (55.32 q ha⁻¹), straw yield (81.14 q ha⁻¹), biological yield (136.46 q ha⁻¹) and harvest index (40.53 %) was associated with the plot

fertilized with 100% RDF +25% N through (vermicompost)+ ZnSO₄ @ 25 kg ha⁻¹ has been statically par with the plot fertilized with 75% RDF + 25% N through FYM + *Azotobacter*. Minimum grain yield (46.10 q ha⁻¹), straw yield (78.65 q ha⁻¹), biological yield (124.75 q ha⁻¹) and harvest index (36.95 %) has been associated with the control plot. The results of the present investigation were also in agreement with the findings of Choudhary et al. [11].

4. SUMMARY AND CONCLUSION

The different component of integrated nutrient management were increased with increase the growth parameters as par duration of the wheat plants. Among the integrated nutrient management, application of 100% RDF +25% N through (vermicompost) + ZnSO₄ @ 25 kg ha⁻¹ recorded maximum growth parameters in respect of plant height, dry matter accumulation plant⁻¹, leaf area index and number of tillers at different growth stages of crop. Similarly maximum yield was also associated with the treatment fertilized with 100% RDF +25% N through (vermicompost) + ZnSO₄ @ 25 kg ha⁻¹. However, the minimum value of growth parameters and yield parameters was received under control during both the years of experimentation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anonymous. Foreign Agriculture Service/USDA office of global analysis; 2019.
2. Kaur R, Kumar S, Kaur R, Kaur J. Effect of Integrated nutrient management on yield of wheat (*Triticum aestivum* L.) under irrigated conditions. 2018;6(4):1800-1803.
3. Kumar P, Dubey SD, Sachan R, Rawat CL, Kumar V. Effect of organic manure, inorganic fertilizers and biofertilizers on nutrient content of maize (*Zea mays* L.) and their residual effect on succeeding wheat (*Triticum aestivum* L.) crop; 2022.
4. Choudhary L, Singh KN, Gangwar K, Sachan R. Effect of FYM and Inorganic fertilizers on growth performance, yield components and yield of wheat (*Triticum aestivum* L.) under indo-gangetic plain of Uttar Pradesh; 2022.
5. Mckee GW. A coefficient for computing leaf area in hybrid corn. *Agron. J.* 1964; 56(2):240-241.
6. Pandey IB, Bharti V, Bharti RC, Mishra SS. Effect of fertilizer level and seed rates on growth and yield of surface seeded wheat (*Triticum aestivum* L.) under low land rice ecosystem of North Bihar. *Indian Journal of Agronomy.* 2004;49 (1):43-45.
7. Khattak SG, Peter JD, Wiqar A. Effect of Zn as soil addition and foliar application on yield and protein content of wheat in alkaline soil. *J. Natn. Sci. Foundation Sri Lanka.* 2015;43(4):303-312.
8. Jain P. Effect of sowing methods, input and varieties levels on growth, yield components, yield and nutrient uptake of durum wheat (*Triticum durum* desf.). *Trends in Biosciences.* 2012;5:22-24.
9. Bachhao KS, Kolekar PT, Nawale SS, Kadlag AD. Response of different wheat varieties to different sowing dates. *Journal of Pharmacognosy and Phytochemistry.* 2018;7(1):2178-2180.
10. Akhtar N, Ramani VB, Yunus M, Femi V. Effect of different nutrient management treatments on growth, yield attributes, yield and quality of wheat (*Triticum aestivum* L.). *International Journal of Current Microbiology and Applied Science.* 2018; 7:3473-3479.
11. Choudhary L, Singh KN, Gangwar K, Sachan R, Pandey HP. Effect of FYM and inorganic fertilizers on nutrient content, uptake and quality traits of wheat (*Triticum aestivum* L.) under Indo-Gangetic Plain of Uttar Pradesh; 2022.

© 2022 Patyal et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/90985>