



Reference Interval of Glycated Hemoglobin in Adults in Port Harcourt, Nigeria

Allison Frederick Igila^{1*} and C. Ojule Aaron¹

¹Department of Chemical Pathology, Faculty of Basic Medical Sciences, University of Port-Harcourt, Rivers State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Author AFI designed the study, performed the statistical analysis, wrote the protocol and wrote the final draft. Author COA managed the analysis of the study and the literature review. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJMAH/2019/v16i430148

Editor(s):

(1) Prof. Janvier Gasana ,Department of Environmental & Occupational Health, EO Epidemiology, and EO Medicine, Robert Stempel College of Public Health & Social Work, Florida International University, USA.

Reviewers:

(1) Awofadeju Stephen Olajide, Obafemi Awolowo University, Nigeria.

(2) Veeravan Lekskulchai, Srinakharinwirot University, Thailand.

(3) Adebayo Olugbenga Adegoke, Madonna University, Nigeria.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/51683>

Original Research Article

Received 25 July 2019
Accepted 27 September 2019
Published 03 October 2019

ABSTRACT

Background: Measurement of glycated haemoglobin is accepted as the gold standard for the diagnosis and management of diabetes mellitus; but it is rarely used in this environment, a resource-poor setting, largely due to the high cost of the assay. There is need to determine its reference intervals in Port-Harcourt and encourage the use of this assay.

Aim: This study was designed to determine the reference interval of glycated haemoglobin of apparently healthy subjects between the ages of 20-80 years in Port Harcourt, Rivers State, Nigeria.

Methods: A total of 172 subjects who met the inclusion criteria were recruited for the study. A cross sectional sampling method was used to recruit subject. Subjects' past medical history, demographic and anthropometric information were obtained with the help of a questionnaire. Blood was collected from subjects into an EDTA bottle and analysed for glycated haemoglobin using the boronate affinity chromatographic method.

Results: Results were analyzed using Statistical Package for Social Sciences version 20.0 (SPSS 20.0) and the reference interval was determined by the nonparametric method due to the skewness

*Corresponding author: E-mail: kuufredo69@gmail.com;

of the data. The mean age of subjects was 35.4 years and the minimum and maximum glycated haemoglobin value obtained were 3.5% and 6.0% after eliminating outliers. This gave a mean glycated haemoglobin value of 4.84% and a reference interval of 4.0-5.9%.

Conclusion: The reference interval so determined (4.0-5.9%) is different from that generated by the manufacturer of the diagnostic kit(4.0-6.5%). The introduction of this new local reference interval will enhance patient management in our local hospitals.

Keywords: Glycated hemoglobin; diabetes mellitus; boronate affinity chromatographic method; obesity.

1. INTRODUCTION

With the increase in the incidence of obesity and metabolic syndrome worldwide,[1] there is associated increase in type-2 diabetes [2]. Port Harcourt, which is the capital city of Rivers state in Nigeria has one of the highest prevalence of obesity and type-2 diabetes in Nigeria.[2] With diabetes on the increase and with studies showing that a persistently high glycated hemoglobin value positively correlates with the development of diabetes complications,[3] the need for tighter control to prevent diabetic complications has made glycated hemoglobin estimation a reliable means of making diagnosis, monitoring patients and preventing complications. Plasma concentration of glycated haemoglobin is not affected by the day to day variation of plasma glucose or by the quality of patient preparation prior to sample collection[4]. This means that this test can be conducted anytime of the day without prior instruction and this has placed it at an advantage over fasting plasma glucose, but not necessarily replacing fasting plasma glucose for day to day monitoring of patient's blood glucose. To this effect, a local population based reference interval is necessary and important. This study was therefore designed to determine the reference interval of glycated hemoglobin in apparently healthy subjects in Port Harcourt, Rivers state, Nigeria and to evaluate the association of gender, hemoglobin concentration and BMI with glycated haemoglobin values in our population.

2. METHODS

The University of Port Harcourt Teaching Hospital, (UPTH) is a tertiary hospital in Rivers State. It has evolved since its inception in 1980 from a 60 bedded hospital to one with over 800 beds and is presently the apex health institution in Rivers State.

A cross sectional survey method was used to select apparently healthy subjects from the

population. Sample size was determined using Reed hypothesis, which suggests a minimum of 120 reference subjects to be used for 90% confidence limits and 150 for 95% confidence limits [5]. The study population consists of apparently healthy individuals between the ages of 20 and 80 years living in Port Harcourt, who were not known diabetic or had history of hemoglobinopathy. Questionnaire was used to exclude hemoglobinopathies, diabetes mellitus and any other chronic illness. Physical examination and packed cell volume was used to exclude subjects with anaemia.

2.1 Measurement of Height

Height of subjects was measured using a height meter. The subject stands with feet together, without shoes or headgear. Back and heels against a vertically ruled bar to which a movable horizontal bar was applied to the vertex of the subject's head and measurement taken in meters (M^2) [6].

2.2 Measurement of Weight

Weight was measured using a weighing scale. Subjects were requested to wear light cloths and then stand erect on the scale. The weight was measured in kilograms (kg). The BMI was then calculated by dividing weight (kg) by the height in meters, squared⁶ (m^2).

2.3 Blood Collection and Processing

Subjects were made to sit comfortably and a tourniquet was tied at any point of the upper arm to make the veins prominent. A vacutainer was used to collect two milliliters (2 ml) of blood into an ethylenediamine tetra acetic acid [EDTA] bottle. These bottles were gently rolled over to allow the anticoagulant in the bottles mix with the blood. The blood samples were analyzed in batches later same day for glycated haemoglobin and PCV.

2.4 Laboratory Analysis

2.4.1 PCV and haemoglobin estimation

The mixed whole blood in EDTA bottle was aspirated into a capillary tube till the tube was about two third filled. One end of the capillary was sealed with plasticine and placed in the haematocrit centrifuge with the sealed end toward the centrifuge periphery. The packed cell volume (PCV) was determined by centrifuging blood in the capillary tube (also known as a micro haematocrit tube) at 10,000 rpm for five minutes. The haematocrit reader was used to calculate the PCV. The average of the duplicate readings in percentage was taken as the PCV value for each subject. The PCV value was divided by 3 to give a rough estimate of haemoglobin concentration in g/dl. [7].

2.4.2 Glycated hemoglobin

The sample in the EDTA bottle was analyzed same day using the chromatographic technique by Bio system for the glycated haemoglobin analysis [8].

2.4.3 Principle

After preparing the haemolysate, where the labile fraction is eliminated, hemoglobins are retained by a cationic exchange resin. Glycated Hemoglobin (HbA_{1c}) is specifically eluted after washing away the hemoglobin A_{1a+b} fraction. (HbA_{1c}) is quantified by direct photometric reading at 415nm. The estimation of the relative concentration of HbA_{1c} is made by the measure of total hemoglobin concentration by direct photometric reading at 415 nm.

Calculations

$$\frac{\text{Absorbance of HbA}_{1c}}{\text{Absorbance of Hb}_{\text{TOTAL}}} \times 100 = \% \text{HbA}_{1c}$$

2.5 Quality Assurance

All guidelines for specimen collection, processing, storage and handling was strictly adhered to. Laboratory analysis of samples was carried out with inclusion of quality control (QC) samples in which within run and between run quality control samples were assayed. Instrument efficiency was checked through regular calibration and recalibration.

3. RESULTS

The data generated was analyzed using Statistical Package for Social Sciences versions 20.0 (SPSS 20.0) Data was ranked and mean, median and mode were calculated and outliers were removed using the Dixon outliers' statistic [9]. The non-parametric method was used to determine the reference interval due to the skewed nature of the data where the 2.5th and the 97.5th percentile of the ranked data represented lower and upper reference limits [10]. Multiple regression analysis was used to determine the association of PCV, age and BMI with glycated haemoglobin values of the reference population and the level of significance was set at p-value < 0.05

A total of 172 subjects recruited for this study were made up of 99(51.7%) female and 83(48.3%) male subjects. A total of 99 subjects had a BMI above 25 kg/m² while 73 subjects had BMI below BMI ≤ 25 kg/m².

The mean age for the study population was 35.4 years and the mean glycated hemoglobin was 4.84%. The data, when divided into male and female groups had a mean of 4.63% for the female and 4.61% for the male group. The reference subjects had minimum and maximum glycated hemoglobin values of 3.5% and 6.0% respectively (Table 1) and a reference interval of 4.0%- 5.9% (Table 1).

Table 1. Results of HbA_{1c}

| | Total subjects(n=172) |
|--------------------------------|------------------------------|
| Mean age | 35.35 |
| Male | 83 |
| Female | 89 |
| HbA _{1c} (%) | 3.5 |
| Min value | 6.0 |
| Max value | 4.84 (0.04) |
| Total population (SEM) | 4.83 (0.05) |
| Males (SEM) | 4.85 (0.05) |
| Females (SEM) | 4.0-5.9 |
| Ref interval HbA _{1c} | |

4. DISCUSSION

The advantages of glycated haemoglobin over fasting plasma glucose have made it an important index in the screening, diagnosis and management of diabetes.

Table 2. Test of association between HbA1c and other factors using multiple linear regression analysis in the total population

| Model | Unstandardized coefficients | | Standardized coefficients | Students' t-test | Sig. |
|------------|-----------------------------|------------|---------------------------|------------------|-------|
| | Beta | Std. error | Beta | | |
| (Constant) | 2.740 | .225 | | 12.18 | 0.001 |
| Ages | .012 | .002 | .324 | 5.69 | 0.001 |
| PCV | .014 | .005 | .120 | 2.57 | 0.01 |
| BMI | .012 | .005 | .128 | 2.38 | 0.02 |
| Alcohol | .042 | .033 | .063 | 1.27 | 0.20 |

a. Dependent variable: hba1c

More so, studies have shown that diabetics with glycated haemoglobin value around the upper reference limit are more likely to develop diabetic complications than those with lower values [11]. To this effect, a population based reference value of glycated haemoglobin is very necessary as the upper limit of normal varies from one population to another. The reference interval of glycated haemoglobin of a local population will greatly influence the management of diabetic subjects, bearing in mind the association of glycated haemoglobin value and diabetic complications.

The changes in lifestyle and eating habits, apart from genetic, age and race factors has been shown to affect blood glucose value and so by extension affect the level of glycation of red blood cells.[12] Though the reference interval of fasting plasma glucose in this region was determined in a study done in 2008, [13] no literature has been cited on the reference interval of glycated haemoglobin in this area.

From this study the mean glycated haemoglobin value was 4.84% and the reference interval of glycated haemoglobin was found to be 4.0 – 5.9%. This is lower than the reference interval given by the diagnostic kit manufacturers, which was given as 4.0 – 6.5% [14] and also different from those obtained in other sub Saharan African studies[15,16,17]. The glycated haemoglobin value determined in a study in Kumasi Ghana using the chromatographic method was 3.7 – 7.0% [15]. This interval was found to be statistically and clinically significant when compared to the reference interval of this study even though both cities are located in the sub-Saharan Africa region. This difference may be due to the type of diet and the level of diabetes awareness which may have led to lifestyle modifications [18] A study done in Pakistan showed a reference interval of 4.6 – 6.6% [19]. This was also found to be both statistically different and clinically significant from the reference interval determined in this study and in

other studies. The variation of the reference intervals shown above is quite remarkable and only re-emphasizes the fact that even when using the same method, every laboratory should determine the reference interval of its population [15,20] for better management of patients. The use of reference interval given by diagnostic kits manufacturers for our population would therefore make for poor screening, diagnosis, monitoring and treatment of diabetic patients.

In this study, it was also noted that the mean glycated haemoglobin value for females was slightly higher than that of males but the difference was not statistically significant. This finding is in agreement with other studies, [21,22] though some had contrary findings [23]. It also may explain why the prevalence of diabetes among African Americans is higher in female than in males [24]. The positive association of glycated haemoglobin with PCV and age in this study, as in other studies, affirms that glycated haemoglobin value may not be too reliable in anemic subjects and in those with haemoglobinopathies. Anemia has been shown to be an independent determinant of glycated haemoglobin value [21]. The positive association of glycated haemoglobin with age may call for a possible age related partitioning of the reference interval of glycated haemoglobin.

5. CONCLUSION

The reference interval for this study is significantly different from that provided by the kit of manufacturer and those from other regions of the world. The introduction of this local reference interval will improve the management of diabetic patient in our region.

CONSENT AND ETHICAL APPROVAL

Ethical approval was obtained from the Ethical committee of the University of Port Harcourt Teaching hospital and all subjects gave informed

written consent. A total of 172 apparently healthy reference subjects who met the inclusion criteria and signed the consent form were recruited for the study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Zhang P, Zhang X, Brown J, Visitisen D, Sicree R, Shaw J, Nicholas G. Global healthcare expenditure on diabetes for 2010 and 2030. *Diabetes Res Clin Pract.* 2010;87:293–301.
2. Nyenwe EA, Odia OJ, Ihekweba AE, Ojule A, Babatunde S. Type 2 diabetes in adult Nigerians: A study of its prevalence and risk factors in Port Harcourt, Nigeria. *Diabetes Res Clin Pract.* 62:177-185.
3. Brownlee M, Hirsch IB. Glycaemic variability: a haemoglobin A_{1c}-independent risk factor for diabetic complications. *JAMA.* 2006;295:1707-1708.
4. Nathan DM, Turgeon H, Regan S. Relationship between glycated haemoglobin levels and mean glucose levels over time. *Diabetologia.* 2007;50:2239-2244.
5. Solberg HE. A guide to IFCC recommendations on reference values. *J Int Fed Clin Chem.* 1993;5:162-165.
6. Keys A, Fidanza F, Karvonen MJ, Kimura N, Taylor H. Indices of relative weight and obesity. *Int J Epidemiol.* 2014;43(3):655-665.
7. Connor GO, Molloy AM, Daly L, Scott JM. Deriving a useful packed cell volume estimate from haemoglobin analysis. *J Clin Pathol.* 1994;47:78–79.
8. Lakshmy R, Gupta R. Measurement of glycated hemoglobin A1c from dried blood by turbidimetric immunoassay. *Journal of Diabetes Science and Technology.* 2009; 3(5):1203-1206.
9. Horn PS, Feng L, Li Y, Pesce AJ. Effect of outliers and non healthy individuals on reference interval estimation. *Clin Chem.* 2001;47:2137-2145.
10. Solberg HE. A guide to IFCC recommendations on reference values. *J Int Fed Clin Chem.* 1993;5:162-165.
11. International Organization for Standardization (ISO). Medical laboratories – particular requirements for quality and competence. Geneva: ISO. 2012;15189: 36-37.
12. Kuhl J, Hilding A, Östenson CG, Grill V, Efendic S, Båvenholm P. Characterization of subjects with early abnormalities of glucose tolerance in the Stockholm Diabetes Prevention Programme: The impact of sex and type 2 diabetes heredity. *Diabetologia.* 2005;48(1):35-40.
13. Ejilemele AA, Ojule AC, Orluwene CG. Plasma glucose in a reference population in Port Harcourt, Nigeria. *PMJ.* 2008;2 (3):228-232.
14. Larsen ML, Horder M, Mogensen EF. Effect of long term monitoring of glycated haemoglobin levels in insulin dependent diabetes mellitus. *N Engl J Med.* 1990; 323:1021-1025.
15. Arthur FKN, Yeboah FA, Nsiah K, Nkrumah PKN, Afreh KA, Agyenim-Boateng K. Fasting blood glucose and glycosylated haemoglobin levels in randomly selected Ghanaian diabetic patients – the clinical implications. *J Sci Technol.* 2005;25(2):13-17.
16. Naby B, Alioune C, Joelle Sobngwi-T, Eric VB, Alain T, Leopold F, Liman S, Ngamani S, Ngapout S, Kengne AP, Sobngwi E. Improving access to HbA_{1c} in sub-Saharan Africa (IA3) cohort: Cohort profile. *Pan Afr Med J.* 2017;27:275.
17. Mbanya JCN, Motala AA, Sobngwi E, Assah FK, Enoru ST. Diabetes in Sub-Saharan Africa. *The Lancet.* 2010; 375(9733):2254–2266.
18. Sánchez-González CM, Castillo-Mora A, Alvarado-Maldonado IN, Carlos Ortega G, Martínez-CN, Arce-Sánchez L, Valencia MR, Hernandez AM, Gutierrez GE, Salvador Espino YS, Lopez YR, Sanchez RH, Munoz ER. Reference intervals for hemoglobin A1c (HbA_{1c}) in healthy Mexican pregnant women: A cross-sectional study. *BMC Pregnancy Childbirth.* 2018;18(1):424.
19. Ali SK, Khan DA, Khan FA. Determination of glycosylated haemoglobin reference range in adult population attending a military care set up in Rawalpindi. *PAFMJ.* 2011;61(4):542-545.

20. Salwa HN, Emad AR. Determining the reference range Values of glycosylated hemoglobin (HbA1c) by immunoturbid assay in Iraqi population. *Natl J Chem.* 2011;41:127-134.
21. Jonsson PM, Sterky G, Gafvels C, Ostman J. Gender equity in health care: The case of Swedish diabetes care. *Health Care Women Int.* 2000;21(5):413-431.
22. Paul U, Dubuc B, Keyes Scott. Sex differences in glycosylated hemoglobin in diabetic and non-diabetic C57BL/6 mice. *Diabetes Res and Clin Pract.* 1993; 21:95-101.
23. Park S, Barrett-Connor E, Wingard DL, Shan J, Edelstein S. Glycated haemoglobin is a better predictor of cardiovascular disease than fasting or post challenge plasma glucose in women without diabetes. *The Rancho Bernardo Study. Diabetes Care.* 1996;19:450–456.
24. William HH, Robert MC. Racial and ethnic differences in the relationship between HbA1c and blood glucose: Implications for the Diagnosis of Diabetes. *J Clin Endocrinol Metab.* 2012; 97(4):1067–1072.

© 2019 Igila and Aaron; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/51683>