



Evaluation of Copper Status and Some Red Cell Parameters of Pregnant Women in Enugu State, South Eastern Nigeria

**Richard Eze¹, Emmanuel Ifeanyi Obeagu^{1,2*}, Arvin Nwakulite¹, I. L. Okoroiwu²,
C. C. N. Vincent³, Chukwuma J. Okafor⁴, Ejike Felix Chukwurah⁵,
Udunma Olive Chijioke⁶ and Chukwudi Ofodile Amaechi⁷**

¹*Department of Medical Laboratory Science, Madonna University, Elele, Rivers State, Nigeria.*

²*Department of Medical Laboratory Science, Imo State University, Owerri, Imo State, Nigeria.*

³*Department of Nursing Science, Imo State University, Owerri, Imo State, Nigeria.*

⁴*Department of Pathology and Biochemistry, State University of Zanzibar, Tanzania.*

⁵*Department of Haematology and Immunology, Faculty of Clinical Medicine, Ebonyi State University, Abakaliki, Nigeria.*

⁶*Department of Health Administration and Management, Faculty of Health Sciences, University of Nigeria, Enugu Campus, Enugu State, Nigeria.*

⁷*Department of Medical Laboratory Science, Nnamdi Azikiwe University, Nnewi Campus, Nnewi, Anambra State, Nigeria.*

Authors' contributions

This work was carried out in collaboration among all authors. Authors RE, EIO, AN and ILO designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors CCNV, CJO, EFC, UOC and COA managed the analyses of the study and managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i30A31616

Editor(s):

(1) Dr. Ozlem Sultan Aslanturk, Adnan Menderes University, Turkey.

Reviewers:

(1) Sofia Febuanti, Poltekkes Kemenkes Tasikmalaya, Indonesia.

(2) Sam Froze Jiee, Malaysia.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/68772>

Original Research Article

Received 15 March 2021

Accepted 21 May 2021

Published 29 May 2021

ABSTRACT

Copper is an essential trace mineral (micronutrient) that is naturally present in some foods and is available as a dietary supplement. Pregnancy, also known as gestation, is the time during which one or more offspring develops inside a woman. This study was designed to assay the copper level

of pregnant women in urban area and to evaluate the effect of the obtained values on some blood parameters of the pregnant women. In this study, a total of 100 subjects and 50 controls was used. subjects were pregnant women while the controls were non-pregnant women. The copper level in subject was assayed with a serum copper test kit using automation with Bio E-lab fully automated chemistry analyzer and result of the complete blood count was gotten using Mindray Hematology analyzer. The mean for subjects ($M=102.83\pm 1.38$) and control ($M=102.68\pm 0.94$), $p = 0.71$. These results suggest that the difference between the copper level for subjects and controls is not really much.. The variables copper level and MCV were found to be slightly correlated, $r(150) = .018$., $p > .005$. Additionally, copper levels in the subjects was also found to negatively correlated with MCH, $r(150) = .907$, $p > .005$. One-way analysis of variance showed that the correlation of copper levels was not significant on all of the red bloodcell parameters in the serum of the pregnant women.

Keywords: Copper status; red cell parameters; pregnant women.

1. INTRODUCTION

Copper is an essential trace mineral (micronutrient) that is naturally present in some foods and is available as a dietary supplement. It is a cofactor for several enzymes (known as "cuproenzymes") involved in energy production, iron metabolism, neuropeptide activation, connective tissue synthesis, and neurotransmitter synthesis [1]. One abundant cuproenzyme is ceruloplasmin (CP), which plays a role in iron metabolism and carries more than 95% of the total copper in healthy human plasma). In humans, copper is essential to the proper functioning of organs and metabolic processes. The human body has complex homeostatic mechanisms which attempt to ensure a constant supply of available copper, while eliminating excess copper whenever this occurs [2].

The availability of copper in suburban areas is usually high because of the use in production of many items like electrical wiring and plumbing materials such as household water pipes. These pipes and plumbing fixtures such as brass, faucet contain copper which can leach into the drinking water when the copper level in the water exceeds the action level can cause problem to health [3].

They may be elevated level of copper in carbonated or acidic beverage that contact copper tubing .carbon dioxide can corrode the plumbing container which can add copper to softdrink. When these contaminated water and softdrinks are consumed together with the dietary copper , it can skyrocket the copper level in the blood of pregnant women which may lead to copper toxicity. unlike the rural areas that lack all of these advanced production that affect the copper level [4].

The average human diet provides approximately 1,400 mcg/day for men and 1,100 mcg/day for women that is primarily absorbed in the upper small intestine . Almost two-thirds of the body's copper is located in the skeleton and muscle [4]. Only small amounts of copper are typically stored in the body, and the average adult has a total body content of 50–120 mg copper . Most copper is excreted in bile, and a small amount is excreted in urine. Copper levels in the body are homeostatically maintained by copper absorption from the intestine and copper release by the liver into bile to provide protection from copper deficiency and toxicity [4].

Pregnancy, also known as gestation, is the time during which one or more offspring develops inside a woman . During pregnancy the plasma volume increases by 40-50% and the red blood cell volume increases only by 20–30%. These changes occur mostly in the second trimester and prior to 32 weeks gestation. Due to dilution, the net result is a decrease in haematocrit or hemoglobin, which is a measure of red blood cell concentration. Erythropoietin, which stimulates red blood cell production, increases throughout pregnancy and reaches approximately 150 percent of their pregnancy levels at term [5].

Copper is important during pregnancy, when blood supply doubles and energy level drops and also helps in creating the fetus heart blood vessels and cells as well as nervous , immune and skeletal systems. In pregnancy ,excess copper levels can be associated with intrauterine growth restriction ,preclampsia and neurological disease because of oxidative damage caused by copper accumulation in the placenta and fetal tissue. Sometimes the changes in certain hormone levels and their effects on their target organs can lead to gestational diabetes and gestational hypertension. A pregnant woman will

also become hypercoagulable, leading to increased risk for developing blood clots and embolisms, such as deep vein thrombosis and pulmonary embolism. Women are 4-5 times more likely to develop a clot during pregnancy and in the postpartum period than when they are not pregnant.

The study was done to estimate the copper level in pregnant women in an urban area.

2. MATERIALS AND METHODS

2.1 Study Area

The research was carried out on apparently healthy pregnant women in a maternity hospital in Enugu town, Enugu state, Nigeria. It is located in the southeastern part of Nigeria.

2.2 Study Population

A total number of 100 pregnant women mean age $20 \pm$ years were recruited from the maternity hospital and their blood sample was collected. A total number of 50 non pregnant women mean age of $20 \pm$ was used as control in the research. They all gave their consent to participate in this study.

2.2.1 Procurement of copper

A commercially prepared serum copper test kit product of Centronic GmbH Am Kleinfeld 11 Wartenberg Germany used to assay the copper serum level.

2.3 Sample Collection Technique

A standard clean venepuncture was used to collect the blood. 5ml of blood was collected from a prominent vein in the antecubital fossa of the arm of the subjects. 2ml of the blood sample was delivered in a plain container and the remaining 3ml of the blood sample was transferred in a commercially prepared Ethylene diamine tetra acetic acid (EDTA) container. The blood sample in the plain container was spun with a centrifuge, the serum extracted using an automatic micropipette and placed in a labelled container. The sample was frozen until analysis and the analysis was carried out in Bioquest diagnostic laboratory in Parklane avenue in Enugu state.

2.4 Inclusion Criteria

Healthy pregnancy women mean age of $20 \pm$ and healthy non pregnant women mean age $20 \pm$ in

urban area were used in the research as test subject and control respectively.

2.5 Exclusion Criteria

Healthy pregnancy women mean age of $20 \pm$ and healthy non pregnant women mean age $20 \pm$ in the rural area were excluded in this study.

2.6 Methods

2.6.1 Estimation of copper level method

Automation Bio E lab fully automated chemistry analyzer model AS – 280 product of Nanjing city, Jiangsu province China.

The Bio E lab fully automated chemistry analyzer is an instrument that uses the pale yellow supernatant portion (serum) of centrifuged blood sample or a urine sample, and induces reactions using reagents to measure various components, such as sugar, cholesterol, protein, enzyme, etc.

2.6.2 Estimation of Red cell parameter

Methods: Automation using Mindray Auto Haematology Analyzer Model BC-2800

2.7 Statistical Analysis

Data obtained were analyzed for mean and standard deviation. Statistical analysis was conducted using Integrated Business Machine Statistical Package for social sciences (IBM – SPSS), version 2.0 and t-test level of significance was considered as <0.05 and >0.05 .

3. RESULTS

The mean copper level of the subjects was 353.39 and S.D of 87.63. That of the control was 322.41 and S.D of 80.27.

4. DISCUSSION

Trace Element (trace metals) are minerals present in living tissues in small amount. Copper is one of the many trace metals and it's widely distributed in food and water. This study compares the copper level in pregnant women and non pregnant as test subject and control respectively while checking the effect the copper level will have on Red blood cell parameters in the pregnant women.

In the study the mean differences in the copper level of the pregnant and non pregnant women

Table 1. Mean \pm S.D of copper level among subjects and control

Parameter	Mean Value (Subjects)	Mean Value (Control)	t-value	p-value
Copper Level(ug/dL)	353.39 \pm 87.63	322.41 \pm 80.27	2.098	.038
Mean Difference: 30.98				

A t-test was conducted to compare copper Level for subjects and controls. The mean for subjects ($M=102.83$, $SD=1.38$) and control ($M=102.68$, $SD=0.94$) conditions; $t(150) = .484$, $p = 0.71$. These results suggest that the difference between the copper level for subjects and controls is not really much. Specifically, our results suggest that when the difference is insignificant.

Table 2. Correlation of copper level in the serum and the various red blood cell parameters

Parameters	Copper Levels			X ²	Sig.	R	sig
	Hypocupraemia	Normal Level	Hypercupraemia				
Hb				16.04	.000	-.401	.000
Severe (<7g/dl)	1(1.0)	83(83.0)	0(0.0)				
Mild (9-11g/dl)	4(4.0)	12(12.0)	0(0.0)				
PCV				.107	.743	-.033	.746
Moderate	0(0.0)	2(2.0)	0(0.0)				
Slight	5(5.0)	93(93.0)	0(0.0)				
MCHC				.336	.562	-.058	.567
Low (200-300g/L)	0(0.0)	6(6.0)	0(0.0)				
Normal (315-360g/L)	5(5.0)	89(89.0)	0(0.0)				
High (>350g/L)	0(0.0)						
MCV				.034	.855	.018	.856
Microcytic Anaemia (20-70 μ g/cell)	1(1.0)	4(4.0)	0(0.0)				
Normal	16(16.0)	79(79.0)	0(0.0)				
Macrocytic Anaemia (>100)	0(0.0)	0(0.0)	0(0.0)				
MCH				.014	.905	.907-	.012
Hypochromia	1(1.0)	4(4.4)	0(0.0)				
Normal	17(17.0)	78(78.0)	0(0.0)				
Others	0(0.0)		0(0.0)				

The variables copper level and MCV were found to be slightly correlated, $r(150) = .018$, $p > .005$. Additionally, copper levels in the subjects was also found to negatively correlated with MCH, $r(150) = .907$, $p > .005$. A positive correlation means the relationship between the Red blood cell parameters and copper level gearing towards the same direction

was low (0.71). The p-value is greater 0.05, this indicates that the differences was not significant. In the relationship between copper level and the Red blood cell parameters the variables were found to be slightly correlated, $r(150) = .018$, $p > .005$. there is no significant difference between copper level and Red blood cell parameters. Copper deficiency has not been documented in humans during pregnancy. It is quite possible that the demonstrated teratogenic effects of the drug penicillamine, which is a copper chelator, may be mediated through copper deficiency, but copper status has not been investigated in the reported cases. Copper level can be analyzed in both blood and water (drinkable water, sea water) using copper kit and this is done to maintain a normal copper level in diet and water to avoid hypocupremia (low blood copper level) and copper toxicity Obeagu, [6-9].

5. CONCLUSION

This study proves from the results obtained from that the difference between the copper level for subjects and controls is not really much, and there is no significant difference, analysis of variance showed that the effect of copper levels was not significant on all of the Red blood cell parameters in the serum of the pregnant women.

CONSENT AND ETHICAL APPROVAL

Approval was gotten from the management of the maternity hospital and the consent of the subject and control was also obtained.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Klevay LM. Copper. In: Coates PM, Betz JM, Blackman MR. edition Encyclopedia

- of Dietary Supplements. 2nd edition London and New York: Information Healthcare. 2010;604-611.
2. Klevay LM. Is the Western diet adequate in copper? Journal of Trace Element in Medicine and Biology. 2011;25:204-212.
3. Rosado JL. Zinc and copper: proposed fortification levels and recommended zinc compounds. Journal of Nutrition. 2003;133:298.
4. Squitti R, Simonelli I, Ventriglia M, Siotto M, Pasqualetti P, Rembach A, et al. Meta-analysis of serum non-ceruloplasmin copper in Alzheimer's disease. Journal of Alzheimers Disease. 2014;38:809-822.
5. Theobald HE. Eating for pregnancy and breast-feeding". The Journal of Family Health Care. 2007;17(2):45-49.
6. Obeagu EI. A Review on Pregnancy and Haematology. International Journal of Current Research in Biology and Medicine. 2018;3(5):26-28.
7. Obeagu EI, Obarezi TN, Eze OBL, Emelike CU. Haematological profile of pregnant women in Umuahia, Abia State, Nigeria. International Journal of Current Microbiology and Applied Science. 2014;1(3):713-718.
8. Obeagu EI, Adepoju OJ, Okafor CJ, Obeagu GU, Ibekwe AM, Okpala PU, et al. Assessment of Haematological Changes in Pregnant Women of Ido, Ondo State, Nigeria, J Res Med Dent Sci. 2021;9(4):145-148.
9. Eze R, Ezeah GAC, Obeagu EI, Omeje C, Nwakulite A. Evaluation of iron status and some haematological parameters of pregnant women in Enugu, South Eastern Nigeria. World Journal of Pharmaceutical and Medical Research. 2021;7(5):251-254.

© 2021 Richard et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/68772>