



## **COVID-19 Vaccination Status amongst College Students and Staff during Offline Classes**

**A. Sai Padma<sup>a\*</sup> and N. Chandan Babu<sup>b#</sup>**

<sup>a</sup> Department of Biochemistry, Bhavan's Vivekananda College, Sainikpuri, Secunderabad, Telangana State, India.

<sup>b</sup> Department of Mathematics and Statistics, Bhavan's Vivekananda College, Sainikpuri, Secunderabad, Telangana State, India.

### **Authors' contributions**

*This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/JPRI/2021/v33i57B34050

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/77739>

**Short Communication**

**Received 06 November 2021**  
**Accepted 12 December 2021**  
**Published 14 December 2021**

## **ABSTRACT**

**Aim:** To understand the status of COVID-19 vaccination amongst students and staff during offline classes.

**Study Design:** Questionnaire with open ended questions was sent to participants through social media.

**Place and Duration of Study:** The present study was conducted in Bhavan's Vivekananda College of Science, Humanities and Commerce, Sainikpuri, Secunderabad from 12<sup>th</sup>-15<sup>th</sup> September, 2021.

**Methodology:** Questionnaire was sent to all staff members, second and third year students of undergraduate and first year students of post graduate programmes who were attending classes during the study period. Total 1263 responses were received out of approximately 2500 students.

**Results:** The survey results have shown that 1128 participants (89.3%) out of 1263 have received vaccination and out of them, 742 participants (65.78%) have received single dose of vaccine and 386 (34.22%) participants have received two doses. Majority of them have received Covishield (74.5%), followed by Covaxin (24.11%).

**Conclusion:** The present study necessitates all educational institutions to conduct similar kind of

# Assistant Professor;

\*Corresponding author: E-mail: saipadmabhavans@yahoo.co.in;

studies to understand the current status of vaccination and safety measures taken by students and staff during Covid-19 pandemic during offline classes. This will help to keep the campuses safe and prevent further spread of infection to avoid third wave of corona virus.

*Keywords: COVID-19 vaccination; college students; types of vaccines; offline classes.*

## 1. INTRODUCTION

Though strong measures were taken by India to contain the spread of COVID-19 through better diagnostics and treatment, actual solution will be provided by vaccines by enhancing immunity and containing the disease spread [1]. For all elements of COVID-19 vaccine administration in India, the Indian government has constituted a National Expert Group on Vaccine Administration for COVID-19 (NEGVAC) in April 2020 [2]. The Guidelines for National COVID Vaccination Program have been reviewed and revised on 21<sup>st</sup> June 2021 and states that vaccination will be prioritized as the following: Health care workers, front line workers, citizens above 45 years of age, citizens with second dose due, citizens 18 years and above. States/UTs may decide their own prioritization of vaccine schedule within the population group of citizens more than 18 years of age [3]. The eligible age for vaccination in India is currently 18 and above and this will surely benefit both undergraduate and postgraduate students as most of them fall in the age group of 18 - 23 years [3]. Approved COVID-19 vaccines provide a high degree of protection against getting seriously ill and dying from the disease, although no vaccine is 100% protective [4]. The safety and efficacy data from clinical trials of vaccine candidates are examined by Drug regulator of India and all the COVID-19 vaccines that receive license will have comparable safety and efficacy [5]. However, it must be ensured that the entire schedule of vaccination is completed by only one type of vaccine as different COVID-19 vaccines are not interchangeable. The Covid Vaccine Intelligence Network (CoWIN) platform is providing convenient and safe pre-booking of vaccination appointments [6]. India's drug regulator has approved emergency use of Covishield (the Oxford-AstraZeneca vaccine in India) and Covaxin, manufactured by Bharat Biotech with restricted use [7].

To comprehend the vaccine confidence and hesitancy, various parameters like vaccine confidence-trust in the effectiveness and safety of vaccine, vaccination complacency where vaccination is not a deemed preventive action,

vaccination convenience where availability is easy, are taken into consideration [8]. A survey in India in December, 2020 indicates that approximately 11,000 respondents revealed that 53% were unsure about taking the COVID-19 vaccine [9]. About 69% of respondents have mentioned in another citizen-survey platform in Delhi that there is no urgent need to get immunized and the key reasons for this kind of hesitancy included restricted information about side-effects, efficacy levels, and perceived high immunity levels [10]. As per available recent data on vaccination status in India at large, 42.2% people have got at least one dose and fully vaccinated are only 13.5% [11]. In account of all the above information stated, there is a need to know the actual status of vaccination in the current scenario where colleges have opened for physical classes and urgent requirement for 100% vaccination.

## 2. METHODOLOGY

The methodology for the study included a questionnaire with most of the questions contained pre-defined answers and circulated as Google form and a total number of 1263 participants have responded to the questionnaire. The responses were analyzed by using statistical tools like Chi-square test, ANOVA one way classification and descriptive statistics through python.

**The major observations from the study are:** 1. Majority of participants are in the age of 18-20 years, followed by 21-23 years. 2. The students who have responded to the questionnaire are mostly undergraduate as percentage of students in colleges belong to undergraduate courses is more than in post graduate courses. 3. A good percentage of participants are vaccinated (89.3%), with majority of them received single dose (Table.1).

4. The percentage of study participants received two doses of vaccine is comparatively higher (34.2%) when compared to reported values of overall population in India at time of study [11]. 5. Participants have not shown much hesitancy towards vaccine administration in the current

situation when compared to earlier studies [9,10] sounding any valid. 7. There is a strong association observed with respect of gender and age to type of vaccine received (Table.2).

**Table 1. Number and percentage indication of responses to various questions in the questionnaire, pertaining to vaccination in the colleges**

| S. No | Parameters  | Category  | No. of responses | % of responses |
|-------|---|---|------------------|----------------|
| 1     | Gender of participants  | Male  | 525              | 41.6%          |
|       |   | Female  | 738              | 58.4%          |
| 2     | Age of participants   | 18-20   | 955              | 75.6%          |
|       |   | 21-23   | 216              | 17.1%          |
|       |   | 24-26   | 15               | 1.2%           |
|       |   | 27-30   | 0                | 0%             |
|       |   | 31-40   | 24               | 1.9%           |
|       |   | 41-50   | 23               | 1.8%           |
|       |   | 51-60   | 17               | 1.3%           |
| 3     | Semester of study /staff  | >60   | 11               | 0.9%           |
|       |   | First   | 51               | 4%             |
|       |   | Second  | 201              | 15.9%          |
|       |   | Third   | 501              | 39.7%          |
|       |   | Fifth   | 450              | 35.6%          |
| 4     | Nature of participants  | Staff   | 60               | 4.8%           |
|       |   | Undergraduates  | 1019             | 80.7%          |
|       |   | Postgraduates   | 184              | 14.6%          |
| 5     | Number and % of participants vaccinated for COVID-19 during study period.     | Staff   | 60               | 4.8%           |
|       |   | Vaccinated  | 1128             | 89.3%          |
| 6     | Non-vaccinated participants planning to take it in near future.               | Not vaccinated  | 135              | 10.7%          |
|       |   | Planning to take  | 121              | 89.3%          |
| 7     | Dose of vaccine completed   | Not Planning to take                                    | 14               | 10.7%          |
|       |   | 1 <sup>st</sup> Dose                                    | 742              | 65.78%         |
| 8     | Inhibitions about vaccine safety due to short time testing in clinical trails | 2 <sup>nd</sup> Dose                                    | 386              | 34.22%         |
|       |   | Having inhibitions                                      | 344              | 27.2%          |
| 9     | Type of vaccine taken based on choice or availability                         | Not having  | 919              | 72.8%          |
|       |   | Covaxin   | 272              | 24.11%         |
|       |   | Covishield  | 841              | 74.55%         |
|       |   | Sputnik V   | 8                | 0.7%           |
| 12    | Reasons for refusing the vaccine  | Other   | 7                | 0.6%           |
|       |   | Feeling as vaccine doesn't have any positive benefits   | 15               | 10.9%          |
|       |   | Person feeling as healthy                               | 16               | 11.9%          |
|       |   | Anticipating as vaccination causes serious side effects | 9                | 7%             |
|       |   | Fear as painful act                                     | 10               | 7.6%           |
|       |   | Vaccine not available                                   | 19               | 13.9%          |
|       |   | Does not want to reveal                                 | 66               | 48.7%          |

**Table 2. Association between Gender and Age to the type of Vaccine received**

| <b>Null Hypothesis H<sub>0</sub>: There is no Association between Gender and Type of Vaccine received</b>         |                                 |                   |                       |                       |              |                    |
|---|---------------------------------|-------------------|-----------------------|-----------------------|--------------|--------------------|
| <b>Alternative Hypothesis H<sub>1</sub>: There is an Association between Gender and Type of Vaccine received.</b> |                                 |                   |                       |                       |              |                    |
| <b>Gender</b>   | <b>Type of Vaccine received</b> |                   |                       |                       |              |                    |
|   | <b>Covaxin</b>                  | <b>Covishield</b> | <b>Sputnik V</b>      | <b>Not vaccinated</b> | <b>Other</b> | <b>Grand Total</b> |
| Female  | 179                             | 479               | 5                     | 69                    | 6            | 738                |
| Male  | 93                              | 362               | 3                     | 66                    | 1            | 525                |
| Grand Total   | 272                             | 841               | 8                     | 135                   | 7            | 1263               |
| Gender Vs Type of Vaccine received  | Test                            | p-value           | Conclusion            |                       |              |                    |
|   | Chi-square                      | 0.017             | Reject H <sub>0</sub> |                       |              |                    |
| <b>b. Null Hypothesis H<sub>0</sub>: There is no Association between Age and Type of Vaccine received</b>         |                                 |                   |                       |                       |              |                    |
| <b>Alternative Hypothesis H<sub>1</sub>: There is an Association between Age and Type of Vaccine received</b>     |                                 |                   |                       |                       |              |                    |
| <b>Row Labels</b>   | <b>Type of Vaccine received</b> |                   |                       |                       |              |                    |
|   | <b>Covaxin</b>                  | <b>Covishield</b> | <b>Sputnik V</b>      | <b>Not vaccinated</b> | <b>Other</b> | <b>Total</b>       |
| 18-20   | 195                             | 650               | 6                     | 97                    | 7            | 955                |
| 21-23   | 55                              | 138               | 1                     | 22                    | 0            | 216                |
| 24-26   | 3                               | 9                 | 0                     | 3                     | 0            | 15                 |
| 27-30   | 0                               | 2                 | 0                     | 0                     | 0            | 2                  |
| 31-40   | 9                               | 11                | 1                     | 3                     | 0            | 24                 |
| 41-50   | 6                               | 17                | 0                     | 0                     | 0            | 23                 |
| 51-60   | 4                               | 13                | 0                     | 0                     | 0            | 17                 |
| Other   | 0                               | 1                 | 0                     | 10                    | 0            | 11                 |
| Grand Total   | 272                             | 841               | 8                     | 135                   | 7            | 1263               |
| Age Vs Type of Vaccine received   | Test                            | p-value           | Conclusion            |                       |              |                    |
|   | Chi-square                      | 0.000000020       | Reject H <sub>0</sub> |                       |              |                    |

The present study will help to understand the current status of vaccination in the country as a sample model and helps to identify unvaccinated students and staff and develop strategies to educate them to receive vaccination and achieve 100% vaccination in the college campus to make protective against COVID-19 infections, apart from following other COVID protocols. Many universities are checking the vaccination data of students joined and if once institutes have this data, it will be easier for them to manage students as and when they come to the campus or hostels [12]. Vaccination status has no bearing on the admission process, and unvaccinated students are still permitted. A similar database for students under 18 will help in the similar lines for higher secondary educational institutions, but the vaccination hasn't been rolled out yet for them, necessitates to be more careful in code of conduct against COVID-19.

**3. CONCLUSION**

Students and staff of all colleges in the country need to be well educated and motivated by conducting similar kinds of studies and promoting vaccination drives in the college campuses themselves to ensure 100% vaccination. Keeping

campuses safe and preventing the spread of infection is the need of the day for any educational organization. It is advised to follow fresh recommendations of the National Expert Group on Vaccine Administration for COVID-19 (NEGVAC) regarding COVID-19 vaccination [13].

**DISCLAIMER**

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

**CONSENT**

As per international standard or university standard, respondents' written consent has been collected and preserved by the author(s).

**ETHICAL APPROVAL**

It is not applicable.

## ACKNOWLEDGEMENT

Authors acknowledge the support given by management and Principal, Bhavan's Viviekananda college to carry out the study and present the data.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Available: <https://www.mohfw.gov.in/pdf/COVID19VaccineOG111Chapter16.pdf> (Accessed on December 8th 2021).
2. Ministry of Health and Family Welfare (2020) COVID-19 Vaccine Operational Guidelines. Ministry of Health and Family Welfare, Government of India; 2020. Available: <https://main.mohfw.gov.in/sites/default/files/COVID19VaccineOG111Chapter16.pdf>. (Accessed on November 19th, 2021)
3. Available: <https://www.mohfw.gov.in/pdf/RevisedVaccinationGuidelines.pdf> (Accessed on November 15<sup>th</sup> 2021).
4. Available: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/covid-19-vaccines/advice> (Accessed on December 8th 2021).
5. Available: [https://www.mohfw.gov.in/covid\\_vaccination/vaccination/faqs.html#who-will-get-the-vaccine](https://www.mohfw.gov.in/covid_vaccination/vaccination/faqs.html#who-will-get-the-vaccine) (Accessed on December 8th 2021).
6. Available: <https://www.mohfw.gov.in/pdf/FAQCoWINforcitizens.pdf> (Accessed on November 15<sup>th</sup> 2021)
7. Kumar VM, Pandi-Perumal SR, Trakht I, Thyagarajan SP. Strategy for COVID-19 vaccination in India: the country with the second highest population and number of cases. *npj Vaccines*. 2021; 6(1):1-7.
8. Dasgupta R, Mishra P, Yadav. COVID-19 vaccination and the power of rumors: Why we must "Tune in". *Indian J Public Health*. 2021;65:206-8.
9. Hesitancy among Respondents; December; 2020. Available: <https://www.expresshealthcare.in/blogs/editors-blog/surveys-find-increasing-vaccine-hesitancy-among-respondents/426418/>.
10. Reuters. Vaccine 'Hesitancy' Rises among Indians as Virus Cases Fall – Survey; December; 2020. Available: <https://in.reuters.com/article/us-health-coronavirus-india-cases/vaccine-hesitancy-rises-among-indians-as-virus-cases-fall-survey-idINKBN28R0HD>.
11. Available: [https://www.google.com/search?q=what+is+the+vaccination+status+of+covid+19+in+india&rlz=1C1SQJL\\_enIN903IN903&oq=&aqs=chrome.69i59i450l8.4418l9720j0j15&sourceid=chrome&ie=UTF-8#wptab=s:H4sIAAAAAAAAAAOMwe8Q4g5Fb4OWPe8](https://www.google.com/search?q=what+is+the+vaccination+status+of+covid+19+in+india&rlz=1C1SQJL_enIN903IN903&oq=&aqs=chrome.69i59i450l8.4418l9720j0j15&sourceid=chrome&ie=UTF-8#wptab=s:H4sIAAAAAAAAAAOMwe8Q4g5Fb4OWPe8) (Accessed on November 14<sup>th</sup> 2021)
12. Available: <https://theprint.in/india/education/universities-colleges-push-for-vaccination-of-students-before-they-return-to-campus/684997/> (Accessed on November 14<sup>th</sup> 2021)
13. Available: <https://health.odisha.gov.in/pdf/Updates-recommendations-NEGVAC-for-Covid-19-vaccination-22052021.pdf> (Accessed on November 30th 2021)

© 2021 Padma and Babu; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/77739>