



Effect of Urine Sources on Some Soil Health Indicators, Maize Yield and Its Heavy Metals Uptake in Abakaliki, Southeastern Nigeria

J. N. Nwite^{1*}

¹*Department of Soil Science and Environmental Management, Faculty of Agriculture and Natural Resources Management, Ebonyi State University, P.M.B. 053, Abakaliki, Nigeria.*

Author's contribution

The sole author designed, analyzed and interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/IJPSS/2016/29016

Editor(s):

- (1) Radim Vacha, Deputy Director of Research and Development, Research Institute for Soil and Water Conservation, Czech Republic.
(2) Slawomir Borek, Faculty of Biology, Department of Plant Physiology, Adam Mickiewicz University, Poland.

Reviewers:

- (1) Makaya M. Joseph, Research and Training Unit, Life and Earth Sciences, University of Ouagadougou, Burkina Faso.
(2) Manuel Jimenez Aguilar, Agricultural and Development Research Institute, Granada, Spain.
(3) Olowoake Adebayo Abayomi, Kwara State University, Malete, Nigeria.
(4) Raghavendra Singh, ICAR – National Organic Farming Research Institute, Tadong, Gangtok, Sikkim – 737 102, India.
(5) Roshan Babu Ojha, Nepal Agricultural Research Council, Nepal.

Complete Peer review History: <http://www.sciencedomain.org/review-history/17407>

Original Research Article

Received 17th August 2016
Accepted 15th November 2016
Published 31st December 2016

ABSTRACT

The need to find alternative to inorganic fertilizer which is costly and not easily affordable to local farmers necessitates research in rare areas. Thus, this experiment was carried out at the Plant and Screen house to study effect of urine sources on some soil health indicators, maize yield and its heavy metals uptake. Completely Randomized Design was used in laying the experiment. 20 kg of sieved soil was treated with different urine sources replicated five times. The result indicates that soil pH, total N and organic matter were significantly ($P<0.05$) higher in different urine sources than control. Human urine had significantly ($P<0.05$) higher treatment effect on soil pH, percent total N and organic matter compared to other sources of urine. Similarly, human urine was 9-10%, 15-27%, and 10-47% higher in number of leaves, plant height, and grain yield of maize when compared to those of cattle and goat urine sources. Significantly ($P<0.05$) higher copper and lead uptake by maize grains were obtained in control relative to those of urine sources. Copper and lead uptake by maize grains were respectively higher by 20, 80, 87% and 87, 47, 7% in control when

*Corresponding author: E-mail: nwitejamesn@yahoo.com;

compared to human, cattle and goat urine sources. Generally, heavy metals uptake by maize grains is below recommended safe limits for toxicity. Urine from adult animals is recommended as credible alternative for improvement of soil health status and sustainable productivity.

Keywords: Alternative source; inorganic fertilizer; organic manure, toxicity; Zea mays L.

1. INTRODUCTION

Traditional agriculture relies heavily on mineral fertilizer NPK for crop production in Nigeria and other developing countries [1] and incidentally, use of fertilizer is confronted with problems of unavailability, high cost and increase in soil acidity. As a result, use of fertilizer is considered to be counterproductive and there is need for its alternative source. This alternative source is urine since it is easily affordable as it could be accessed from livestock and man. It has been reported by [2] that urine contains useful nutrients which if carefully harnessed could sustain soil health status and increase its productivity. Well preserved urine has good quality and could have the same effect as inorganic fertilizer in optimizing soil fertility status [1]. Research shows that urine contains major nutrients including nitrogen, phosphorus, potassium as well as calcium and magnesium which is dependent on age and feed of the animals [3].

When there is no planned disposal of urine it naturally constitutes health hazard due to its pungent odour which could be curtailed through its proper treatment and conversion in treating soil for higher productivity [1]. This offensive odour is attributed to freshly accumulated urine at pH of 6.7 [4]. [5-9] have shown that human urine source was successfully used as fertilizer in crop production and raising flowers in Europe and other countries. Confirmatory studies have been carried out using Barley and under crop and field trials or even under home gardening [8].

With the wide spread scarcity of inorganic fertilizer and its associated problems in food production, there is need for alternative source. If appropriate quantity of urine is applied to the soil at right time, its nitrogen contents could have the same value as that of inorganic fertilizer [2]. For instance, 100 kg N per hectare of urine improved Barley production between 90 – 110 days of planting in Sweden [8].

Naturally, human being could not easily accept food crops produced with urine due to suspicion

of its health hazard status and safe for consumption. This, however, could be overcome by treating urine for quality assurance and safe from health hazards [1]. In Nigeria food crops that grow around urinals or where urine is disposed are normally eaten by human beings and animals without any complaints of health problems. The objective of this experiment was to study effect of urine sources on some soil health indicators, maize yield and its heavy metals uptake under Abakaliki agroecological environment.

2. MATERIALS AND METHODS

2.1 Experimental Site

The research was conducted in 2015 at Plant and Screen House of Teaching and Research Farm, Faculty of Agriculture and Natural Resources Management, Ebonyi State University, Abakaliki. The area is located between latitude (06° 4' N and 08° 65' E) in the South-Eastern zone of Nigeria. The area experiences bimodal pattern of rainfall which is spread from April-July and September-November of each year. There is a break in August normally referred by residents as "August break". At the beginning of rainfall, it is torrential and violent and is characterized by thunderstorm and lightning. The minimum and maximum rainfalls are 1700 and 2000 mm with a mean of 1800 mm [10]. The temperature during rainy season is usually low (27°C) but increases to 31°C in dry season. Relative humidity is 80% in rainy season which declines to 60% during the cold Harmattan periods and dry season of the year [10] being characteristics of tropical climate.

The soil is derived from sedimentary deposits from cretaceous and tertiary periods. According to Federal Department of Agricultural Land Resources [11], Abakaliki agricultural zone lies within "Asu River" and is associated with Olive brown shale, fine grained sandstones and mudstone. It is unconsolidated within 1 m depth (Shale residuum) and belongs to the order ultisol classified as *typic haplustult*. The area was grown of short vegetation and medium to tall trees. There is also growth of native grasses, herbs and shrubs with patches of ground.

2.2 Experimental Design and Treatment Application

The study was carried out in 2015 between October and mid of January. The experimental design used in this study was Completely Randomized Design (CRD). Human urine of male adult was collected from prison inmates while cattle and goat urine was sourced from Cattle and Goats' market located at Gariki and Hossana, Abakaliki respectively. This was to reduce the time needed to collect enough quantity of urine for the study. Plastic containers of 5litres each were provided to both prison and Animal attendants for the purpose of collection of urine. The animals used were of matured age. The choice of these animals was based on ease of accessibility since every farming family in the locality can afford to keep them. These animals too are omnivorous and have common feeding habit at adult age. The urine was stored in air-tight plastic containers for 6 months before application to ensure sanitation process. The urine treatments were based on hectare equivalence of 20 kg soil.

Human urine = 50,000 mg ha⁻¹ equivalent to 100 mg kg⁻¹ soil
Cattle urine = 50000 mg ha⁻¹ equivalent to 100 mg kg⁻¹ soil
Goat urine = 50000 mg ha⁻¹ equivalent to 100 mg kg⁻¹ soil
Control = 0 mg ha⁻¹ equivalent to 0 mg kg⁻¹ soil

The urine rates were applied to 20 kg of soil weighed into perforated polybags two weeks after germination of maize seeds. These treatments were replicated six times to give a total of twenty four experimental polybags in the experiment. The polybags were watered to field capacity as often as moisture is required. The polybags were separated by 0.5 m spaces while replicates were set 1 m apart.

2.3 Planting of Maize

Maize variety (Oba super II hybrid) (*Zea mays* L.) collected from Ebonyi State Agricultural Development Programme (EBADEP), Onu Ebonyi Izzi, Abakaliki was used as a test crop. The maize seeds were planted at two seeds per hole and at 5 cm depth in each pot. Two weeks after germination (WAG), thinning was carried out to allow one plant per stand. Weeds were

removed by handpicking at regular intervals till harvest.

2.4 Agronomic Parameters

A total of ten maize plants were randomly selected and tagged for study. When the husks were dried, the cobs were harvested, dehusked, shelled and grain yield adjusted to 14% moisture content. Plant height was measured with metric ruler from the base of plant to tallest plant leaf at tasseling.

2.5 Soil Sampling

Auger sampler was used to collect soil samples at 0-20 cm depth from site where soil used for experiment was collected. The samples were bulked, air dried and passed through 2mm sieve and used for routine laboratory analysis. Samples were further collected from each polybag for some post-harvest chemical properties determination.

2.6 Laboratory Methods

Particle size distribution of the experimental soil was determined using the Bouyoucous method as outlined in [12] procedure. Soil pH determination was carried out in soil/water solution ratio of 1:2.5. The pH values were read off using pH meter with glass electrode [13]. Total nitrogen was determined using Microkjeldahl procedure [14]. Available phosphorus determination was done using Bray-2 method as outlined in [15]. Organic matter was determined by [16] digestion method. Exchangeable bases of calcium (Ca), Magnesium (Mg), Potassium (K), and Sodium (Na) were extracted using ammonium acetate (NH₄OAC) extraction method. Potassium and sodium were determined using flame photometer. The compositions of urine were determined by Atomic Absorption spectrophotometer as well as crop uptake copper (Cu) lead (Pb) using [17] procedure.

2.7 Data Analysis

Data collected from the experiment were subjected to Analysis of Variance (ANOVA). Means were separated using Fishers' Least Significant Difference (FLSD) as outlined in [18]. Significance was reported at 5% probability level.

3. RESULTS AND DISCUSSION

3.1 Composition of Urine

Table 1 shows some major nutrients and heavy metals composition of urine source. There were variations in values of nutrients and heavy metals in urine source. Nevertheless, human urine have highest values of nutrients when compared to livestock sources although, comparable. Cattle and goat urine contained 0.10 mgkg⁻¹ each of copper (Cu) and lead (Pb) but was not found in human urine. The comparable composition of elemental concentrations in animal urine could be attributed to their adult age, omnivorous nature as well as similarity in their dietary needs.

Table 1. Compositions of some major nutrients and heavy metals in urine sources

Parameter	Human urine	Cattle urine	Goat urine
pH H ₂ O	9.1	9.0	8.9
Ammonia mgkg ⁻¹	0.01	0.01	0.01
Nitrogen %	4.54	4.52	4.51
Phosphorus mgkg ⁻¹	0.04	0.03	0.02
Potassium cmolkg ⁻¹	0.05	0.03	0.03
Sodium cmolkg ⁻¹	0.29	0.28	0.28
Copper mgkg ⁻¹	-	0.10	-
Lead mgkg ⁻¹	-	-	0.10

3.2 Properties of Soil before Initiation of Study

Table 2 shows physicochemical properties of soil before initiation of study. Sand fraction was dominant in the particle size distribution. The textural class was sandy loam. The pH was 5.0 which indicates strongly acidic soil [19]. Nitrogen was 0.13% and according to [20] is very low and organic matter which had 2.2% value was moderate using Benchmark of [21] for Tropical soils. Phosphorus (20.40 mgkg⁻¹) was high [22]. Exchangeable calcium was of medium value but magnesium, potassium and sodium were very low [23]. Cation exchange capacity recorded very low values [23]. It implies that the soil was of low fertility status as obtained in Abakaliki areas for soils used for maize production as well as other crops.

3.3 Effect of Urine Sources on Some Soil Health Indicators

Effect of urine sources on some soil health indicators is shown in Table 3. Soil health

indicators are used in this study to indicate parameters that are determinant of soil fertility status. Urine sources had significantly ($P < 0.05$) higher treatment effect on pH when compared with the control. Human source of urine had significantly ($P < 0.05$) higher pH than those of cattle and goat urine sources, respectively. On the other hand, human urine was 5 and 6% higher in pH than the urine from cattle and goat. Similarly, significantly ($P < 0.05$) higher treatment effect was obtained in percent total nitrogen in human and cattle sources of urine relative to control. Furthermore, human urine showed significantly ($P < 0.05$) higher treatment effect on percent total nitrogen compared to those of cattle and goat sources of urine. Different sources of urine had significantly ($P < 0.05$) higher available phosphorus when compared to its corresponding value obtained in control. There were also significant differences in available phosphorus among the treatments. The available phosphorus of human source of urine was 14% higher than control and generally marginally higher than those of cattle and goat sources of urine. There was significantly ($P < 0.05$) higher treatment effect of urine sources on percent organic matter relative to control. Urine obtained from human and goat was significantly ($P < 0.05$) higher in percent organic matter than the one from cattle. This represents 21 and 14% increments in percent organic matter in human and goat sources of urine compared to that of cattle source.

Table 2. Properties of soil before initiation of study

Sand gkg ⁻¹	750
Silt gkg ⁻¹	140
Clay gkg ⁻¹	110
Texture class	Sandy loam
pH H ₂ O	5.0
Total Nitrogen %	0.13
Organic matter %	2.2
Available phosphorus mgkg ⁻¹	20.40
Calcium cmolkg ⁻¹	3.10
Magnesium cmolkg ⁻¹	0.92
Potassium cmolkg ⁻¹	0.17
Sodium cmolkg ⁻¹	0.10
Cation exchange capacity cmolkg ⁻¹	7.50

The significant increments of pH, percent nitrogen, organic matter and available phosphorus show that these soil health indicators were released into the soil by urine sources. This finding indicates that urine could substitute

Table 3. Effect of urine sources on some soil health indicators

Treatment	pH H ₂ O	Total N%	P mgkg ⁻¹	OM%
Control	5.1d	0.10b	25.60d	1.05d
Human urine	6.0a	0.14a	29.65a	1.76a
Cattle urine	5.7b	0.12b	28.24c	1.40a
Goat urine	5.6c	0.11b	28.30b	1.62b
FLSD (0.05)	0.1	0.02	0.1	0.05

P – Available phosphorus, *OM* % – Percent organic matter, *N* % -Percent Total nitrogen. Treatment means with different letters indicate significant differences ($P < 0.05$)

mineral inorganic fertilizer as it could be used as fertilizer to supply essential and major nutrients to soil on one hand and on the other improve soil health status. These findings are in line with the report of [2] that urine used as fertilizer improved soil health status. Several researchers [24-26] reported positive impact of urine on nitrogen which increased and sustained soil fertility. Higher significant positive effect of human urine source on soil health indicators suggests that it could be more superior to other urine sources in improving soil health indicators [1]. This finding had earlier been reported by [27,2]. This implies that urine sources especially human urine could serve as useful alternative fertilizer for crop production.

The positive impacts of the urine fertilizer on soil health indicators indicate an improvement on the soil health status. The human urine source improved the soil pH keeping it within a safe range of 5.6-6.0, nitrogen and organic matter at significant levels than other sources. [28] stated that organic fertilizers perform better with some crops. This is further supported by [26] findings that urine fertilizer improved some health indicators.

3.4 Effect of Urine Sources on Agronomic Yield of Maize

Table 4 shows effect of urine sources on agronomic yield of maize. Urine sources had significantly ($P < 0.05$) higher number of leaves, taller maize plants and grain yields of maize except yields of maize in urine source when compared with the control, respectively. Human urine source was 9-10%, 15%-27%, and 10-47% higher in these maize agronomic yields than those of cattle and goat urine sources. Furthermore, there were significant differences in number of leaves, height of maize and grain yield of maize among different sources of urine and control. The significant effect of urine sources on agronomic parameters of maize could be linked to inherent capability of the maize plant [27] and

nutrients supplied to the soil by urine sources (Table 3). Result further showed that urine sources could substitute inorganic fertilization that could sustain profitable maize production. The differences observed in agronomic yield of maize could be attributed to inherent capabilities of the different urine sources. The generally higher agronomic yield of maize in urine sources relative to control could be attributed to improved soil health status by urine treatment. This by implication underscores the usefulness of urine as credible alternative to inorganic fertilizer for sustenance of soil fertility status and in increased maize production that is safe for human consumption. Human urine increased agronomic yield of maize due to its high nutrients (Table 1) and its ability to release same into the soil. [27] and [2] pointed out that human urine increased soil fertility and *Jathropha* production. Low grain yield of maize obtained in this study could be attributed to drought and high incidence of disease attack. This affected optimal watering of the polybags to field capacity moisture contents as there was scarcity of water. This caused moisture stress which coincided with flowering and ear formation of maize plants. There were shriveling of cobs with their consequent poor grain filling.

3.5 Effect of Urine Sources on Heavy Metals Uptake by Maize Grains

Effect of urine sources on heavy metals uptake by maize grains is shown in Table 5. The result showed significantly ($P < 0.05$) higher effect of copper and lead uptake by maize grains in control compared to urine sources of human, cattle and goat. This accounted for 20, 80 and 37% and 87%, 47% and 7% increments of copper and lead uptake in maize grains in control respectively compared to human, cattle and goat sources of urine. There were significant differences in copper and lead uptake in maize grains among the treatments except copper in human and goat urine sources.

Table 4. Effect of urine sources on agronomic yield of maize

Treatment	No. of leaf	Plant height (cm)	Grain yield (g/pot)
Control	11.2a	58.12d	2.0c
Human urine	13.4b	85.08a	4.2a
Cattle urine	12.2c	72.14b	3.8b
Goat urine	12.0c	61.98c	2.2c
FLSD (0.05)	0.3	0.1	0.4

Treatment means with different letters indicate significant differences ($P < 0.05$)

Table 5. Effect of urine sources on heavy metals uptake by maize grains

Treatment	Cu mgkg ⁻¹	Pb mgkg ⁻¹
Control	0.30a	0.30a
Human urine	0.04c	0.04d
Cattle urine	0.24b	0.16b
Goat urine	0.06c	0.28c
FLSD (0.05)	0.05	0.08

Cu – Copper, Pb – Lead, Treatment means with different letters indicate significant differences from each other ($P < 0.05$)

The significantly higher copper and lead uptake by maize grains grown in control plot compared to those grown in urine sources treated plots could be attributed to inputs from soil rather than urine fertilizer. Analysis of urine sources indicated very low presence of copper and lead (Table 1). These findings show that urine could be used as fertilizer for crop production without placing man at a risk of ecotoxicity of heavy metals. This finding could be attributed to improved health status of soil (Table 3) and low presence of heavy metals in urine sources. The likelihood of heavy metals to build up in soil amended with urine fertilizer appears to be higher in soils treated with goat and cattle urine sources than human urine. Comparatively, lead has the propensity to build up faster than copper due to urine contamination of soil. [29] reported heavy metal uptake by crops in their work and noted that these heavy metals were stored in crop parts. [30] and corroborated by [31] in their findings which observed that human beings were at risk of heavy metals toxicity if they could utilize crops grown around areas contaminated with heavy metals due to eco-toxicity. This could be possible through recycling of heavy metals through food chain. Heavy metal of lead has the capacity to cause brain, renal or reproductive disorders in human beings. The heavy metals of copper and lead are below 0.0-2.0 and 0.01

rated as medium to low [32] values and far below 0.0-5.0 [33] or 2-1500 and 2-300 recommended as normal by [34]. However, the interesting result is that heavy metal uptake by maize grains could not be linked to urine treatment of soil.

4. CONCLUSION

This study has shown that urine sources could improve soil health status and serve as useful alternative fertilizer for maize crop production. Urine sources significantly improved soil health indicators. Agronomic parameters responded significantly to improved soil health status and were superior in urine sources than control. Perhaps, the greatest beneficial aspect of use of urine as fertilizer is low input of heavy metals which keep them below safe limits and without any danger of eco-toxicity. In view of its superior performance over other urine sources, human urine could be harvested for treatment of soil for higher and sustainable productivity rather than be allowed to be wasted through improper disposal. However, there is need to apply caution and monitor unrestricted use of urine sources for soil fertility improvement and crop production.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. Nwite JN. Effect of urine source on soil chemical properties and maize yield in Abakaliki, South-Eastern Nigeria. International Journal of Advance Agricultural Research. 2015;3:31-36.
2. Adeoluwa OO, Suleiman ON. Effect of human urine on the growth performances of *Jathropha curcas* seedlings and some soil health indices. Nigerian Journal of Soil Science. 2012;22(2):186-193.
3. Marino C. Urine composition depends on certain factors. Journal of Soil Science Society of America. 2008;73:159-219.
4. Høglund C. Evaluation of microbial health risks associated with the reuse of source-separated human urine. Ph. D Thesis, Department of Biotechnology, Applied Microbiology, Royal Institute of Technology (KTH) Stockholm Sweden. 2001;62-64.
5. Heinnonen-Tanski, Van Wijk-Sibesma C. Human excreta for plant production.

- Bioresource Technology. 2005;96:403-411.
6. Kirchmann H, Peterson S. Human urine-chemical composition and fertilizer use efficiency. *Fertilizer Resources*. 1995; 40:149-154.
 7. Steineck S, Richert Stintzing A, Rodhe L, Elinquist H, Jakobsson MC. Plant nutrients in human and food refuse. *In: Proceedings of Nigerian Journal of Forestry Seminar 292*. DIAS Report. 1999;13:125-130.
 8. Richert Stintzing A, Rodhe L, Akerhieln H, Steniecek S. Human urine as a fertilizer and plant nutrients application technique and environmental effects. *In: Venglosk, J., Greserova, G. (eds). Proceedings of 10 International Conference Ramiran 2002 Network*. FAO European System of Cooperative Research Network. 2002;161-162.
 9. Malkki S, Heinonen-Tanski. Composition of toilets in permanent houses. *In: 1 Use of municipal organic wastes*. Proceedings of Nigerian Journal of Forestry Seminar 292 DIAS-report. 1999;13:147-154.
 10. Overseas Development of Natural Resources Institute (ODNR). Nigeria Profile of Agricultural Potential ODA, United Kingdom. 1989;3.
 11. Federal Department of Agricultural Land Resources (FDALR). Reconnaissance soil survey of Anambra State, Nigeria. Soil Report, Kaduna. 1985;3.
 12. Gee GW, Or D. Particle size analysis. *In: Dane. J. H. and Topp, G. C. (eds). Methods of Soil Analysis. Physical Methods*. Soil Science Society America. 2002;5(4):255-293.
 13. Peech M. Hydrogen activity. Methods of soil analysis. *In: Black CA (ed). American Society of Agronomy*. 1965;9(1):914-926.
 14. Bremner JM. Nitrogen-total. *In: Sparks, D. L. (ed). Methods of soil analysis. Chemical Methods*. American Society of Agronomy. 1996;5(3):1085-1121.
 15. Page AL, Miller RH, Keeney DR. Methods of soil analysis. American Society of Agronomy. 1982;9:539-579.
 16. Walkey A, Black A. An examination of the degtjareff method for determining soil organic matter and a proposed modification of the chronic acid titration method. *Soil Science*. 1934;37:29-38.
 17. Dewis J, Freitas F. Physical and chemical methods of soil and water analysis. *Soil Bulletin 18*, FAO, UN, Rome; 1976.
 18. Steel GD, Torrie JH. Procedures of statistics. A biometrical approach, 2nd ed. New York, McGraw Hill, Book Company. 1980;63.
 19. Landon JR. (eds). Booker, tropical soil manual. A hand book for soil survey and agricultural land evaluation in tropics and subtropics New York, USA, John Wiley and Sons: Inc. Third Avenue; 1991.
 20. Enwezor WO, Udo FJ, Sobulo RA. Fertility status and productivity of acid sands: *In: Acid sands of Southeastern Nigeria*. Soil Science Society of Nigeria. 1981;1:56-73.
 21. Federal Ministry of Agriculture, Water and Rural Development (FMAWRD). Fertilizer use and management practice for crops in Nigeria. Produced by the Federal Fertilizer Department. *In: Aduayi, E. A., Chude, V. O. Adebusi, B. A. and Olayiwola, S. O. (eds), Abuja*. 2002;188.
 22. Enwezor WO, Udo EJ, Usoroh NJ, Ayoade KA, Adepetu JA, Chude VO, Udegbe CJ. Fertilizer use and management practices for crops in Nigeria, Series No 2. Fertilizer Procurement and Distribution Division, Federal Ministry of Agriculture, Water Resources and Rural Development, Lagos, Nigeria; 1989.
 23. Asadu CLA, Ucheonye Oliobi C, Agada C. Assessment of sewage application in southeastern Nigeria. Part 1 Impact on selected soil morphological and physical properties. *Outlook on Agriculture*. 2008; 37(1):57-62.
 24. Gutser R, Ebertseder T, Weber A, Schrami M, Schmmidhlter U. Short-term and residual availability of nitrogen after long-time application of organic fertilizers on arable land. *Journal of Plant Nutrition and Soil Science* 2005;168:439-446.
 25. Schonning C. Urine diversion-hygienic risks and microbial guidelines for reuse. Department of Parasitology, Mycology and Environmental Microbiology. Swedish Institute for Infections; 2001.
 26. Adeluwa OO, Cofie O. Urine as an alternative fertilizer in agriculture: *In: Effects on amaranths (Amaranthus caudatus) production*. Renewable Agriculture and Food Systems. 2012;8:1. DOI: 10.1017/51742170511000512
 27. Bengé M. Assessment of the potential of *Jathropa curcas*, (*biodiesel*) for the energy production and other uses in developing countries. Posted on EC Ho's website with permission of the author. 2006;2:22.

- Available:<http://www.ascension-publishing.com/B/Z/Jathropha>
(Pdf download on January 9, 2012)
28. Perverly JH, Gates PB. Utilization of municipal solid waste and sludge compost in crop production systems. *In: Sewage sludge, Land utilization and the Environment. Proceedings of a conference by American Society of Agronomy*; 1993.
 29. Adewole MB, Adeoye GO, Sridhar MKC. Effect of inorganic and organo mineral fertilizers on the uptake of selected heavy metals by *Helianthus annuus L* and *Tithornia diversifolia* (Hems h) under green house condition. *Journal of Toxicological and Environmental. Chemistry*. 2008; 91(5):970-980.
 30. Anikwe MAN, Nwobodo KCA. Long-term effect of municipal wastes disposal on soil properties and productivity in sites used for urban agriculture in Abakaliki, Nigeria. *Bio-resources Technology*. 2002; 83:24-50.
 31. Asadu CLA, Nweke FI. Soils of arable crop fields in sub-Sahara Africa: Focus on cassava growing areas, collaborative study of cassava in Africa. Working Paper No. 18. Resources and Crop -Management Division, IITA, Ibadan, Nigeria. 1999; 1782.
 32. Lagos State Environmental Protection Agency (LASEPA). Lagos State Environmental Reports. 2005;3:28-33.
 33. World Health Organization (WHO). Guideline for drinking quality water. Switzerland. 1996;1(2&3):50-57.
 34. Alloway BJ. Heavy metals on soils. New York: John Wiley and Sons, Inc. 1990; 280.

© 2016 Nwite; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://sciencedomain.org/review-history/17407>